List of Drugs approved

List of Drugs approved during 2006 to 2021

| Sl. No. | Name of Drug | Pharmacological Classification | Date of Approval |
|---------|--|---|------------------|
| 1. | Oxandrolone Tablet (2.5/10 mg) | To promote weight gain | 4-8-06 |
| 2. | Tenofovir Disoproxil Fumarate 300 mg + Emtricitabin 200 mg + Efavirenz 600 mg tablet | For HIV-1 infection | 1-8-06 |
| 3. | Lumiracoxib Tablet (100/200/400 mg) | For OA, RA acute pain & primary dysmenorrhoea | 1-8-06 |
| 4. | Quetiapine S.R tablet 200/300 mg | For psychotic disorder | 1-8-06 |
| 5. | Amisulpiride Tablet (50/100/200 mg) | For schizophrenic disorder | 1-8-06 |
| 6. | Baclofen E.R capsule (30 mg) | For spasticity | 1-8-06 |
| 7. | Forskolin Eye drop (1%) | For open angle glaucoma | 27-7-06 |
| 8. | Sodium Hyaluronate Injection (1%) | For OA | 26-7-06 |
| 9. | Clobestasol Propionate (0.05%) + Calcipotriol Ointment (0.005%). | For plaque psoriasis | 20-7-06 |
| 10. | ISMN SR 30 mg + Aspirin 75/150 mg tablet (additional strength). | Same as approved | 20-7-06 |
| 11. | Cefixime 200 mg + Pot. Clavulanate (125 mg) tablet | Antibiotic | 20-7-06 |
| 12. | Levocetrizine 2.5/5 mg + Ambroxol 30 mg/5ml of syrup | For productive cough | 20-7-06 |
| 13. | Lamivudine 40 mg + Stavudine 10 mg dispersible tablet | For HIV infection | 12-7-06 |
| 14. | Vitamin C 500 mcg + Zinc Citrate 2.2 mg + Selenium 60 mcg tablet | For Vitamin C deficiency | 12-7-06 |
| 15. | Deferasirox dispersible tablet 125/250/500 mg | For chronic iron overload due to blood transfusion | 10-7-06 |
| 16. | Zolpidem Tartrate E.R tablet (6.25 mg/12.5 mg) | For insomnia | 10-7-06 |
| 17. | Solifenacin Succinate Tablet (5 mg/10 mg) | For overactive bladder with urge urinary incontinence urgency & urinary frequency | 6-7-06 |
| 18. | Mebeverine HCl 135 mg + Isabgula husk 3.5 gm sachet | Irritable bowel syndrome | 6-7-06 |
| 19. | Bimatoprost 300 mg + Timolol (as maleate) 5 mg/ml eye drops | For reduction of IOP | 5-7-06 |
| 20. | Ceftriaxone (as sodium) 250 mg + Tazobactam 31.25 mg for injection (additional strength) | Antibiotic | 5-7-06 |
| 21. | Buclizine HCl tablet (25 mg) & Syrup 6 mg/5 ml | As appetite stimulant | 28-6-06 |
| 22. | Olmesartan 40 mg + HCTZ 25 mg tablet (additional strength) | Anti-hypertensive | 28-6-06 |
| 23. | Levofloxacin 0.5% Ear drops | For otitis media | 27-6-06 |
| 24. | Aspirin 150 mg + Clopidogrel (as bisulphate) 300 mg tablet (additional strength) | As loading dose for prevention of in-stent restenosis in PCI | 22-6-06 |
| 25. | Methyl Prednisolone Aceponate Cream (0.1%) | For atopic dermatitis | 15-6-06 |
| 26. | Ferrous ascorbate (equivalent to 50 mg Iron) + Folic acid 1.5 mg | For iron deficiency anaemia | 8-6-06 |
| 27. | Olopatadine Ophthalmic solution (0.1%) | For allergic conjunctivitis | 8-6-06 |

| 28. | Lopinavir 200 mg + Ritonavir 50 mg Tablet (additional strength) | Anti-HIV | 8-6-06 |
|-----|--|---|---------|
| 29. | Amoxycillin 1000 mg + Clavulanate Potassium 62.5 mg E.R tablet | Antibiotic | 8-6-06 |
| 30. | Lamotrigin 200 mg tablet (additional strength) dispersible tablet (5 mg/50 mg/100 mg/200 mg) | Anti-epileptic | 8-6-06 |
| 31. | Icodextrin 7.5% peritoneal dialysis solution | For peritoneal dialysis | 8-6-06 |
| 32. | Bemiparin Sodium Solution for injection (5000/7500/10,000 I.U) | For deep vein thrombosis | 8-6-06 |
| 33. | Nadifloxacin (1%) + Miconazole Nitrate (2%) + Mometasone Furoate (0.1%) Cream | For dermatoses | 6-6-06 |
| 34. | Amlodipine (as besylate) 5 mg + Metoprolol Succinate equivalent to Metoprolol Tartrate 50 mg | For hypertension | 5-6-06 |
| 35. | Gatifloxacin (0.3%) + Prednisolone Acetate (1%) eye drops | For Ophthalmic use | 2-6-06 |
| 36. | Cyclosporin Eye drops (0.1%) (additional strength) | For Ophthalmic use | 2-6-06 |
| 37. | Perflutren liquid microsphere injectable suspension | For diagnostic use | 31-5-06 |
| 38. | Zonisamide capsule 25/50 mg (additional strength) | Same as approved | 25-5-06 |
| 39. | Ibandronic Acid Tablet 150 mg (additional strength) | Same as approved | 25-5-06 |
| 40. | Levocetrizine 5 mg + Ambroxol Hydrochloride SR (75 mg) tablet | For production cough | 24-5-06 |
| 41. | Nevibolol (as HCl) 5 mg + Amlodipine (as besylate) 5 mg/10 mg | For hypertension | 24-5-06 |
| 42. | Itopride 150 mg capsule (S.R) | For GERD | 22-5-06 |
| 43. | Ciclesonide rotacaps (200/400mcg) | For asthma | 19-5-06 |
| 44. | Oxcarbazepine S.R 450 mg/900 mg (additional strength). | Anti-epileptic | 19-5-06 |
| 45. | Pregabalin Capsule 25/50/100/200 mg (additional strength) | Same as approved. | 19-5-06 |
| 46. | Olmesartan (as Medoxomil) 10 mg (additional strength) | Anti-hypertensive | 18-5-06 |
| 47. | Midazolam (as maleate) 7.5 mg/15 mg tablet | For insomnia | 10-5-06 |
| 48. | Formoterol Fumarate (6mcg) + Fluticasone Propionate (100/250/500mcg) dry powder inhaler | For asthma & COPD | 10-5-06 |
| 49. | Nadifloxacin Gel 1% | Acne vulgaris | 10-5-06 |
| 50. | Amlodipine (as besylate) 5 mg + HCTZ 12.5 mg Tablet | For hypertension | 24-4-06 |
| 51. | Nicorandil E.R Tablet (10 mg/20 mg) | Angina Pectoris | 24-4-06 |
| 52. | Atazanavir (100/150/200 mg) | Anti-HIV in adults. | 24-4-06 |
| 53. | Oseltamivir formulation (additional indication) | For Prophylaxis of influenzae in adult and children > 14 years of age | 24-4-06 |
| 54. | Pantoprazole (as Sodium) (E.C) 40 mg + Itopride (150 mg) S.R | For GERD | 8-4-06 |
| 55. | D-trans Fentantyl Transdermal patches | For pain | 5-4-06 |
| 56. | Resperidone Tablet & Syrup (additional Ind) | Acute mania & mixed episode in bipolar-I disorder. | 4-4-06 |
| 57. | Doxofylline Syrup (20 mg/ml) | For bronchial asthma & COPD. | 30-3-06 |
| 58. | Cholestyramine Powder for oral suspension (4gm/sachet) | For hypercholesterolaemia | 30-3-06 |
| 59. | Haloperidol Tablet (0.5 mg) (additional Lower strength) | As approved earlier. | 30-3-06 |
| 60. | Lactulose Enema (20%) | For constipation & hepatic encepthalopathy | 28-3-06 |
| 61. | Tobramycin Inhalation solution (300 mg/5ml) | For chronic pulmonary infection cystic fibrosis. | 22-3-06 |

| 62. | Propionyl-L-Carnitine HCl Tablet (500 mg) & injection 300 mg/vial | Chronic congestive heart failure | 22-3-06 |
|-----|--|--|---------|
| 63. | Eflornithine HCl Cream (139mg/gm) | For reduction of unwanted facial hair in women. | 22-3-06 |
| 64. | Saquinavir (500 mg) tablet (additional strength) | Anti-AIDS | 20-3-06 |
| 65. | Sodium Hyaluronate Cream 0.2% | For leg ulcer, diabetic ulcer etc. | 16-3-06 |
| 66. | Cefditeron (as pivoxil) 200 mg tablet | Antibiotic | 3-3-06 |
| 67. | Fluorescein Sodium Solution for Injection (100 mg/ml) | Ophthalmological diagnostic agent | 2-3-06 |
| 68. | Diclofenac Sodium (75/50 mg) as (E.C) + Misoprostol 200mcg. | For OA & RA | 28-2-06 |
| 69. | Atomoxetine 60 mg capsule (Addl. Strength) | For ADHD | 27-2-06 |
| 70. | Levetiracetam Oral Solution (100 mg/ml). | Anti-epileptic | 24-2-06 |
| 71. | Tenofovir Disoproxil Fumarate (300 mg) + Emtricitabine (200 mg) tablet | For HIV-1 infection | 23-2-06 |
| 72. | Aceclofenac 200 mg (S.R) tablet | NSAID | 22-2-06 |
| 73. | Desflurane liquid for inhalation (240 mg/bottle). | For induction & maintenance of anaesthesia. | 22-2-06 |
| 74. | Voriconazole Powder for oral solution (40 mg/ml) | Anti-fungal | 21-2-06 |
| 75. | Cefetamet Pivoxil HCl oral suspension 250 mg/ml | Antibiotic | 16-2-06 |
| 76. | Testosterone Transdermal Spray (12.5 mg/spray) (additional Lower strength) | For replacement therapy in males with Testosterone deficiency. | 15-2-06 |
| 77. | Zanamavir inhalation powder 5 mg/blister | For influenza | 13-2-06 |
| 78. | Doxofylline Tablet (400 mg) | For bronchial asthma & COPD. | 8-2-06 |
| 79. | Metoprolol Succinate equiv. to Metoprolol Tartrate 50 mg + Amlodipine (as besylate) 5 mg | For hypertension | 7-2-06 |
| 80. | Ciclesonide Inhaler 80/160mcg per actuation | For persistent asthma | 6-2-06 |
| 81. | Pegaptinib Sodium Injection (0.3mg). | For neovascular (wet) age related macular degeneration. | 6-2-06 |
| 82. | Dexibuprofen Tablet (200 mg/300 mg/400 mg) | For OA, RA & ankylosing spondylitis | 6-2-06 |
| 83. | Moxifloxacin Eye Ointment (5 mg/gm) | For ophthalmic use | 6-2-06 |
| 84. | Erdosteine Powder for oral suspension 35 mg/ml | For chronic obstructive bronchitis | 6-2-06 |
| 85. | Tamsulosin HCl (MR) 0.4mg + Dutasteride 0.5 mg tablet | For BPH | 6-2-06 |
| 86. | Atenolol 50 mg + HCTZ 25 mg (additional strength) | For hypertension | 3-2-06 |
| 87. | Alendronate Sodium (equiv. to Alendronic acid 70 mg) + Cholecalciferol 70mcg (equiv. to 2800 I.U Vit.D) Tablet | For osteoporosis | 01-2-06 |
| 88. | Combikit of Tamsulosin HCl 0.4mg M.R capsule (10 no.) & Dutasteride Soft Gelatin Capsule 0.5 mg (10 no.) | For BPH | 19-1-06 |
| 89. | Nebivolol (as HCl) 5 mg + Valsartan 80 mg | For hypertension | 17-1-06 |
| 90. | Cefprozil Powder for oral suspension (125 mg/5ml) | Antibiotic | 6-1-06 |
| 91. | Erdosteine 150 mg + Guaiphenesin 50 mg + Terbutaline 1.25 mg per 5ml syrup | Mucolytic | 8-8-06 |
| 92. | Rosiglitazone (as maleate) 2mg + Glibenclamide 5 mg + MetFormin 500 mg tablet | Anti Diabetic | 8-8-06 |
| 93. | L - Lysine HCl Ointment (150 mg/gm) | For wounds of various etiology | 8-8-06 |
| 94. | Levocitirizine 2 HCl 5 mg+ Ambroxol HCl SR 75 mg + Pseudoephedrine HCl 120 mg SR Capsules | For SAR, PAR & Chronic idiopathic urticatia | 9-8-06 |
| | | | |

| 96. | Dexibuprofen 200 mg/300 mg + Paracetamol 500 mg tablet | For RA, OA & ankolysing spondylosis | 10-8-06 |
|------|---|--|---------|
| 97. | ISMN SR 60 mg + Metoprolol Succinate ER 25/50 mg | For angina & Hypertension | 10-8-06 |
| 98. | Aspirin 75 mg/150 mg + Glycine 37.5 mg/75 mg tablet | For prevention of acute coronary syndrome in patients with CAD prophy-laxis against stroke | 10-8-06 |
| 99. | Combipack of 21 tablet of Quinin sulphate 600 mg each & 8 capsule of Doxycycline (as HCl) 100 mg each | For P. Falciparam malaria | 10-8-06 |
| 100. | Pravastatin 40 mg tablet (Additional strength) | As approved | 10-8-06 |
| 101. | Olanzapine powder for oral suspension (10 mg/5ml) | For schizophrenia in adults | 11-8-06 |
| 102. | Cefepem (as HCl) 0.5/1gm + Amikacin (as sulphate) 0.125/ 0.250gm Inj. | Antibiotic | 11-8-06 |
| 103. | Lamivudine 20 mg + Nevirapine 35 mg + Stavudine 5 mg tablet | Anti HIV | 11-8-06 |
| 104. | Nadifloxacin (10 mg/gm) + Mometasone Furoate (1mg/gm) cream | For mixed infection of skin | 17-8-06 |
| 105. | Premetrexed (Disodium) powder for solution for Inj. 500 mg/vial | Anti Cancer | 21-8-06 |
| 106. | Gliclazide 40 mg/80 mg + MetFormin HCl 500 mg + Rosiglitazone 2mg tablet | Anti-diabetic | 22-8-06 |
| 107. | Tiropramide HCl (100 mg) tablet | For abdominal spasmodic pain, colics of hepatobilliary track, abdominal colics | 22-8-06 |
| 108. | Eperisone HCl (50 mg) tablet | For neck-shoulder orm syndrome scapula hum oral per arthritis and low back pain | 23-8-06 |
| 109. | Caroverine cap. 20 mg & Inj. 40 mg/2ml | For abdominal pain & tinnitus in adult | 23-8-06 |
| 110. | Paclitaxel nanoparticle Inj. (20 mg/ml) | For metastatic breast cancer | 23-8-06 |
| 111. | Amoxycillin 250 mg + Dicloxacillin 250 mg cap. | Antibiotic | 28-8-06 |
| 112. | Rifaximine 200 mg tab. | For infections diarrhoea in adults | 28-8-06 |
| 113. | Tigecycline powder For Inj. 50 mg/vial | For skin & skin structure infection & intra abdominal infection | 28-8-06 |
| 114. | Stavudine SR 100 mg tablet | Anti HIV | 29-8-06 |
| 115. | Cefixime (50 mg/100 mg) + Clavulanic acid 31.125 mg/62.5 mg per ml dry syrup & tablet | Antibiotic | 29-8-06 |
| 116. | Ciclsonide 80 mcg/160 mcg + Formotorol 4.5 mg inhaler | Anti asthratic | 29-8-06 |
| 117. | Ciclsonide 100 mcg/ 200 mcg/400 mcg + Formotorol Fumarate 6 mg rotacaps | Anti asthmatic | 29-8-06 |
| 118. | Abacavir Sulphate syrup 20 mg/ml | For HIV Infection | 29-8-06 |
| 119. | S(-) pantoprazole (as sodium) 20 mg (E.C.) + Domperidome SR 30 mg tablet | For GERD | 30-8-06 |
| 120. | Cefadroxil 500 mg + Clavulanic acid 125 mg tablet | Antibiotic | 30-8-06 |
| 121. | Cefadroxil 250 mg + Clavulanic acid 62.5 mg dispersible tablet | Antibiotic | 30-8-06 |
| 122. | Cefadroxil 250 mg + Clavulanic acid 62.5 mg per 5ml syrup | Antibiotic | 30-8-06 |
| 123. | Human Insulin (r-DNA) Powder inhaler | For Type-I & Type-2 Diabetes | 30-8-06 |
| 124. | Fulvestrant Inj. 250 mg/5ml pre-filled syringe | Anti Cancer | 30-8-06 |
| 125. | S(-) Pantoprazole (as Sodium) Inj. 20 mg/vial | For pepticulcer, GERD & Zollinger Ellison Syndrome | 30-8-06 |
| 126. | Dasatinib tablet 20 mg/50 mg/70 mg tablet | For chronic Hepatitis B infection | 30-8-06 |
| 127. | Telbivudine 600 mg tablet | For chronic Hepatitis B infection | 30-8-06 |
| 128. | Aprepitant cap (80 mg/125 mg) | For chemotherapy induced nausea & vomiting | 30-8-06 |

| 129. | Telmisartan 40 mg + Amlodipine (as Besylate) 5 mg tablet | For essential hypertension | 30-8-06 |
|------|---|---|----------|
| 130. | Gemifloxacin (as msylate) 320 mg tablet | Antibiotic | 30-8-06 |
| 131. | Ceftazidime 500 mg/1000 mg + Tazobactum (as sodium) 62.5 mg/125 mg per vial for injection | Antibiotic | 30-8-06 |
| 132. | Cefepime (As HCl) 500 mg/1000 mg + Tazobactum (as Sodium) 62.5 mg/125 mg | Antibiotic | 30-8-06 |
| 133. | Paromomycin Inj. 375 mg/ml | For visceral leishma-niasis (Kalaazar) | 30-8-06 |
| 134. | Cefixime 100 mg + Cloxacillin (as Sodium) 500 mg + Lactobacillus (45 million spore) tab. | Antibiotic | 31-8-06 |
| 135. | Alprazolam Dispersible Tab. 0.125 mg/0.25 mg/0.50 mg | For anxiety | 20-9-06 |
| 136. | Epinastine HCl (0.05%) Opthlamic solution | For allergic conjunc-tivitis | 22-9-06 |
| 137. | Levofloxacin tablet (250 mg/500 mg) & Infusion 5 mg/ml (For additional indication) | For prostatitis | 26-9-06 |
| 138. | Feracrylum 1% mouth gargle 10 mg/ml | As haemostatic agent in dental surgery | 28-9-06 |
| 139. | Aceclofenac Inj. (150 mg/ml) | For pain in post operative & post traumatic cases | 28-9-06 |
| 140. | Combipack of one cap. of aprepitant .125 mg & two cap. of aprepitant 80 mg each (combipack) | As Approved | 10-10-06 |
| 141. | Formoterol fumarate 6mcg + Fluficasone propinate 50 mcg/125 mcg/250mcg MDI | For Asthma & COPD | 14-10-06 |
| 142. | Amlodipine 25 mg + Metoprolol 25 mg (additional lower strength) | | 01-11-06 |
| 143. | Estrogen vaginal tablet 25 mcg | For atropic viginitis | 2-11-06 |
| 144. | Esomeprazole 40 mg (E.C.) + Itopride 150 mg SR Capsules | For GERD | 13-11-06 |
| 145. | Cefixime 200 mg + Cloxacillin 500 mg (ER) + Lactobacillus 90 million Spore tab. (additional strength) | Antibiotic | 14-11-06 |
| 146. | Cefixime dispersible tablet 400 mg (new dosage form) | Antibiotic | 22-11-06 |
| 147. | Trandalopril 1mg/2mg/4mg/2mg + Verapamil (ER) 240 mg/240 mg/240 mg/180 mg | Anti hypertensive | 23-11-06 |
| 148. | Ciclesonide 160mcg/320mcg cap. for inhalation | Anti asthmatic | 28-11-06 |
| 149. | Rabeprazole 20 mg (EC) + Itopride SR 150 mg | For GERD | 28-11-06 |
| 150. | Metoprolol succinate 25 mg/50 mg ER + HCTZ 12.5 mg | Anti hypertensive | 30-11-06 |
| 151. | Atorvastatin 5 mg + Fenofibrate 160 mg (additional strength) | As approved | 01-12-06 |
| 152. | Aceclofenac Inj. 50 mg/ml (additional strength) | As approved | 01-12-06 |
| 153. | Amisulpride 400 mg tablet (additional strength) | As approved | 01-12-06 |
| 154. | Misoprostol tablet 25 mcg/100 mcg/ 200 mcg (additional strength indication) | For cervical ripening, prevention of post partum haemorrhage & First trimester of aboration with mifepristone | 8-12-06 |
| 155. | Atazanavir 150 mg + Ritonavir 50 mg capsules | Anti HIV | 12-12-06 |
| 156. | Mebeverine HCl 135 mg + Chlordiazepoxide 5 mg tablet | For IBS | 12-12-06 |
| 157. | Ampicillin 250 mg + Dicloxacillin 250 mg tablet | Antibiotic | 13-12-06 |
| 158. | Etoricoxib 10 mg + Methyl Salicylate 100 mg + Mentol 50 mg + Linseedoril 30 mg per gm. of Gel | For acute Musculoskeletal pain | 14-12-06 |
| 159. | Racecadotril dispersible tablet (10 mg/30 mg) (new dosage Form) | For diarrhoea | 15-12-06 |
| | | | |

| 161. | Aceclofenac dispersible tablet 100 mg (new dosage form) | as Approved | 21-12-06 |
|------|---|-------------|----------|
|------|---|-------------|----------|

List of Drugs approved during January to December 2007

| Sl. No. | Drug Name | Indication | Date of Approval |
|---------|--|--|------------------|
| 1. | R-Ondansetron (As HCl Dihydrate) (1 mg/ml) injection | Chemotherapy induced nausea & vomiting | 12-1-07 |
| 2. | Torsemide 10 mg + Amiloride 5mg tablets | Edema associated with congestive heart failure & hypertension | 17-1-07 |
| 3. | Fexofenadine Suspension (30 mg/5 ml) | Seasonal allergic rhinitis & chronic idiopathic urticaria | 19-1-07 |
| 4. | Sildenafil Citrate 20 mg Tablet | For pulmonary osterial hypertension | 23-1-07 |
| 5. | Pioglitazone 15 mg/30 mg + MetFormin ER 1000 mg tablet (Additional Strength) | As approved | 23-1-07 |
| 6. | FluvAstatin Tablet (As calcium) 10 mg/20 mg/40 mg (Additional Strength) | As approved | 29-1-07 |
| 7. | Halobetazol Propionate Cream (0.05%) | Corticosteroid responsive dermatomes | 2-2-07 |
| 8. | Eszopiclone tablet (1 mg/2 mg/ 3 mg) | For insomnia | 7-2-07 |
| 9. | Diltiazem HCl 2% gel | For anal fissure | 7-2-07 |
| 10. | Nalbuphine injection 10 mg/20 mg per ml | For moderate to severe pain | 7-2-07 |
| 11. | Levonorgestrel 1.5 mg tablets | For emergency contraceptive | 7-2-07 |
| 12. | Pantaprazole Sodium 20 mg/40 mg Tablet (with Sodium Bicarbonate As buffer) | For gastric ulcer, duodenal ulcer, Z. E Syndrome, GERD | 7-2-07 |
| 13. | Levofloxacin (250 mg/500 mg) + Ambroxol SR 75 mg tablet | For lower respiratory tract infection in adult | 8-2-07 |
| 14. | Testosterone Cypionate (100 mg/200 mg) + Cotton seed oil 736/566 mg per ml injection | For testosterone deficiency (i) Primary hypogonadism (ii) Hypogonadotropine hypogonadism | 8-2-07 |
| 15. | Rosiglitazone 1 mg + Metofmin HCl 500 mg tablet (Additional Strength) | As approved | 8-2-07 |
| 16. | Loteprednol Etabonate 5 mg + Tobramycine 3 mg per ml ophthalmic solution | For steroid responsive inflammatory ocular condition | 14-2-07 |
| 17. | Amlodipine (As Besylate) 5mg + Metoprolol (As Succinate) ER 25 mg (Additional Strength) | As approved | 14-2-07 |
| 18. | Glibeneclamide 5 mg + MetFormin SR 850 mg tablet (Additional Strength) | As approved | 15-2-07 |
| 19. | Aceclofenac injection (150 mg/3 ml) | For short term treatment of pain due to post operative and post traumatic cases | 19-2-07 |
| 20. | Metoprolol (As Succinate) 50 mg/ 25 mg ER) + Amlodipine 5 mg/2.5 mg tablet (For Additional Indication) | Angina pectoris in adults | 20-2-07 |
| 21. | Ramipril 10 mg + HCTZ 12.5 mg tablet (Additional Strength) | As approved | 20-2-07 |
| 22. | Amlodipine (As Besylate) 5 mg/5 mg/10 mg + Valsartan 80 mg/160 mg/160 mg tablet | Essential hypertension | 21-2-07 |
| 23. | Glatiramer Acetate injection (20 mg/ml) | Multiple sclerosis | 21-2-07 |
| 24. | Baclofen ER (GRS) Capsule 45 mg/60 mg (Additional Strength) | As approved | 22-2-07 |
| 25. | Amlodipine (As besylate) 5 mg/10 mg + Perindopril 4 mg tablet | For hypertension | 23-2-07 |
| 26. | Olopatadine HCl Tablet 2.5 mg/5 mg | Allergic nasal inflammation hives, itching with skin diseases | 26-2-07 |

| 27. | Metoprolol (As Succinate) ER 25 mg/50 mg + Ramipril 2.5 mg/5 mg | Essential hypertension | 26-2-07 |
|-----|---|---|---------|
| 28. | Rabeprazole Sodium 20 mg/10 mg with Sodium Bicarbonate as buffer | For Duodenal ulcer, G.U, Z.E.S. & GERD | 26-2-07 |
| 29. | Ciclesonide 160 mcg/320 mcg + Formoterol Fumarate (12 mcg) (Dry powder inhaler additional strength) | As approved | 27-2-07 |
| 30. | Moxonidine tablet 0.2 mg/0.3 mg/0.4 mg | For hypertension | 27-2-07 |
| 31. | Ceftriaxone (As Sodium) 125 mg/250 mg + Sulbactam (As sodium) 62.5 mg/125 mg for injection (Additional Strength) | As approved | 1-3-07 |
| 32. | Quetiapine (As Fumarate) SR 400 mg tablet (Additional Strength) | As approved | 5-3-07 |
| 33. | Cabergoline 1mg tablet (Additional Strength) | For Hyperprolactinemia | 6-3-07 |
| 34. | Levofloxacin Oral Solution (each 240 ml bottle contains Levofloxacin 125 mg/5 ml) | Antibiotic (For adult use only) | 7-3-07 |
| 35. | Clindamycin 1% + Isotretinoin 0.05% gel | For acne | 8-3-07 |
| 36. | Estradiol topical emulsion 2.5 mg/gm | Vasomotor symptoms Associated with menopause | 16-3-07 |
| 37. | Levosalbutamol MDI (50 mcg/actuation) & Dry Powder inhaler (100 mcg/ capsule) (New Dosage Form) | As approved | 16-3-07 |
| 38. | Polonosetron (As HCl) for Injection 0.25 mg/5 ml | Chemotherapy induced nausea & Vomiting | 20-3-07 |
| 39. | Cifixime 200 mg + Erdosteine 300 mg Capsules | Acute Exacerbation of chronic bronchitis | 21-3-07 |
| 40. | Aminophylline CR 225/350 mg | Asthma, COPD, Bronchitis | 21-3-07 |
| 41. | Etoricoxib 10 mg + Methyl salicylate 20 mg per gm of cream | Acute Musenlosleteal pain | 21-3-07 |
| 42. | Eberconazole cream (1%) | Dermatophyte infection of skin such As T-corporis, T-c, T-Pedis | 22-3-07 |
| 43. | ISMN 60 mg Sr + Atenolol 50 mg/100 mg | For angina pectoris | 23-3-07 |
| 44. | Levonorgestrel 0.03 mg tablet | For control of fertility | 23-3-07 |
| 45. | Bortezomib 2 mg/vial (Additional pack size) | As approved | 28-3-07 |
| 46. | Comb kit of two capsule of Azaranari 150 mg each + 1 tablet of Ritonavir 100 mg) | For HIV infection | 2-4-07 |
| 47. | Mecobalamine 500 mcg + Alpha Lipoic acid 100 mg Soft Gelatin capsule | For diabetic neuropathy | 2-4-07 |
| 48. | Propofol 10 mg + Lignocaem 1 mg per ml injection | For induction of anesthesia | 4-4-07 |
| 49. | Ezetimibe 10 mg + Fenofibrate 160 mg tablet | For combined hypertipidemia | 4-4-07 |
| 50. | Methylprednisolone Acrponate topical solution 0.1% | Atopic dermatitis (eczema) inflammatory and allergic skin reaction | 4-4-07 |
| 51. | Citicoline (750 mg) tablet (Additional Strength), syrup 500 mg/5 ml (new dosage form) (not for children) | As approved | 10-4-07 |
| 52. | Ertapenem for injection (1 gm/vial) | Antibiotic | 10-4-07 |
| 53. | Letrozole 2.5 mg tablet (for additional indication) | For induction of ovulation in anovulatory infertility (additional indication) | 10-4-07 |
| 54. | Quetiapine SR Tablets 50mg (Additional Strength) | As approved | 10-4-07 |
| 55. | Metoprolol (As Succinate) 25 mg + Amlodipine (As Besylate) 5 mg (Additional Strength) | As approved | 10-4-07 |
| 56. | Sunitinib Maleate Capsules 12.5 mg/25 mg/50 mg | For G.I. Stromal tumor, advanced renal cell carcinoma | 10-4-07 |
| 57. | Comb pack of 1 bottle of Ciprofloxacin Injection (200 mg/ 100 ml) & 1 bottle of Ornidazole injection 500 mg/100 ml) | Antibiotic | 20-4-07 |

| 58. | Aprepitant 40 mg capsules (Additional Strength) | Prevention of post-operative nausea & vomiting (additional indication) | 24-4-07 |
|-----|--|--|---------|
| 59. | Olopatadime Ophthalmic solution (0.2%) (Additional Strength) | As approved | 24-4-07 |
| 60. | Combi pack of 1 tablet of Residronate sodium (35 mg) + 6 | For osteoporosis in postmenopausal women | 1-5-07 |
| 61. | tablet of Calcium carbonate equivalent to 500 mg calcium each Levocetirizem 2 HCl Dispersible tablet (2.5 mg) | For allergic rhinitis & chronic urticaria | 8-5-07 |
| 62. | Amlodipine (As Besylate) 10 mg + Perindopril 8 mg tablets (Additional Strength) | As approved | 7-5-07 |
| 63. | Etodolac ER tablets 400 mg/500 mg | For osteoarthritis & rheumatoid arthritis | 9-5-07 |
| 64. | Levosulpiride tablet (25 mg) | For G.I. problem like fanatical dyspepsia, nausea, vomiting and diabetic gastroperisis | 10-5-07 |
| 65. | Rimonabant tablet 20 mg | For Obese Patients BMI≥ 30 kg/m2 or overweight patients BMI≥ 27 kg/m2 with Associated risk factors such As type-I diabetes or dyslipidemia | 11-5-07 |
| 66. | Lactitol monohydrate Syrup 66.67/100 ml | For constipation | 14-5-07 |
| 67. | S(-) Metoprolol (As Succinate) 25 mg + S(-) Amlodipine (As Besylate 2.5/5 mg | For essential hypertension | 16-5-07 |
| 68. | Amlexanox Oral Paste (5%) | For apthous ulcers in people with normal immune system | 16-5-07 |
| 69. | Lenalidomide capsule (5mg/10mg/15mg/25mg) | For transfusion dependent anemia due to myalodysplastic syndrome associated with deletion 5q cytogenic abnormality | 16-5-07 |
| 70. | Cefotaxim (As Sodium) 0.5 mg/1 gm + Sulbactam (As Sodium) 250 mg/500 mg injection (for Additional Indication) | For surgical prophylaxis (additional indication) | 18-5-07 |
| 71. | Lornoxicam tablet 4 mg/8 mg | For Osteoarthritis rheumatoid arthritis | 21-5-07 |
| 72. | Atenolol 25 mg/50 mg + Indapamide SR 1.5 mg | For essential hypertension | 21-5-07 |
| 73. | Livetiracetam concentration for infusion 100 mg/ml (new dosage form) | As adjunctive therapy in partial onset seizures in adults with epilepsy when oral administration is temporarily not feasible | 21-5-07 |
| 74. | Eplerstat tablet 50 mg | For abnormality of vibration sense and abnormal change in heart rate associated with diabetic peripheral neuropathy | 25-5-07 |
| 75. | Citicoline tablet 200 mg (Additional Strength) | As approved | 25-5-07 |
| 76. | Varenicline (As Tartrate) 0.5 mg/ 1 mg tablets | As an aid to smoking cessation treatment in adults | 29-5-07 |
| 77. | Azithromycin eye drops 1% | Conjunctivitis | 6-6-07 |
| 78. | Ciclesonide Nasal Spray 50 mcg/spray | For seasonal & perennial allergic rhinitis in adults, adolescents aged ≥ 12 years | 8-6-07 |
| 79. | Ceftazidime 500 mg/1000 mg + Sulbactam (As Sodium) 250 mg/500 mg for injection | Antibiotic | 8-6-07 |
| 80. | Isotretinoin 0.05% + Erythromycin 2% gel | For acne | 8-6-07 |
| 81. | Aliskiren Tablet 150 mg/300 mg | For hypertension | 8-6-07 |
| 82. | Glimepiride 1 mg/2 mg + Metformin SR 1000 mg tablet (Additional Strength) | As approved | 8-6-07 |
| 83. | Cefditorene pivoxil dispersible tablet (30 mg/50 mg) (New Dosage form for children) | As approved | 8-6-07 |
| 84. | Dexrabeprazole Sodium Enteric Coated tablet 5mg/10mg | For Duodenal ulcer, Gastric ulcer & GERD | 13-6-07 |
| 85. | Renolazine Extend Release tablet (500 mg) | For Chronic Angina | 14-6-07 |
| | | - | |

| 86. | Phenytoin Sodium ER capsules (300 mg) (New dosage form) | For control of generalized toni-clonic and treatment & | 14-6-07 |
|------|--|---|---------|
| | | prevention of seizures occurring during or following neurosurgery | |
| 87. | Alfuzocin HCl ER (10 mg) + Dutasteride 0.5 mg tablet | For BPH in men with enlarged prostate | 14-6-07 |
| 88. | Cefepime injection 500 mg/1 gm/2 gm (For additional indication) | For bacterial meningitis in children | 14-6-07 |
| 89. | Tranexamic acid mouthwash (50 mg/ml) | For prevention of oral hemorrhage in anticoagulant treated patients undergoing oral surgery | 15-6-07 |
| 90. | Deflazocort oral suspension (6 mg/5 ml) (new Dosage Form) | As approved | 15-6-07 |
| 91. | Tolterodine tartrate ER 4 mg + Tamsulocin HCl ER 0.4 mg tablet | For BPH with overactive bladder | 15-6-07 |
| 92. | Omeprazole 20 mg/40 mg + Sodium Bicarbonate 1100 mg capsules | For duodenal ulcer, gastric ulcer & GERD | 15-6-07 |
| 93. | Dexrabeprazole Sodium E. coated tablet 5mg/10mg | For duodenal ulcer, gastric ulcer and GERD | 13-6-07 |
| 94. | Cefepime Injection (New indication) | For bacterial meningitis in children | 14-6-07 |
| 95. | Phenytoin Sodium E.R (300mg) | For control of generalised tonic-clonic (grand-mal) and prevention and treatment of seizures occuring during or following neurosurgery | 14-6-07 |
| 96. | Tranexamic acid Mouth Wash (50mg/ml) | For prevention of oral hemorrhage in anticoagulant treated patients undergoing oral surgery | 15-6-07 |
| 97. | Deflazacort Oral Suspension (6mg/5ml) (New dosage form) | For asthma, rheumatoid arthritis when glucocorticosteriod therapy is warranted | 15-6-07 |
| 98. | Tolterodine Tartrate 4mg (ER) + Tamsulosin HCl 0.4mg Capsule | For BPH with overaction bladder in Adults only | 15-6-07 |
| 99. | Omeprazole (E.coated) 40mg + Sodium Bicarbonate 1100mg capsule | For gastric ulcer, duodenal ulcer and GERD | 18-6-07 |
| 100. | Mebeverine HCl prolonged release capsule (200mg) | For irritable bowel syndrome | 19-6-07 |
| 101. | Azithromycin 500mg + Ambroxol S.R (75mg) Tablet | For RTI caused by susceptible organisms in adults | 5-7-07 |
| 102. | Pholcodiene Syrup (4mg/5ml) | For symptomatic relief of dry ticklish and unproductive coughs in children above 1 years | 6-7-07 |
| 103. | Exenatide Injection pre-filled pen (250mcg/ml) | For adjunctive therapy to improve glycemic control in patients with Type-II diabetes mellitus, who have not achieved adequate glycemic control with Metformin, a sulfonyl urea or a combination of metformin and sulfonylurea | 6-7-07 |
| 104. | Moxifloxacin HCl (0.5%) + Ketorolac Tromethamine 0.5% Ophthalmic Solution | For reduction of post-operative inflammatory conditions of the eyes | 11-7-07 |
| 105. | Levofloxacin (15mg) + Prednisolone Acetate 10mg per ml eye drops | For post-operative steriod responsive inflammatory ocular conditions when ocular bacterial infections or a risk of bacterial infection exists | 11-7-07 |
| 106. | Loteprednol Etabonate (0.5%) + Levofloxacin (1.5%) Eye drops | For post-operative steriod responsive inflammatory ocular conditions when ocular bacterial infections or a risk of bacterial infection exists | 12-7-07 |
| 107. | Clotrimazole (10mg) + Methyl-prednisolone aceponate 1mg/gm of cream | For fungal infection of skin associated with inflammation | 12-7-07 |
| | | | |

| 108. | Capecitabin Tablet (150mg/ 500mg) (for additional indication) | (i) First line treatment of advanced gastric cancer in combination with platinum based regimen. | 16-7-07 |
|------|---|--|----------|
| | | (ii) Treatment of patients with locally advanced or metastatic breast cancer in combination with docetaxel following failure of cytotoxic chemotherapy | |
| 109. | Rifabutin Capsule (150mg) | For treatment of pulmonary T.B. only in patients co- infected with HIV. (It should never be used as a substitute to Rifampicin in T.B. patients without HIV co-infection) | 16-7-07 |
| 110. | Lapatinib (as Ditosylate) (250mg) Tablet | In combination with capecitabine it is indicated in patients with advanced or metastatic breast cancer whose tumour overexpress HER2 and who have received prior therapy including trastuzumab | 24-7-07 |
| 111. | Levocetirizine Di-HCl mouth dissolving tablets (5mg) (new dosage form) | For the treatment of allergic rhinitis and chronic urticaria | 25-7-07 |
| 112. | Idaravone Injection (1.5mg/ml) | For improvement of neurologic symptoms, disorders of activitis of daily living and functional disorder associated with acute ischemic stroke | 26-7-07 |
| 113. | Diclofenac Sodium (75mg/ml) Injection (Additional Strength) | For management of painful conditions in post-operative cases, rheumatoid arthritis and osteoarthritis | 30-7-07 |
| 114. | Cefpodoxime (as Proxetil) (100mg/200mg) + Clavulanic acid (62.5/125mg) tablet | For LRTI, URTI and UTI | 30-7-07 |
| 115. | Atorvastatin (40mg) + Ezetimibe (10mg) Tablet (Additional Strength) | As approved | 30-7-07 |
| 116. | Cefepime (250mg) + Tazobactam (31.25mg) Injection (Additional Strength) | As approved | 30-7-07 |
| 117. | VSL#3 (Probiotic Blend) sachet | For rotavirus diarrhoea in children | 31-7-07 |
| 118. | Sorafenib (as Tosylate) (200mg) Tablet | For advanced renal cell carcinoma | 31-7-07 |
| 119. | Methylcobalamin (500mcg) + Alpha Lipoic Acid (100mg) + Pyridoxine (3mg) + Folic acid (1.5mg) Soft Gelatin Capsule | For diabetic neuropathy | 31-7-07 |
| 120. | Halobetasol Propionate (0.5mg) + Salicylic Acid 30mg per gm of ointment | For plaque psoriasis | 1-8-07 |
| 121. | Combipack of 12 tablet of Artesunate (50mg) each + 3 tablet of each containing Sulfadoxine (500mg) + Pyrimethamine (25mg) Tablet. (Additional Strength) | As Approved | 1-8-07 |
| 122. | Omeprazole (20mg/40mg)+Sod. Bicarbonate-600mg+Mg.Hydroxide-700mg chew tab | For duodenal, gastric ulcer, GERD, and erosive esophagitis | 1-8-2007 |
| 123. | Deferasirox dispersible tablet (400mg) (Additional Strength) | As approved | 17-8-07 |
| 124. | Tegafur (100mg) + Uracil (224mg) capsule | For first line treatment of metastatic colorectal cancer in combination with Calcium folinate | 22-8-07 |
| 125. | Bortezomib Injection (Additional Indication) | For the treatment of patients with mantle cell lymphoma who have received at least one prior therapy | 5-9-07 |
| 126. | Amlodipine (5mg) + Nebivolol (5mg) tablet (Additional Strength) | As approved | 6-9-07 |
| 127. | Ceftazidine (2gm) + Sulbactam (1gm) Injection (Additional Strength for Additional Indication) | For serious abdominal infection. | 6-9-07 |

| 128. | Polyethylin Glycol 400 (0.4%) + Propylene Glycol (0.3%) | For temprory relief of dryness, irritation burning, stinging, grittiness or foreign body sensation caused due to environmental factors | 18-9-07 |
|------|--|---|----------|
| 129. | Resperidone E.R. (long-acting) injection (12.5mg/vial) (Additional Strength) | As approved | 25-9-07 |
| 130. | Levocetirizine Di-HCl syrup (2.5mg/5ml) (For use in children 2 years and above) | As approved | 26-9-07 |
| 131. | R-Sibutramine Capsule (2.5mg/5mg) | As an adjunctive therapy for the management of obesity including weight loss and maintenance of weight loss in pateints whose BMI is 30Kg/m2 or higher | 26-9-07 |
| 132. | Cefuroxime Axetil (250mg) + Clavulanic Acid (125mg) Tablet | For adult patients with infection caused susceptible micro-organisms including S.aureus viz UTI, upper and lower RTI and gonococcal urethritis | 27-9-07 |
| 133. | Sulphur Hexafluoride Powder and solvent for dispersion for injection (45mcg/ml) | For diagnostic use in echocardiography doppler of macrovasculature an doppler of microvasculature | 28-9-07 |
| 134. | Ondansetron orally disintegrating tablet (4mg/8mg) (Additional dosage form) | For chemotherapy induced nausea and vomiting | 28-9-07 |
| 135. | Cefuroxime Axetil (250mg) + Clavulanic acid (125mg) tablet | For UTI, URTI and LRTI and gonococcal urethritis | 5-10-07 |
| 136. | Cefuroxime-125/250/500mg+Clav.Acid-31.25/62.5/125mg | For adult patients with infection caused by susceptible microbs,including s.aureus.(e.g-UTI, Upper & lower RTI & gonococcal urethritis) | 10/05/07 |
| 137. | Amiodarone Injection (Additional Indication) | For cardiopulmonary resuscitation in the event of cardiac arrest related to ventricular fibrilation resistant to external electric shock | 5-10-07 |
| 138. | Risedronate Sodium tablet (35mg) (Additional Indication) | For osteoporosis in men at high risk of fracture | 8-10-07 |
| 139. | Sucralfate (70mg) + Metronidazole (10mg) + Povidone Iodine (50mg) per gram of ointment | For infected wounds and ulcers | 9-10-07 |
| 140. | Levetiracetam Tablet (Additional Indication) | (i) As monotherapy in partial onset seizures with or without secondary generalisation in patients with 16 years of age with newly diagnosed epilepsy.(ii) As adjunctive therapy in myoclonic seizures in | 9-10-07 |
| | | adults and adolescents from 12 years of age with Juvenile myoclonic epilepsy. In primary generalised tonic-clonic seizures in adults and adolescents from 12 years of age with idiopathic generalised epilepsy | |
| 141. | Combipack of two tablet of chloroquine phosphate (250mg) each + 14 tablet of Proguanil HCl (100mg) each | For drug resistant malaria | 10-10-07 |
| 142. | Paclitaxel (protein bound particle) for injectable suspension (paclitaxel (100mg) + Albumin Human 900mg per 20ml | For breast cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy | 11-10-07 |
| 143. | Olmesartan Medoxomil (20mg) + Amlodipine Besylate (5mg) Tablet | For mild to moderate hypertension | 15-10-07 |
| 144. | Atorvastatin Calcium (10mg) + Ramipril (2.5mg/5mg) tablet | For treatment of patients with both hypertension and hypercholesterolemia | 15-10-07 |
| 145. | Omeprazole(20mg) + Sodium bicarbonate (1100mg) tablet (Additional Strength) | As approved | 15-10-07 |
| 146. | Moxifloxacin HCl (5mg) + Prednisolone Acetate (10mg) per ml (eye drops) | For inflammation after ocular surgery and where a risk of bacterial ocular infection exists | 16-10-07 |
| _ | | | |

| 147. | Methylcobalamine (500mcg) + Alpha Lipoic acid (100mg) + Pyridoxine HCl (1.5mg) + Thiamine (2mg) + Calcium Picolinate (200mg) tablet | For diabetic neuropathy | 16-10-07 |
|------|---|--|----------|
| 148. | Satranidazole (300mg) + Ofloxacin (200mg) Tablet | For diarrhoea of mixed infection in adults | 18-10-07 |
| 149. | Methylcobalamin 500mcg+beta carotine-10.33mg+alpha lipoic acid 100mg+B6 1.5mg+B1 2mg & Cr.polynicotinate | For adult patients with diabetic neuropathy | 18-10-07 |
| 150. | Calcium Pidolate equivalent to Calcium 500mg + Vit-D3 250 IU | For osteoporosis and calcium deficiency state | 23-10-07 |
| 151. | Etizolam Tablet (0.5mg/1mg) | For anxiety disorder associated with depression | 29-10-07 |
| 152. | Oxaprozine Tablet (600mg) | For relief of signs and symptoms of osteoarthritis, rheumatoid arthritis | 30-10-07 |
| 153. | Prulifloxacin (600mg) Tablet | For acute uncomplicated lower urinary tract infection (simple cystitis), complicated lower UTI, acute exacerbation of chronic bronchitis | 30-10-07 |
| 154. | Cilostazole E.R. Tablet (100mg/200mg) | For management intermittent claudication | 30-10-07 |
| 155. | Atorvastatin Calcium (10mg) + Fenofibrate (145mg) (Additional Strength) | As Approved | 30-10-07 |
| 156. | Sitagliptin Tablet (25mg/50mg/100mg) | As adjunct to diet and exercise to improve glycemic control in patients with type-II diabetes | 30-10-07 |
| 157. | Miglitol (50mg) + Metformin HCl SR (500mg) tablet | For adult with Type-II diabetes mellitus where simple drug therapy, diet and exercise do not result in adequate glycemic control | 31-10-07 |
| 158. | Mupirocin (2%) + Methylprednisolone Aceponate (0.1%) Cream | For atopic dermatitis | 31-10-07 |
| 159. | Methyl cobalamine 500mcg+Folic acid1.5mg+B5-5mg,+Zn.sulphate 25mg+Cr.Polynicotinate 200mcg+Selenium-65mcg+Inositol-1.5mg + Alpha lipoic acid -100mg capsule | For vitamins and mineral deficiency states | 31-10-07 |
| 160. | Methylcobalamine 500mcg+Folic acid1.5mg+ Alpha lipoic acid-300mg+Benfetiamin-50mg capsule | For diabetic neuropathy | 13-11-07 |
| 161. | Methyl prednisolone aceponate-1mg + Salicylic acid-50mg ointment | For treatment of psoriasis | 19-11-07 |
| 162. | Propanol 50gm+Povidone iodine-1gm in 100ml solution | For surgical & hygenic hand disinfection | 19-11-07 |
| 163. | Propanol-18 gm+Ethenol(100%)-45gm in 100ml solution | -do- | 19-11-07 |
| 164. | Voglibose dispersible tab.0.2mg/0.3mg | As Approved | 20-11-07 |
| 165. | | | |
| 166. | Ceftriaxone-2gm+ Tazobactum 250mg for injection | As Approved | 21-11-07 |
| 167. | Methylprednisolone aceponate-1 mg + Mupirocin-20 mg cream | For the treatment of atopic dermatitis | 21-11-07 |
| 168. | Citicholine (As Sod.)-200mg/500mg/750mg granules per sachet | As Approved | 26-11-07 |
| 169. | Butarphenol tartrate 10mg/ml nasal spray | For management of pain in patients using opioid analgesics | 26-11-07 |
| 170. | Deferasirox 100mg dispersible tablet (Additional Strength) | As Approved | 28-11-07 |
| 171. | Methyl cobalamine 500mcg+Folic acid1.5mg+B6- 3mg+Zn.sulphate 61.8mg+Ca.Pantothenate 10mg+ Niacinamide-50mg+Chrimium-50mcg capsule | For vitamins and mineral deficiency patients | 29-11-07 |

| 172. | Methylcobalamine 500mcg+Folic acid 1.5mg+ Alpha lipoic acid—100mg+B6-3mg+Thiamine 10mg capsule | For diabetic neuropathy | 29-11-07 |
|------|--|---|----------|
| 173. | Doxophylline-400mg+Salbutamol (as sulphate)-2mg/4mg tablets | For treatment of Bronchial asthma in adults | 29-11-07 |
| 174. | Gatifloxacin (As Sequihydrate)-3mg + Flurbiprofen sod 0.3mg/ml eye drops | For reduction of post-operative inflammatory conditions of eye, when bacterial infection exists | 29-11-07 |
| 175. | Sertanidazole-300mg+ Ofloxacin-200mg tablet | For treatment of diarrhoea of mixed infection in adults patients | 29-11-07 |
| 176. | Zoledronic acid 5mg/100ml solution for infusion (Additional Indication) | Post-menopausal osteoporosis | 6-12-07 |
| 177. | Olmesartan medoximil-20mg + Ramipril-5mg tablet | For treatment of essential hypertention | 6-12-07 |
| 178. | Levosulpiride Tablet (50mg/100mg) (Additional Strength for Additional Indication) | For depression and schizophrenia in adults | 6-12-07 |
| 179. | Combipack of 12 Artesunate –50mg tablets & 6 Mefloquin-250mg tablets | For drug resistant P.falciparum malaria infection | 6-12-07 |
| 180. | Pregabalin-75mg/150mg +Methylcobalamine 500mcg cap | For the treatment of adult patients with peripheral neuropathy | 10-12-07 |
| 181. | Doxycycline-100mg +L.bacillus spores-5 billion | For adult patients prone to intra-adominal bacterial infection & Antibiotic associated diarrhoea | 11-12-07 |
| 182. | Cefprozil 250mg/500mg+Clav. Acid-62.5mg/125mg tab | For the treatment of adult patients with upper & lower RTI, Skin & Skin structure infection | 11-12-07 |
| 183. | Dextrometharpahan HBr-10mg+Chlorphenaramine maleate- 2mg/5ml syrup | For the temporary relief of cough due to throat irritation, sneezing & running nose | 11-12-07 |
| 184. | Lornoxicam 8mg/2ml inj | For short-term treatment of mild to moderate pain, when oral administration is inappropriate(i.e. after dental surgery) | 11-12-07 |
| 185. | Apomorphine HCl-2mg/3mg Sublingual tablet | For erectile dysfunction in adult men | 11-12-07 |
| 186. | Escitlopram (as Oxalate)10mg + Clonazepam-0.5mg tablet (Additional Strength) | For comobid depression and anxiety disorder | 12-12-07 |
| 187. | Doxophylline injection-10mg/ml | For bronchial asthama & COPD in adult patients, when oral therapy is inappropriate | 31-12-07 |

List of Drugs approved during January to December 2008

| Sl. No. | Drug Name | Indication | Date of Approval |
|---------|--|---|------------------|
| 1. | Me-salicylate 30% + Menthol 10% + Camphor 4% Cream | For temporary relief from pain, swelling & inflammation due to musculoskeletal disorder | 29-11-2007 |
| 2. | Voglibose 0.2mg/0.3mg tablet (Mouth dissolving) | For improvement of parantial hyperphemia in diabetes mellitus only. When diet and/or exercise or oral, hypoglycaemic drug or insulin preparation in addition to diet and or exercise do not result in adequate glycemic control | 28-12-2007 |
| 3. | Hydroxychloroquine 400mg tablets (Additional Strength) | Same as Approved | 31-1-2008 |
| 4. | ISMN 30 mg SR+Metoprolol Succinate equivalent to Metoprolol Tartrate 25mg/ 50mg tablets (Additional Strength) | Same as Approved | 2-1-2008 |
| 5. | Halobetasol 0.05%+Clotrimazole 1.0% cream | For the treatment of steroid responsive fungal infections including eczema and psoriasis | 4-1-2008 |

| 6. | Citicholine Syrup 750mg/5ml | For the treatment of disturbance of consciousness resulting from head injuries, brain operation and acute stage of cerebral infraction in adults only | 4-1-2008 |
|-----|---|---|-----------|
| 7. | Levodopa 200mg + Carbidopa 50mg C.R. tablet | For Parkinson's Disease | 8-1-2008 |
| 8. | Nanocrystalline Silver Gel 0.002% | For the treatment of burns of would infections | 4-1-2008 |
| 9. | Cinitapride (as Hydrogen tartrate) 1 mg tablet | For the treatment of gastroesophageal reflux disease and functional inconveniences of the gastrointestinal motility | 4-1-2008 |
| 10. | Tenofovir disoproxil fumarate 300mg + Lamivudine 300mg tablet | For the treatment of adult patients with HIV infection | 7-1-2008 |
| 11. | Cefixime 200mg/100mg+Lactic acid bacillus 2.5 billion spores tablet & dispersible tablet | For the treatment of Urinary tract infections, respiratory tract infections, biliary tract infections in adult patients prone to antibiotic associated diarrhoea | 7-1-2008 |
| 12. | D-Trans Fentanyl Metrix system (12mcg/hr) (Additional Strength) | Same as Approved | 7-1-2008 |
| 13. | Ceftazidime 500mg+Tobramycin 60mg per vial (Injection) | For the treatment of lower respiratory tract infection caused by pseudomonas aeruginosa | 8-1-2008 |
| 14. | Atazanavir 300mg capsules (Additional Strength) | Same as approved | 8-1-2008 |
| 15. | Ertapenem for injection 1gm/vial (Additional Indication) | For the prophylaxis of surgical site infections following elective colorectal surgery | 15-1-2008 |
| 16. | Vildagliptin Tablets 50mg/100mg | As an adjunct to diet and exercise to improve glycemic control in patients with type-II diabetes mellitus | 18-1-2008 |
| 17. | Atorvastatin 10mg+Aspirin 75/150mg Capsules | For the treatment of dyslipidemia associated with atherosclerotic arterial disease with risk of myocardial infraction, stroke or peripheral vascular disease | 18-1-2008 |
| 18. | Smatriptan (as Succinate) 50mg+ Naproxen Sodium 550 mg tablets | For the treatment of acute migraine | 22-1-2008 |
| 19. | Sodium Valproate 666mg + Valproic acid 290 mg equivalent as Sodium valproate 1000 mg C.R. tablet (Additional Strength) | Same as Approved | 24-1-2008 |
| 20. | Cefepime (as Hydrochloride) 2000/1000/500mg + Sulbactam (as Sodium) 1000/500/250 mg per vial of injection | For serious CRTI, uncomplicated and complicated UTI, Uncomplicated skin & skin structure, infection acute exacerbation of chronic bronchitis & intra-abdominal infection | 24-1-2008 |
| 21. | Paclitaxel Nanoparticle Solution for injection 20mg/ml (Additional Indication) | Ovarian cancer, non-small cell lung cancer and AIDS related Kaposis Sarcoma | 22-1-2008 |
| 22. | Sorafenib tablets 200mg (Additional Indication) | For Hepatocellular Carcenoma | 22-1-2008 |
| 23. | Zanamvir inhalation powder 5mg/blister (Additional Indication) | For prophylaxes and treatment of both influenza A&B in adults and children (greater than or equal to 5 years) | 22-1-2008 |
| 24. | Mycophenolate Mofetil Capsule 250mg, 500mg & Oral suspension 200mg/ml (Additional Indication) | For the prophylaxis of acute organ rejection in patients receiving allogeneic hepatic transplantation | 23-1-2008 |
| 25. | Combikit 4 tablets of Alendronate sodium trihydrate equivalent to Alendronic acid 70mg each + 24 tablets of Calcium carbonate 1250mg (equivalent to 500 mg Calcium) + Cholecalciferol-250IU | For the treatment of osteoporosis in post-menopausal women | 23-1-2008 |
| 26. | Aceclofenac injection 150mg/ml | For short term treatment of pain due to post operative and post traumatic cases | 23-1-2008 |
| 27. | Atorvastatin 10mg + Fenofibrate 80mg tablet (Additional Indication) | For the treatment of uncomplicated and complicated UTI, uncomplicated skin and skin structure infections, and complicated intra-abdominal infections | 25-1-2008 |

| 28. | Sodium Valproate 499.5mg+ Valproic acid 217.5mg equivalent as Sodium Valproate-750mg CR tablet (Additional Strength) | Same as Approved | 24-1-2008 |
|-----|---|---|-----------|
| 29. | Diclofenac 50mg + Serratiopeptidase 10mg enteric coated tablet | For the treatment of acute pain in adults | 24-1-2008 |
| 30. | Gabapentin 100mg+Methylcobalamin 500mcg tablets (Additional Strength) | Same as Approved | 24-1-2008 |
| 31. | Cefpirome (as Sulphate) 500/1000mg/2000mg + Sulbactam (as Sodium) 250mg/500mg/1000mg powder for injection | For the treatment of infection caused by sensitive organisms viz. lower RT1, complicated UTI, Septicaemia, bacteremia | 24-1-2008 |
| 32. | Metformin HCl ER 500 mg + Fenofibrate 80mg/160mg tablet | For the treatment of Type-II diabetes associated with mixed dyslipidemia | 25-1-2008 |
| 33. | Salicylic acid 10mg/20mg (Foaming facewash solution) | For the treatment of acne vulgaris | 28-1-2008 |
| 34. | Ozagrel HCl tablets 100 mg/200mg | For the treatment of bronchial asthma | 28-1-2008 |
| 35. | Rivastigmine transdermal patch (each patch of 5m2, 10m2, 15m2 and 20m2 contain Rivastigmine base 9/18/27/36 mg respectively | For the treatment of patient with mild to moderately severe dementia of the Alzheimer type and dementia associated with Parkinsons disease | 28-1-2008 |
| 36. | Daptomycin powder for concentrate solution for infusion (350mg/500mg per vial) | For the treatment of complicated skin and skin structure infections and for staphylococcus aureus blood-stream infections (bacteremia) | 28-1-2008 |
| 37. | Dextromethorphan Orodispersible strip 5.5 mg/11mg | For the relief of persistent dry irritating cough in adult patients only | 28-1-2008 |
| 38. | Diclofenac sodium 50mg + Dicyclomine HCl 20 mg per ml of injection | For the treatment of acute colicky pain in adults | 28-1-2008 |
| 39. | Travoprost 40mcg+ Timolol (as Maleate) 5 mg per ml of eye drops | For decrease of intraocular pressure in patients with open angle glaucoma or ocular hypertension, who are insufficiently responsive to topical beta blockers or prostaglandin analogues | 28-1-2008 |
| 40. | Amoxicillin 250mg/125mg + Cloxacillin 250mg/125mg + Lactic Acid Bacillus 1.7 Billion spores | For RTI, SRS of tissue infection, bones & joint infection, ENT infection in adults prone to antibiotic associated diarrhoea | 30-1-2008 |
| 41. | Amoxicillin 250mg + Lacto-Bacillus 1.7 billion Spores | For RTI, SRS of tissue infection, bones & joint infection, ENT infection in adults prone to antibiotic associated diarrhoea | 31-1-2008 |
| 42. | Amoxycillin 250mg + Cloxacillin 250mg+Lactolbacillus 2 billion spores capsules | For the treatment of lower RTI, skin & soft tissue bone & joint infections, ENT infections in adult patients prone to antibiotic associated diarrhoea | 30-1-2008 |
| 43. | Tiotropium (as Tiotropium Bromide Monohydrate) 18mcg + Formoterol Fumarate Dihydro 12mcg + Ciclesomide 400mcg per capsules | As 3rd line treatment of severe cases of COPD when monotherapy and second line therapy with two drugs do not respond adequately | 31-1-2008 |
| 44. | Metoprolol Succinate Equivalent to Metoprolol Tartrate [ER 25/50mg] + Atorvastatin (As Calcium) 10 mg tablets | For the treatment of patient with both essential hypertension and hypercholesterolemia | 28-1-2008 |
| 45. | Levocetrizine Di HCl Dispersibel tablet 5mg/10mg (New Dosage form) | For allergic rhinitis & chronic urticaria | 28-1-2008 |
| 46. | Oseltamivir 30mg & 45 mg Capsules (Additional Strength) | Same as Approved | 28-1-2008 |
| 47. | Fondaparinax Sodium injection (Additional Indication) | Treatment of acute coronary syndrome (ACS) with and without ST-segment elevation | 28-1-2008 |
| 48. | Camostat Mesylate tablets 100mg | For remission of acute symptoms of chronic pancreatitis and post-operative reflux esophagitis | 29-1-2008 |
| 49. | Hyaluronic Acid Slat 25mg + Cross-Linked Hyaluronic acid 25mg per ml in prefilled syringe | For intradermal implantation and facial soft tissue agumentation | 30-1-2008 |
| | | | |

| Ursodeoxycholic acid tablets 600 mg | For the treatment of patients with chronic cholestatic | 20.1.2009 |
|--|--|--|
| | liver disease | 30-1-2008 |
| Dalteparin Sodium Injection (Additional Indication) | For the extended treatment of symptomatic venous thromboembolism (VTE) proximal deep thrombosis (DVT) and/or pulmonary embolism (PE), to reduce the recurrence of VTE in patients with cancer | 30-1-2008 |
| Beclomethasone Dipropionate 0.25mg +Econazole 10mg+ Neomycin 5mg per gm of Cream | For the topical treatment of bacterial and fungal infections | 30-1-2008 |
| Exenatide injection (Each 2ml vial contains-Exenatide 5mcg/10mcg per 0.2ml) Additional Strength) | For adjunctive therapy to improve glycemic control in patients with type-II diabetes mellitus, who have not achieved adequate glycemic control with metformin, a sylfonylurea or a combination of metformin and a sulfonylureas | 30-1-2008 |
| Diclofenac Sodium 50mg+Pacacetamol 500mg tablets | For the symptomatic treatment of acute pain and inflammation in patients with osteoarthritis, rheumatoid arthiritis and ankylosing spondylitis | 30-1-2008 |
| Gabapentin SR 600 mg tablets | As adjunctive therapy in the treatment of partial seizures of epilepsy | 30-1-2008 |
| Memantine 5mg/10mg+Donepezil 5 mg Tablets | For moderate to severe Alzheimer's disease | 31-1-2008 |
| Doxifluridine Capsules 100mg/200mg | For carcinoma of the stomach, colon, rectum, breast, uterine, cervix and bladder | 31-1-2008 |
| Haemodialysis Powder/Liquid Form containing citrate | For use for patients being treated by hemodialysis | 31-1-2008 |
| Carmustine (Polifeprosan 20 with carmustine Implant (Wafer) | In newly diagnosed high-grade malignant glioma patients as an adjunct to surgery and radiation. In recurrent glioblastoma multiforme patients as an | 31-1-2008 |
| | adjunct to surgery | |
| Cimetropium Bromide tablet 50mg | For the treatment of irritable bowel syndrome | 31-1-2008 |
| Cyclobenzaprine HCl Tablets 5mg/10mg | As adjunct to rest and physical therapy for relief of muscle spasm associated with acute painful musculoskeletal conditions | 31-1-2008 |
| Sertaconazole Nitrate Cream 2% | For the topical treatment of the superficial mycosis like dermatophytosis | 31-1-2008 |
| Cefpodoxime 50mg+Clavulanic acid 31.25mg dispersible tablet & dry syrup (Additional Strength) | Same as Approved | 31-1-2008 |
| Darifenacin (as Hydrobromide) 7.5 mg/15mg ER tablets | For the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency and frequency | 31-1-2008 |
| Amoxycillin 125mg/250mg/500mg + Lactobacillus 1.66 billion Spores (Capsules) | For the treatment of respiratory tract infection, skin and soft tissue infections & antibiotics associated diarrhoea in adult patients | 31-1-2008 |
| Halobetasol 0.5mg+Miconazole 20mg per gram cream | For steroid responsive dermatoses associated with superficial fungal infection of skin | 31-1-2008 |
| Montelucast 10mg+Levocetirizine 5mg film coated tablets | For the treatment of allergic rhinitis in adults only | 31-1-2008 |
| Hydroquinone 2% + Tretinoin 0.025% (w/w) Momentasone 0.10% Gel | For short term intermittent treatment of moderate to severe melasma of the face in the presence of measures for sun avoidance including use of sun screen | 31-1-2008 |
| | 8 | |
| | Beclomethasone Dipropionate 0.25mg +Econazole 10mg+ Neomycin 5mg per gm of Cream Exenatide injection (Each 2ml vial contains-Exenatide 5mcg/10mcg per 0.2ml) Additional Strength) Diclofenac Sodium 50mg+Pacacetamol 500mg tablets Memantine 5mg/10mg+Donepezil 5 mg Tablets Doxifluridine Capsules 100mg/200mg Haemodialysis Powder/Liquid Form containing citrate Carmustine (Polifeprosan 20 with carmustine Implant (Wafer) Cimetropium Bromide tablet 50mg Cyclobenzaprine HCl Tablets 5mg/10mg Sertaconazole Nitrate Cream 2% Cefpodoxime 50mg+Clavulanic acid 31.25mg dispersible tablet & dry syrup (Additional Strength) Darifenacin (as Hydrobromide) 7.5 mg/15mg ER tablets Amoxycillin 125mg/250mg/500mg + Lactobacillus 1.66 billion Spores (Capsules) Halobetasol 0.5mg+Miconazole 20mg per gram cream Montelucast 10mg+Levocetirizine 5mg film coated tablets Hydroquinone 2% + Tretinoin 0.025% (w/w) Momentasone | thromboembolism (VTE) proximal deep thrombosis (DVT) and/or pulmonary embolism (PE), to reduce the recurrence of VTEIn patients with cancer Beclomethasone Dipropionate 0.25mg +Econazole 10mg+ Neomycin 5mg per gm of Cream Exenatide injection (Each 2ml vial contains-Exenatide 5mcg/10mcg per 0.2ml) Additional Strength) Diclosenae Sodium 50mg+Pacacetamol 500mg tablets Diclosenae Sodium 50mg+Pacacetamol 500mg tablets Diclosenae Sodium 50mg+Pacacetamol 500mg tablets For the symptomatic treatment of acute pain and inflammation in patients with osteoarthritis, rheumatoid arthritis and ankylosing spondylitis As adjunctive therapy in the treatment of partial seizures of epilepsy Memantine 5mg/10mg+Donepezil 5 mg Tablets Doxistluridine Capsales 100mg/200mg For carcinoma of the stomach, colon, rectum, breast uterine, cervix and bladder Haemodialysis Powder/Liquid Form containing citrate Carmustine (Polifeprosan 20 with carmustine Implant (Wafer) Lin newly diagnosed high-grade malignant glioma patients as an adjunct to surgery and radiation. 2. In recurrent glioblastoma multiforme patients as an adjunct to surgery and radiation. 2. In recurrent glioblastoma multiforme patients as an adjunct to surgery and radiation. Sertaconazole Nitrate Cream 2% Ceripodoxime 50mg+Clavulanic acid 31.25mg dispersible tablet & dry-syrup (Additional Strength) Darifenacin (as Hydrobromide) 7.5 mg/15mg ER tablets For the treatment of overactive bladder with symptoms of the treatment of respiratory tract infection, skin and soft tissue infections & ambitotics associated with superficial mycosis like dermatophytosis For the treatment of respiratory tract infection, skin and soft tissue infections & ambitotics associated with superficial fungal infection of skin Montelucast 10mg+Levocetirizine 5mg film coated tablets For the treatment of allergic rhinitis in adults only For the treatment of allergic rhinitis in adults only For the treatment of allergic rhinitis in adults only |

| 70. | Diclofenac Sodium 100mg (SR Pellets) + Rabeprazole Sodium 20mg (E.C Pellets) Capsules | For treatment of painful inflammatory conditions associated with gastroduodenal ulcers | 31-1-2008 |
|-----|--|--|-----------|
| 71. | Ciclesonide 16mcg+320mcg/Formoterol 12mcg+Tiotropium 18mcg Dry powder inhaler | As third line treatment of severe cases of COPD, when monotherapy and second line therapy with two drugs do not respond adequately | 31-1-2008 |
| 72. | VSL# (Combination of Lactic acid bacteria (VSL#3) | For the treatment of irritable bowel syndrome | 31-1-2008 |
| 73. | Clostimethate Sodium of 1,00,000 IU Injection | For the treatment of serious infection caused Gram-Ve bacteria including LRTI, UTI, where commonly used anti-bacterial agent may be infective because of bacterial resistance | 31-1-2008 |
| 74. | Ramipril 5mg+HCTZ 12.5mg tablets (Additional Strength) | Same as Approved | 20-2-2008 |
| 75. | Metronidazole CR tablet 600mg | For the treatment of amoebiasis, urogenital trichomoniasis & giardiasis | 21-2-2008 |
| 76. | Quetiapine fumarate SR tablets | For the additional indication "acute manic episode associated with bipolar disorder | 21-2-2008 |
| 77. | Atorvastatin Calcium 80mg tablets (Additional Strength) | Additional strength (it should not be used routinely because of increased risk of adverse events. It should be reserved for patients who are considered to be at higher risk after careful risk assessment | 25-2-2008 |
| 78. | Euphorbia Prostrata Dry Extract (Ethanolic 80% (v/v) (35-70): 100 mg tablet | For the treatment of haemorrhoid | 29-2-2008 |
| 79. | Amtolmetin Guacil tablets 600mg | For the treatment of osteoarthritis & rheumatoid arthiritis | 7-3-2008 |
| 80. | Brimonidine Tartrate Opthalmic solution 0.1% (with Purite as preservation 0.0005%) (Additional Strength) | For lowering of IOP in patients with open angle glaucoma occular hypertension | 7-3-2008 |
| 81. | Methoxsalen USP-20mg Tablet | For repigmentation of idiopathic vitiligo along with phototherapy. For symptomatic control of severe, recalcitrant, disabling psoriasis in conjunction with phototherapy | 7-3-2008 |
| 82. | Cefditoren pivoxil 400mg tablets (Additional Strength) | Same as Approved | 11-3-2008 |
| 83. | Doxofylline ER tablets 400mg/800mg (Additional Strength) | For the treatment of bronchial asthma & COPD in adults | 12-3-2008 |
| 84. | Betamethasone 0.05% + Miconazole 2% Cream | For the treatment of steroid responsive dermatoses associated with superficial fungal infections of the skin | 12-3-2008 |
| 85. | Premetrxed disodium injection 100mg/vial (Additional Strength) | Same as Approved | 12-3-2008 |
| 86. | Omeprazole 20mg/40mg+Sodium bicarbonate 1680mg per sachet powder for oral suspension | For gastric ulcer, duodenal ulcer and GERD | 12-3-2008 |
| 87. | ISMN 30mg SR+Atenolol 50mg tablets (Additional Strength) | Same as Approved | 19-3-2008 |
| 88. | Temozolamide capsules 140mg/180mg (Additional Strength) | Same as Approved | 31-3-2008 |
| 89. | Amisulpiride 300mg tablets (Additional Strength) | Same as Approved | 20-3-2008 |
| 90. | Nifedipine SR Tablets 10mg/20mg | For the treatment of essential hypertension | 24-3-2008 |
| 91. | Ambroxol HCl 15mg+Guiphenesin 50mg+Terbutalin sulphate 1.25mg+Menthol 2.5mg per 5ml of syrup | For symptomatic relief of bronchospasm in bronchial asthma and chronic bronchitis | 26-3-2008 |
| 92. | Ambroxol HCl 30mg + Guaiphenesin 100mg+Terbutaline sulphate 2.5mg per 10ml syrup | For the symptomatic relief of bronchospasm in bronchial asthma & chronic bronchitis | 26-3-2008 |
| 93. | Boswellia Serrata 0.05gm/gm of cream | For the treatment of mild to moderate psoriasis | 28-3-2008 |
| 94. | Etoricoxib injection 90mg/ml | For short term use in acute painful condition in adults only | 28-3-2008 |
| | | | |

| 95. | Drospirenone 3mg+Ethinyl estradiol 0.02mg tablets (Additional Strength) | Oral contraceptive in women | 28-3-2008 |
|------|--|---|-----------|
| 96. | Nimesulide 100mg ER + Levocetirizine HCl 5mg Tablets | For symptomatic treatment of acute rhinosinusitis | 28-3-2008 |
| 97. | Dexketoprofen (as Trometamol) tablets 12.5mg/25 mg | For the treatment of mild to moderate pain, such as dental pain, dysmenorrhoea & musculoskeletal pain | 28-3-2008 |
| 98. | Nepafenac Opthalmic suspension (1mg/ml) | For the treatment of pain and inflammation associated with cataract surgery | 31-3-2008 |
| 99. | Hylan GF 20 Injection (Synvisc) (Additional Indication) | For pain management in patients suffering with osteoarthritis of the knee, hip, shoulder and ankle | 31-3-2008 |
| 100. | Docetaxel injection conc. | For additional indication "advanced gastric adenocarcinoma, locally advanced squamous cell carcinoma of head & neck, Harmone refraetory metastatic prostrate cancer, locally advanced or metastatic breast cancer | 31-3-2008 |
| 101. | Methylcobalamin 1000 mcg + Pyridoxine HCl IP 100mg+Nicotinamide IP 100mg+D-Panthenol IP 50mg per 2 ml of injection | For peripheral neuropathy | 31-3-2008 |
| 102. | Lamivudine 20mg + Stavudine 5mg dispersible tablets (Additional Strength) | Same as Approved | 1-4-2008 |
| 103. | Nicardipine HCl Injection 2.5mg/ml | For short-term treatment of hypertension when oral therapy is not feasible or not desirable | 7-4-2008 |
| 104. | Diclofenac Potassium 50mg+ Serratiopeptidase (as enteric coated pellets) 15 mg capsules (Additional Strength) | For the treatment of acute pain in adults | 8-4-2008 |
| 105. | Diacerin 50mg+Aceclofenac 100mg tablets | For the treatment of adult patients with osteoarthritis of knee and or hip joints | 11-4-2008 |
| 106. | Rabeprazole Sodium Enteric coated pellets 15.4% w/w (20mg/129.83mg) | Same as Approved | 16-4-2008 |
| 107. | Live freeze-dried Lactic acid bacteria and Bifidobacteria capsules | For the treatment of irritable bowel syndrome | 10-4-2008 |
| 108. | Miltefosin Capsule 10mg/50mg (Additional Indication) | Treatment of cutaneous leishmaniasis | 16-4-2008 |
| 109. | Silver Wound Dressing | As an antimicrobial barrier for management of partial thickness burns, incisions, skin grafts, donor sites, lacerations, abrasions and Stage I-IV dermal ulcers (vascular, venous, pressure and diabetic) | 16-4-2008 |
| 110. | Diacerin 50mg+Glucosamine sulphate 500mg tablets | For the treatment of adult patients with osteoarthritis of knee and hip joints | 21-4-2008 |
| 111. | Clostridium Botulinum type-A toxin haemagglutinin complex (Additional Indication) | For focal spasticity, including the treatment of arm symptoms associated with focal spasticity in conjunction with physiotherapy | 22-4-2008 |
| 112. | Fluconazole Dispersible tablets 50mg | For the treatment of systemic candidiasis, mucosal candidiasis, prevention of fungal infections in patients with malignancy | 22-4-2008 |
| 113. | Lenalidomide Capsule 5mg/10mg/15mg/25 mg (Additional Indication) | In combination with dexamethasone for the treatment of multiple myeloma patients who have received at least one prior therapy | 22-4-2008 |
| 114. | Sitagliptin (As Phosphate) 50mg+ Metformin HCl 500mg/1000mg film coated tablets | As an adjunct to diet & exercise to improve glycemic control in patients with type-two diabetes mellitus | 28-4-2008 |
| 115. | Lansoprazole for injection 30mg/vial | For short term treatment of erosive esophagitis when patients are unable to take the oral formulation | 28-4-2008 |
| | | | |

| 116 | I | | |
|------|---|--|-----------|
| 116. | Naproxen Suspension 125mg/5ml | For the treatment of RA in children 5 years of age and above | 28-4-2008 |
| 117. | Sevelamer Carbonate 400mg/800mg tablets & sachet 400mg/800 mg | For control of serum phosphorus in patients with chronic kidney disease on dialysis | 28-4-2008 |
| 118. | Rabeprazole Sodium (With Sodium bicarbonate as buffer) 10mg/20mg tablets | For the treatment of gastric ulcer, duodenal ulcer, Zollinger Ellison Syndrome and GERD | 28-4-2008 |
| 119. | Pantoprazole Sodium (With Sodium bicarbonate as buffer) 20mg/40mg tablets | For the treatment of gastric ulcer, duodenal ulcer, Zollinger Ellison Syndrome and GERD | 28-4-2008 |
| 120. | Cefdinir 250mg dispersible tablets and oral suspension 250mg/5ml | For the treatment of community acquired pneumonia, acute exacerabation of chronic bronchitis, uncomplicated skin and skin structure infection, Pharyngitis/tonsillitis | 29-4-2008 |
| 121. | Pregabalin capsules (Additional Indication) | Management of fibromyalgia syndrome | 29-4-2008 |
| 122. | Pregabalin 75mg/150mg + Methylcobalamin 750mcg capsules (Additional Strength) | Same as Approved | 8-5-2008 |
| 123. | Aliskiren 150/150/300/300mg+HCTZ 12.5/25/12.5/25mg tablets | For the treatment of essential hypertension in adult patients whose blood pressure is not adequately controlled on monotherapy | 8-5-2008 |
| 124. | Desmoprissionlyophilisate Oral tablets 60mcg/120mcg/240mcg per tablets | For diabetes insipidus and noctural emuresis | 8-5-2008 |
| 125. | Losartan Potassium 25mcg + Amlodipine (as Besylate) 5mg tablets (Additional Strength) | Same as Approved | 14-5-2008 |
| 126. | Lecithin Liquid (each 5 ml contains Lecithin 500mg equivalent to essential Phospholipids 250mg | For alcoholic liver disease, drug induced hepatitis | 14-5-2008 |
| 127. | Halobetasol 0.05% + Salicylic acid 6% Ointment (Additional Strength) | For the treatment of plaque psoriasis | 14-5-2008 |
| 128. | Benzoyl Peroxide Medicated Solution (each textured pad contains 4.5%/6.5%/8.5% of solution) | For the topical treatment of acne vulgaris | 14-5-2008 |
| 129. | Deflazocort 24mg tablets (Additional Strength) | Same as Approved | 14-5-2008 |
| 130. | Sodium Chondroitin sulphate 40mg + Sodium Hyaluronate 30mg per ml Intraocular Injection (Additional Strength) | For use as a surgical aid in anterior segments procedure including cataract extraction & intra-ocular lens implantation | 14-5-2008 |
| 131. | Esomeprazole (with Sodium bicarbonate as buffer) 20mg/40mg tablets | For the treatment of GERD, gastric and duodenal ulcer, Zollinger-Ellison syndrome | 14-5-2008 |
| 132. | Telmisartan 40mg + Atorvastatin (as Calcium) 10 mg Tablets | For coexisting essential hypertension & hyperlipidemia | 22-5-2008 |
| 133. | Milnacipran HCl 25mg/50mg equivalent to Milnacipran free base 21.77 or 43.55 mg tablet | For the treatment of adult patients with major depressive disorder | 22-5-2008 |
| 134. | Metolazone 2.5mg + Losartan Potassium 25mg/50mg tablets | For hypertension in patients whose blood pressure is not adequately controlled in monotherapy | 28-5-2008 |
| 135. | Metolazone 2.5mg + Ramipril 2.5mg/5mg | For essential hypertension in adult patients whose blood pressure is not adequately controlled on monotherapy | 28-5-2008 |
| 136. | Doxercalciferol 0.5mcg/2.5mcg Soft Gelatin capsules | For the treatment of secondary hyperthyroidism in patients with chronic kidney disease on dialysis | 29-5-2008 |
| 137. | Acebrophylline Capsules 100mg | For the treatment of adult patients with chronic obstructive pulmonary disease and bronchial asthma | 29-5-2008 |
| 138. | Sofalcone Capsule (100mg/50mg) | For the treatment of gastric ulcer and management of gastric mucosal lesions (erosion, haemorrhage, redness or edema) associated with acute gastritis or acute exacerbation of chronic gastritis | 2-6-2008 |

| 139. | S(+) Etodolac ER tablets 200mg/300mg | For the relief of pain and inflammation in patients with rheumatoid arthritis and osteoarthritis | 2-6-2008 |
|------|---|--|-----------|
| 140. | Diclofenac Sodium 50mg+Codeine Phosphate 50mg tablets | For the treatment of adult patients with signs & symptoms of acute painful conditions | 2-6-2008 |
| 141. | Sodium Hyaluronate injectable gel (each 0.6 ml 1 ml pre-filled syringe contains Sodium Hyaluronate 20mg/ml, 22.5mg/ml | For intradermal implantation and facial soft tissue augmentation | 3-6-2008 |
| 142. | Amphotericin B Lipid suspension injection (25 mg/vial) with lipid suspension | For invasive fungal infection in patients who are refractory to or intolerant of conventional amphotericin B therapy | 3-6-2008 |
| 143. | Dexrabeprazole 10mg (EC) + Domperidone 30mg SR Capsules | For the treatment of Gastroeasophageal Reflux Disease (GERD) in patients not responding to dexrabeprazole alone | 09-6-2008 |
| 144. | Thiocolchicoside 4mg/8mg + Ketoprofen 50mg/100mg tablets | For acute low back pain & other musculoskeletal painful and inflammatory condition in adults | 11-6-2008 |
| 145. | Cefotaxime sodium + Sulbactam sodium for injection (Additional Indication) | For the treatment of lower respiratory tract infection in children | 24-6-2008 |
| 146. | Artemether 40mg/80mg+ Lumefantrine 240mg/480mg tablets (Additional Strength) | Same as Approved | 24-6-2008 |
| 147. | Ivabradine (HCl) 5mg/7.5mg Tablets | For the treatment of chronic stable anginapectoris in patients with normal sinus rhythm who have a contraindication or intolerance for Beta-blockers | 3-7-2008 |
| 148. | Amino Acid, Glucose Parentral Solution (Solution for intravenous infusion) | Indicated to meet daily requirement of nitrogen (amino acids), energy (glucose) for patients needing parentral Nutrition | 8-7-2008 |
| 149. | Temoporfin 1mg/ml Injection (Additional Strength) | Same as Approved | 8-7-2008 |
| 150. | Pazufloxacin Mesylate i.v. infusion 300mg/100ml & 500mg/100ml | For the treatment of RTI, UTI and abdominal infections in adults | 9-7-2008 |
| 151. | Metoprolol ER + Amlodipine tablets (Additional Indication) | For coronary artery disease/ischaemic heart disease | 16-7-2008 |
| 152. | Fenofovir Disoproxil fumarate 300mg + Lamivudine 300mg + Efavirenz 600mg tablets | | 18-7-2008 |
| 153. | Vildagliptin 50mg/50mg/50mg + Metformin HCl +850 mg + 1000mg tablets | For the treatment of type-II diabetes mellitus when single drug therapy along with diet, exercise do not result in adequate glycemic control | 21-7-2008 |
| 154. | Testosterone Undecanoate 250mg/ml Injection | For Testosterone replacement therapy of male hypogonadism when testosterone deficiency has been confirmed by clinical feature or biochemical tests | 22-7-2008 |
| 155. | Abatacept powder for injection 250mg/vial | For adult patients with severe active rheumatoid arthritis who had an inadequate response to one more DMARDs such as methotixate or TNF antagonists | 22-7-2008 |
| 156. | Actarit 100mg Tablets | For the treatment of rheumatoid arthritis patients | 22-7-2008 |
| 157. | Tenofovir Fumarate 300mg+Lamivudine 300mg+Efavirenz 600mg Tablets | For the treatment of HIV infection in adults | 23-7-2008 |
| 158. | Bacillus Clausii Spores (2 Billion) Capsules (Additional dosage form) | Same as Approved | 23-7-2008 |
| 159. | Gatifloxacin 0.3% + Ketorolac Tramthamine 0.4% Eye Drop | For reduction of part operative inflammatory condition of eyes | 24-7-2008 |
| 160. | Ketorolac Tromethamine 4mg+Gatifloxacin 3mg Eye Drops | For the reduction of post-operative inflammatory conditions of the eye | 24-7-2008 |
| _ | | | |

| 161. | Cefadroxyl (as Monohydrate) 125 mg + Clavulanic acid (as Potassium) 31.25 per 5ml Syrup (Additional Strength) | Same as Approved | 25-7-2008 |
|------|---|--|-----------|
| 162. | Doxercalciferol (0.5/2.5mcg) Soft Gelatin Capsule (Additional Indication) | For the treatment of secondary hyperparathyroidism in patients with Stage 3 or Stage 4 chronic kidney disease | 5-8-2008 |
| 163. | Artemether + Lumefantrine Dispersible tablet (40/80mg + 240/480mg) (Additional Strength) | Same as approved | 5-8-2008 |
| 164. | Ceftazidime + Tobramycin injection (1gm + 120mg) (Additional Strength) | Same as approved | 5-8-2008 |
| 165. | Olopatadine Nasal Spray (0.6%) | For the relief of the symptoms of seasonal allergic rhinitis in patients 12 years of age and older | 5-8-2008 |
| 166. | Loxoprofen Sodium tablet (60mg) | For the symptomatic treatment of pain and inflammation condition in patients with osteoarthritis | 14-8-2008 |
| 167. | Dexibuprofen Suspension 50mg/5ml | For the treatment of mild to moderate pain, reduction of fever | 14-8-2008 |
| 168. | Bromfenac Sodium Eye Drop (0.09%) | For the treatment of post-operative inflammation & reduction of post-operative ocular pain in patients who have undergone cataract extraction | 14-8-2008 |
| 169. | Ceftriaxone + Sulbactam injection (Additional Indication) | For the treatment of acute bacterial otitis media, skin & soft structure infections, UTI, uncomplicated gonorrhoea, pelvic inflammatory disease, bacterial septicemia, bone and joint infections, intra abdominal infections, meningitis, surgical prophylaxis | 18-8-2008 |
| 170. | Citicoline Chewable tablet (500mg) | For the treatment of disturbance of consciousness as resulting from head injuries, brain operation and acute stage of cerebral infraction in adult only | 18-8-2008 |
| 171. | Ambroxol + Loratadine + Guaifenesin Syrup (30mg + 5mg + 50mg per 5ml) | For the treatment of productive cough | 18-8-2008 |
| 172. | Ferrous Ascorbate + Folic Acid + Zinc FC Tablets (100mg + 1.5 mg + 22.5 mg) | For the treatment of iron deficiency anaemia | 18-8-2008 |
| 173. | Cyclobenzaprine 7.5mg tablets (Additional Strength) | Same as approved | 18-8-2008 |
| 174. | S-Metoprolol Succinate Oral Solution (Each ml contains: | For the treatment of haemo-dynamically stably patients | 19-8-2008 |
| | S-Metoprolol Succinate 9.5mg equal to S-Metoprolol Tartrate 10mg) | with definite or suspected acute Myocardial Infarction to reduce cardio-vascular mortality. | |
| 175. | Doxorubicin HCl Pegylated Liposomal Injection (20/50MG) (Additional Indication) | In combination with Bortezumab for the treatment of progressive multiple myeloma in patients who have received at least one prior therapy and who have already undergone or are unsuitable for bone marrow transplant | 23-8-2008 |
| 176. | Povidone Iodine sachet (Containing Effervescent Granules). Each 2gm sachet contains: | As mouthwash for oral hygiene in various oropharyngeal conditions, surgery, dental infections, and halitosis | 25-8-2008 |
| | 175mg (Available Iodine 17.5mg) | | |
| 177. | Sodium Chondroitin Sulphate + Sodium Hyaluronate injection (40mg + not more than 17mg per ml) | Same as approved | 25-8-2008 |
| 178. | Metolazone + Spironolactone FC Tablets (2.5/5mg + 50mg) | For the treatment of oedema associated with congestive heart failure, cirrhosis and nephritic syndrome | 25-8-2008 |
| 179. | Tretinoin Microsphere Gel (0.1% w/v) | For topical treatment of acne vulgaris | 26-8-2008 |
| 180. | Misoprostol 50mcg tablet (Additional Strength) | Same as approved | 26-8-2008 |
| 181. | Ferrous Ascorbate + Folic Acid suspension (Equiv. to elemental | Same as approved | 27-8-2008 |

| 28-8-2008 2-9-2008 tatic colorectal 2-9-2008 ection 8-9-2008 the treatment of ed breast cancer acycline and a sistant and for intra-indicated c manifestation dermatoses like 13-9-2008 |
|--|
| tatic colorectal 2-9-2008 ection 8-9-2008 the treatment of ed breast cancer acycline and a sistant and for intra-indicated c manifestation dermatoses like |
| the treatment of ed breast cancer acycline and a sistant and for intra-indicated c manifestation dermatoses like |
| the treatment of ed breast cancer acycline and a sistant and for intra-indicated c manifestation dermatoses like |
| ed breast cancer acycline and a sistant and for antra-indicated c manifestation dermatoses like |
| dermatoses like |
| caemia tetanies 13-9-2008 |
| niry cattle. |
| n 15-9-2008 |
| muscular pain 15-9-2008 |
| hypertension in 15-9-2008 not adequately |
| 15-9-2008 |
| s 15-9-2008 |
| |
| rerebral injuries 16-9-2008 vithout loss of rative damages ile dementia. |
| nic treatment of 16-9-2008 fage & older |
| ergic rhinitis in er & perennial onths of age & |
| inflammatory 16-9-2008 |
| 16-9-2008 |
| hizophrenia in 26-9-2008 |
| nent of 26-9-2008 |
| disease (BRD, vith Mannhenia and Histophilus lactating dairy |
| |

| 201. | Ceftiofur HCl Suspension (As HCl salt) 500mg (veterinary) | For treatment of sub-clinical mastitis in dairy cattle at the time of dry off associated with Staphylococcus aureus, Streptococcus dysgalactiae, and Streptococcus uberis | 26-9-2008 |
|------|---|---|------------|
| 202. | Ceftiofur HCl Suspension (as HCl salt) 125mg (veterinary) | For treatment of clinical mastitis in lactating dairy cattle asssociated with coagulase- negative staphylococci, Streptococcus dysgalactiae, and Escherichia coli | 26-9-2008 |
| 203. | Nitrofurantoin SR Tab. 100mg | For the treatment of acute uncomplicated urinary tract infections (acute cystitis) caused by susceptible strains of Escherichia coli or Staphylococcus saprophyticus | 26-9-2008 |
| 204. | Tolperisone HCl Tablet 50/100/150 mg | For the relief of painful muscle spasms of the skeletal musculature | 26-9-2008 |
| 205. | Etoricoxib+PCM (60mg+500mg) Tablets | For the short term use in acute painful and inflammatory condition | 30-9-2008 |
| 206. | Citicoline Oral Drops | For the treatment of disturbance of consciousness resulting from head injuries, brain operation & acute stage of cerebral infraction in adult | 30-9-2008 |
| 207. | Paracetamol + Meloxicam + Benzyl Alcohol + Lignocaine (5mg/ml + 150mg/ml + 1% + 1%) Injection | Indicated for the treatment of mastitis, metritis, pneumonia, pleuritis, fever and relieving of acute pain in musculoskeletal disorders in animals | 30-9-2008 |
| 208. | Calcium (as Calcium Citrate Malate) + Colecalciferol (as Colecalciferol Concentrate) + Folic Acid (250mg + 100IU + 50mcg) Tablets | For the treatment of calcium and Vitamin D deficiency states (pregnancy, lactation, growing children) | 30-9-2008 |
| 209. | Artemether + Lumefantrine (80mg + 480mg/5ml) Dry syrup | For the treatment of chloroquine resistant/multidrug resitant on complicated malaria | 30-9-2008 |
| 210. | Cabergoline 1mg Tablets (Additional indication) | For the treatment of hyperprolactinemia and inhibition of lactation | 1-10-2008 |
| 211. | Pemetrexed disodium 500 mg/100 mg powder for sol. (Additional Indication) | In combination with cisplatin is indicated for the first line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology | 8-10-2008 |
| 212. | Tylvalosin tartrate oral soluble granules sachet 62.5% w/w (Veterinary) | For the prevention and treatment of mycoplasma induced chronic respiratory disease in broilers and breeders, the prevention and treatment of diseases due to clostridium perfringens (necrotic enteritis, wet litter, cholangiohepatitis), the prevention and treatment of ornithobacterium rhinotacheale (ORT) | 8-10-2008 |
| 213. | Bortezomib inj.3.5mg/vial (Additional Indication) | Treatment of patients with multiple myeloma | 9-10-2008 |
| 214. | Sertaconazole Nitrate Vaginal Tab.500mg | For the treatment of vulvo-vaginal candidiasis in women | 9-10-2008 |
| 215. | Fosfomycin Tromethamine sachet 3 gms | For the treatment of uncomplicated urinary tract infections (acute cystitis) in women due to susceptible strains of E.Coli and Entreococcus faecalis | 9-10-2008 |
| 216. | Ixabepilone lyophilized powder (Additional Indication) | As monotherapy for the treatment of metastatic or locally advanced breast cancer in patients whose tumors are resistant or refractory to cytotoxic chemotherapy. Previous therapy should have included as anthracycline, a taxane, and capecitabine | 9-10-2008 |
| 217. | Eprosartan Mesylate 400 mg/600 mg Tablets | For the treatment of essential hypertension | 11-10-2008 |
| 218. | Tretinoine Gel as Micro Sphere 0.04% 0.1% | For topical treatment of Acne vulgaris | 14-10-2008 |
| 219. | Hydroxychloroquine Sulphate (Additional Indication) | For the treatment of systemic lupus erythematosu | 17-10-2008 |

| 220. | Budesonide + Formoterol (80/160 mcg + 4.5mcg) Inhalational Powder (Additional Indication) | Indicated in the regular treatment of patients with moderate to severe COPD with frequent symptoms and a history of exacerbations | 17-10-2008 |
|------|---|--|------------|
| 221. | Azithromycin + Dexamethasone (10mg + 1mg) Eye Drops | For post-operative steroid responsive inflammatory ocular condition when bacterial ocular infections or a risk of bacterial ocular infections exists. | 21-10-2008 |
| 222. | Diclofenac + Thiocolchicoside (50mg (EC) + 8mg) Capsules | For the treatment of acute inflammation conditions associated with spasm in adults only | 21-10-2008 |
| 223. | Adapalene 0.1% + Benzoyl Peroxide 2.5% gel | For the treatment of acne vulgaris | 24-10-2008 |
| 224. | Aceclofenac Roll - On 1% | For the treatment of acute musculoskeletal pain in patients | 24-10-2008 |
| 225. | Citicoline Troches 200/500mg | For the treatment of patients with serious cerebral injuries of vascular traumatic nature with or without loss of consciousness and for treat-ment of degenerative damages and chronic cerebral vascular injuries in senile dementia | 31-10-2008 |
| 226. | Amisulpride S.R.Bilayered tablet 200/400mg | For acute and chronic schizophrenic disorders, in which positive symptoms and or negative symptoms are prominent, including patients characterised by predominant negative symptoms | 3-11-2008 |
| 227. | Calcitriol + Calcium + Zinc (0.25mcg + 200mg + 7.5mg) Capsules | For the treatment of hypocalcaemia and/or osteoporosis | 4-11-2008 |
| 228. | Atazanavir + Ritonavir (300mg + 100mg) FC Tablets | For the treatment of patients with HIV infection | 4-11-2008 |
| 229. | Timolol Maleate SR Eye Drops 5.0 mg/ml | For the treatment of elevated intraocular pressure in patients with ocular hypertension or open angle glaucoma | 8-11-2008 |
| 230. | Rosuvastatin Calcium tablets 5/10/ 20/40 mg (Additional Indication) | (1) As an adjunctive therapy to diet for the treatment of adult patients with hypertriglyceridemia. (2) As an adjunctive therapy to diet to slow the progression of atherosclerosis in adult patients as part of a treatment strategy to lower Total-C and LDL-C to target levels | 8-11-2008 |
| 231. | Paclitaxel inj.30/100/300mg vial (Additional Indication) | For the treatment of advanced gastric carcinoma | 8-11-2008 |
| 232. | Montelucast + Desloratidine(10mg + 5mg) FC Tablets | For the treatment of allergic rhinitis in adults only | 10-11-2008 |
| 233. | Amlodipine + Metoprolol (2.5/5 mg + 25/50mg (ER))Tablets | Additional Lower Strength | 10-11-2008 |
| 234. | Losartan + HCTZ (100 mg + 25 mg) Tablets | Additional Strength | 10-11-2008 |
| 235. | Phenyramidol HCl tablet 400mg | For the treatment of lumbago, integumental pain, musculoskeletal pain | 14-11-2008 |
| 236. | Salicylic Acid + Coal Tar (3% + 1%) Solution | For scaling of seborrhea, seborrheic dermatitis, psoriasis of scalp | 17-11-2008 |
| 237. | Salicylic Acid + Coal Tar + Precipitated Sulphur (3% + 1% + 3%) Ointment | For the treatment of psoriasis and fungal skin infection | 17-11-2008 |
| 238. | Thiocolchicoside + Etoricoxib(4mg + 60 mg) FC Tablets | For the acute treatment of inflammatory musculoskeletal disorders associated with painful muscle spasm in adults | 17-11-2008 |
| 239. | Tamsulosin + Tolterodine (0.4mg MR Pellets + 2mg/4mg ER Pellets) Capsules | Additional Lower Strength | 17-11-2008 |
| 240. | Calcium Hypochlorite 4gm Sachet (Calcium Hypochlorite | For purification of contaminated ground water sources | 23-11-2008 |
| | 0.542% Other Ingredients: 99.458%) | | |

| 242. | Amoxycillin + Clavulanic acid(1gm+62.5mg) FC Tablets | For the treatment and management of patients with community acquired pneumonia or acute bacterial sinusitis due to beta lactamase producing pathogens | 25-11-2008 |
|------|--|---|------------|
| 243. | Buprenorphine + Naloxone (2 mg/0.4mg + 0.5/0.1 mg) Tablets | Additional Lower Strength | 2-12-2008 |
| 244. | Atorvastatin + Metformin(10mg + 500mg SR) Tablets | As adjunctive therapy to diet and exercise in patients with diabetic dyslipidemia | 4-12-2008 |
| 245. | Pemetrexed disodium 500 mg/100 mg powder for solution (Additional Indication) | As monotherapy for the second-line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology | 4-12-2008 |
| 246. | Magnesuim Valproate Tablet 200mg/400mg | For the treatment of generalized or partial seizures as monotherapy or adjunctive therapy. For the treatment of maniac episodes associated with bipolar disorders. For prophylaxis of migraine headache in adults | 4-12-2008 |
| 247. | Ursodeoxycholic Acid 300/600 SR Tablets (Additional Strength) | For the dissolution of Radiolucent collestral goll stone, chronic cholestatic liver disease in particular primary biliary cirrhosis, primary sclerosing cholangitis & cholestasis associated with cyatic fibrosis | 8-12-2008 |
| 248. | Rifampicin + INH + Piperine (200mg + 300mg + 10mg) Capsules | For the treatment of adult patients with pulmonary tuberculosis | 11-12-2008 |
| 249. | Rifampicin +Piperine (200mg + 10mg) Capsules | For the treatment of adult patients with pulmonary tuberculosis | 11-12-2008 |
| 250. | Amphotericin B Lipid suspension 50mg/ vial (Additional Strength) | Additional Strength | 12-12-2008 |
| 251. | Desmopressin Acetate tablet 0.1/0.2 mg | For the treatment of cranial diabetes insipidus, primary noctumal enuresis in patients. | 12-12-2008 |
| 252. | Mifepristone + Misoprostol (1 tab of mifepristone 200mg and 4 tab of misoprostol 200mcg) Combipack | For the medical termination of intrauterine pregnancy (MTP) of up to 63 days gestation based on the first day of the last menstrual period | 16-12-2008 |
| 253. | Lamotrigine MR Tablets 50/100 mg | For the treatment of bipolar I disorder (Additional Indication) | 17-12-2008 |
| 254. | Levosulpride Injection 25mg/2ml (Additional Strength) | For the treatment of different gastro-intestinal problems like functional dyspepsia, nausea, vomiting & diabetic gastroparesis. For the treatment of schizophrenia and depression in adults | 17-12-2008 |
| 255. | Cyclobenzaprine HCl ER Capsule 15/30mg | As an adjunct to rest and physical therapy for relief of muscle spasm associated with acute, painful, musculoskeletal conditions | 17-12-2008 |
| 256. | Docetaxel injection concentrate 80mg/2ml, 20mg/0.5ml (Additional Indication) | In combination with cisplatin and 5-fluorouracil is indicated for the induction treatment of patients with locally advanced squamous cell carcinoma of the head and neck | 17-12-2008 |
| 257. | Salicylic Acid + Cetyl Alcohol + Stearyl Alcohol (2% + 2.65% + 0.26%) Lotion | For the treatment of scaly dermatoses | 18-12-2008 |
| 258. | Azathioprine 25mg Tablets | Additional Strength | 19-12-2008 |
| 259. | Paracetamol solution for infusion 10mg/ml | For short term treatment of moderate pain, especially following surgery and for the short term treatment of fever, when administration by i.v route is clinically justified | 23-12-2008 |
| 260. | Gemifloxacin + Ambroxol (320mg + 75mg SR) FC Tablets | Indicated for the treatment of lower respiratory tract infections in adult patients | 24-12-2008 |

| 261. | Ceftazidime + Sulbactam (500/1000mg + 250/500mg) Injection | 1. For LRTI infections, | 24-12-2008 |
|------|--|--|------------|
| | | 2. For UTI - Complicated and uncomplicated, | |
| | | 3. For bacterial septicemia, | |
| | | 4. For bone and joint infections, | |
| | | 5. For gynaecologic infections including endometritis, pelvic cellutitis and other infections of female genital tract, | |
| | | 6. For CNS infections including meningitis | |
| 262. | Paracetamol + Phenylephrine Hydrochloride (Each 6gm sachet contains: | For the treatment of fever associated with cold, nasal congestion in adults | 24-12-2008 |
| | 500mg + 10mg) | | |
| 263. | Carboinoxamine Maleate + Ammonium Chloride + Sodium Citrate (Each 5 ml contains: | For symptomatic treatment of cough | 24-12-2008 |
| | 2 mg + 120 mg + 120 mg) syrup | | |
| 264. | Diclofenac + Thiocolchicoside (50 mg +4 mg) Capsules | Additional Lower Strength | 24-12-2008 |

List of Drugs approved during January to December 2009

| Sl. No. | Name of Drug | Indication | Date of Approval |
|---------|--|---|------------------|
| 1. | Zoledronic Acid Injection 5 mg/100 ml (Addl. Indication) | For the treatment of osteoporosis in post-menopausal women and in men who are at increased risk of fracture, including those with a recent low-trauma hip fracture | 7-1-2009 |
| 2. | Misoprostol Tablet 600 mcg | Addl. Strength | 9-1-2009 |
| 3. | Acebrophylline Syrup 10 mg/ml | For the treatment of adult patients with chronic obstructive pulmonary disease (COPD) and bronchial asthma | 9-1-2009 |
| 4. | Selamectin Tropical solution (For verterinary use) 60 mg/120 mg per ml | (I) For the treatment and lasting prevention of flea infestations, treatment and control of flea allergy dermatitis, prevention of heartworm disease, treatment and prevention of ear mites in dogs and cats; | 9-1-2009 |
| | | (II) For the treatment and prevention of sarcoptic mange, roundworms and for control of tick infestations in dogs | |
| 5. | Methylcobalamin 1500 mcg + Calcium 200 mg + Calcitriol 0.25 mcg + Folic Acid 1.5 mg + Pyridoxine 3 mg Capsules | For the treatment of vitamin deficiency states in adults | 14-1-2009 |
| 6. | Metoprolol tartrate tablets 50 mg (Addl. Indication) | Adjunct to the treatment of Hyperthyroidism | 14-1-2009 |
| 7. | Cinacalet (as HCl) Tablet 30/60/90 mg | For the treatment of secondary hyper parathyroidism in patients with chronic kidney diseases (CKD) on dialysis and for hypercalcemia in patients with parathyroid carcinoma | 14-1-2009 |
| 8. | Mycophenolate mofetil Capsules 250 mg, Tablets 500 mg, Oral suspension 200 mg/ml (Addl. Indication) | For the prophylaxis of acute organ rejection in patients recieving allogenic hepatic transplantation | 23-1-2009 |
| 9. | Liquid Paraffin 1.25 ml + Milk of Magnesia 3.75 ml + Sodium Picosulphate 3.33 mg per 5 ml | For symptomatic treatment of constipation in adults | 29-1-2009 |
| 10. | Ranolazine 1000 mg ER Tablets (Additional Strength) | Same as approved | 31-1-2009 |
| 11. | Hyaluronic acid (cross linked) Pre-filled syringe 18 mg/0.6 ml and 24 mg/0.8 ml | For intradermal implantation and facial soft tissue augmentation | 31-1-2009 |
| 12. | Caspofungin Acetate Injection (Addl. Indication) | For use in following: | 31-1-2009 |
| | | (I) Empirical therapy for presumed fungal infections in febrile neutropenic patients | |
| | | (II) Treatment of Candidemia and the following Candida | |

| | T | | |
|-----|--|--|-----------|
| | | infections: Intra-abdominal abscesses, Peritonitis and Pleural space infections. | |
| | | (III) Treatment of Esophageal Candidiasis. | |
| | | (IV) Treatment of invasive Aspergillosis in patients who are refractory to or intolerant of other therapies (i.e., amphotericin B, lipid formulations of amphotericin B, and/or itraconazole) | |
| 13. | Micafungin (as sodium) Powder for solution for infusion 50 mg/vial | For the treatment of patients with candidemia, acute disseminated candidiasis, candida peritonitis, abscess and esophageal candidiasis. For the prophylaxis of Candida infection in patients undergoing hematopoietic stem cell transplantation (HSCT) | 31-1-2009 |
| 14. | Loteprednol Etabonate Nasal spray 0.1% w/v | For prophylaxis and treatment of the nasal symptoms of seasonal allergic rhinitis | 31-1-2009 |
| 15. | Rasagiline (as mesylate) Tablet 0.5 mg/1 mg | For the treatment of signs and symptoms of Idiopathic Parkinson's disease as initial monotherapy and as an addition to Levodopa | 31-1-2009 |
| 16. | Nicotine Polacrilex Gum 2 mg & 4 mg (Addl. Indication) | To reduce the withdrawal symptoms, including nicotine craving, associated with quitting the smoking and quitting chewed tobacco and guthka containing tobacco | 3-2-2009 |
| 17. | Closantel Injection 100 mg/ml (for verterinary use) | For the infestation of trematodes, nematodes, cestodes and arthropods of domestic animals, including milch and other food producing animals | 5-2-2009 |
| 18. | Lornoxicam 4 mg + Paraceta-mol 500 mg film coated tablets | For the short term use in acute painful & inflammatory conditions in adults | 5-2-2009 |
| 19. | Cetrimide 15% w/v + Chlorhexidine Gluconate 7.5% v/v + Isopropyl Alcohol 7.5% v/v Concentrate Solution | For cleansing of physically contaminated wounds and for pre-operative disinfection | 9-2-2009 |
| 20. | Ketoconazole 2% w/v + Prepared Coal Tar 2% w/v (I.P. 1966) Solution | For the treatment of dandruff | 9-2-2009 |
| 21. | Doripenem (as monohydrate) Injection 250 mg/500 mg | For the treatment of nosocomial pneumonia in adults | 16-2-2009 |
| 22. | Halobetasol Scalp Approved 0.05% w/v | For the relief of inflammatory and puritic manifestation of corticosteroid respensive dermatosis | 18-2-2009 |
| 23. | Nicotine (as Polacrilex) Lozenges 2 mg & 4 mg | To reduce the withdrawal symptoms, including nicotine craving, associated with quitting the smoking and quitting chewed tobacco and guthka containing tobacco | 18-2-2009 |
| 24. | Tranexamic Acid Tab SR 750 mg | For the treatment of menorrhagia | 20-2-2009 |
| 25. | Betahistine Mesilate Tablet 6 mg/12 mg | For the treatment of dizziness and feeling of dizziness resulting from Meniere's disease, Meniere's syndrome and vertigo | 20-2-2009 |
| 26. | Abatacept Injection 250 mg/vial (Addl. Indication) | In moderately to severly active rheumatoid arithritis in adults and may be used as monotherapy or concomitantly with DMARD's other than TNF antagonists | 20-2-2009 |
| 27. | Anidulafungin Injection (Lyophilized powder) 50 mg/ 100 mg per vial | For the treatment of invasive candidiasis including candidemia and esophageal candidiasis in adult patients | 23-2-2009 |
| 28. | Hydroquinone 2%+ Tretinoin 0.05% + Hydrocortisone 1% Cream | For the treatment of patients with skin hyperpigmentation | 23-2-2009 |
| 29. | Doxyphylline 400 mg + Terbutaline 5 mg tablets | For the treatment of asthma and chronic obstructive pulmonary disease in adult patients only | 23-2-2009 |
| 30. | Hydroquinone 20 mg + Tretinoin 0.25 mg + Fluocinolone Acetonide 0.1 mg per gm Cream | For the treatment of melasma of face in adults | 23-2-2009 |
| | I . | | |

| 31. | Ketoconazole 2% + Zinc Pyrithone 1% Lotion | For the treatment of pityriasis and dandruff of the scalp | 23-2-2009 |
|-----|---|--|-----------|
| 32. | Lamivudine 30 mg + Zidovudine 60 mg + Nevirapine 50 mg Oral Suspension (Tablets for Oral Suspension) | For the treatment of HIV patients | 23-2-2009 |
| 33. | Glibenclamide 5 mg + Pioglitazone 15 mg + Metformin 500 mg film coated tablets | As 3rd line treatment of Type II Diabetes mellitus when diet, exercise and the single agents and the second line therapy with two drugs do not result in adequate glycemic control | 23-2-2009 |
| 34. | Aceclofenac 100 mg + Tizanidine 2 mg film coated tablets | For the treatment of acute low back pain in adult patients only | 23-2-2009 |
| 35. | Octinoxate 7.5% + Avobenzone 3% + Oxybenzone 3% + Titanium dioxide 2% lotion | As sunscreen lotion for prevention of sunburn skin | 23-2-2009 |
| 36. | Cetrizine 5/10 mg + Ambroxol 30 mg per 5 ml syrup | For the symptomatic relief of productive cough associated with allergic rhinitis, when both anti-histamine and mucolytic agents are desired | 23-2-2009 |
| 37. | Lidocaine Lozenges 200/300 mg | For single agent anesthesia in upper gastro intestinal endoscopy | 26-2-2009 |
| 38. | Calamine 8% + Liquid Paraffin 10% lotion | As antipruritic and smoothening agent | 26-2-2009 |
| 39. | Nifedipine 0.3% + Lidocaine 1.5% cream | For the treatment of anal fissures | 26-2-2009 |
| 40. | Quinapril 20 mg + HCTZ 25 mg film coated tablet | For the treatment of mild to moderate hypertension in adult patients (in whom combination therapy is appropriate) who have been stabilized on the individual components given in the same proportion | 26-2-2009 |
| 41. | Levodopa 100/100/250 + Carbidopa 10/25/25 mg mouth dissolving tablet | For the symptomatic symptoms of idiopathic Parkinson's Disease | 4-3-2009 |
| 42. | Paliperidone ER Tablets 1.5 mg | Additional Strength | 4-3-2009 |
| 43. | Losartan 100 mg+ HCTZ 25 mg tablet | Additional higher strength | 16-3-2009 |
| 44. | Darunavir (as ethanolate) Tablet 300 mg | For the treatment of human immunodeficiency virus (HIV) infection in antiretroviral treatment experienced adult patients, such as those patients with HIV-1 strains resistant to more than one protease inhibitor when co-administered with 100 mg ritonavir, and with other antiretroviral agents | 19-3-2009 |
| 45. | Ropivacaine HCl Inj. 2/5/7.5/ 10 mg | For the production of local or regional anasthesia for surgery and for acute pain management | 19-3-2009 |
| 46. | Butaphosphan 100g + Methyl Hydroxybenzoate 1gm + Cyanocobalamin 50 gm Oral powder (Veterinary Drug) | As vitamin and phosphorous supplement for livestock, shrimp and fish | 19-3-2009 |
| 47. | Fluticasone Furoate Nasal spray 27.5 mcg per spray | For the treatment of symptoms of allergic rhinitis | 19-3-2009 |
| 48. | Octinoxate 7.5% + Avobenzone 2% + Oxybenzone 3% + Octorylene 3%+Zinc Oxide 2% lotion | A sunscreen lotion for prevention of sunburn skin | 19-3-2009 |
| 49. | Omeprazole 20/40 mg + Sodium Bicarbonate 600 mg+ Magnesium Hydroxide 700 mg sachet | Additional Strength | 19-3-2009 |
| 50. | Ambroxol 30 mg + Levo Salbutamol 1 mg+ Guiphenesin 50 mg /5 ml syrup | For the symptomatic relief of bronchospasm in bronchial asthma & chronic bronchitis | 19-3-2009 |
| 51. | Montelucast 4 mg+ Levocetrizine 2.5 mg tablet | Additional lower paedriatric dose | 19-3-2009 |
| 52. | Surgical Sealant | To be indicated for use in sealing suture lines along arterial and venous reconstructions | 26-3-2009 |
| 53. | Lidocaine Patch 700 mg in aqueous base (50 mg per gram adhesive) | For the relief of pain associated with post-herpetic neuralgia. It should be applied only to intact skin | 26-3-2009 |
| 54. | Bimatoprost Opthalmic solution 0.03% | For the treatment of hypotrichosis of the eyelashes by increasing their growth including length, thickness and darkness | 30-3-2009 |

| 55. | Trabectedin powder for concentrate for solution for infusion 1 mg/vial | For the treatment of patients with advanced soft tissue sarcoma, after failure of anthracyclines and ifosfamide, or who are unsuited to recieve these agents | 30-3-2009 |
|-----|--|--|-----------|
| 56. | Acebrophylline Syrup 10 mg/ml (Additional Indication) | For the treatment of children (age 6 to 12 years) with chronic obstructive pulmonary disease (COPD) and bronchial asthma | 30-3-2009 |
| 57. | Mesalamine PR Tablets 1.2 gm | For the induction of remission in patients with active, mild to moderate ulcerative colitis | 30-3-2009 |
| 58. | Mycophenolate Mofetil Tablets 750 mg Additional Strength | Same as approved | 31-3-2009 |
| 59. | Ketorolac Tromethamine Gel 2% w/w | For the treatment of muscle, joint and post-traumatic stress | 31-3-2009 |
| 60. | Theophylline 400 mg SR + Montelukast 10 mg Tablet | For the treatment of patients with bronchial asthma | 31-3-2009 |
| 61. | Hyaluronic Acid Sodium Salt 0.2 gm + Silver Sulfadiazine 1.0 gm/25 gm tube Hyaluronic Acid Sodium Salt 50 mg + Silver Sulfadiazine 1.0 gm/100 gm gauze pads impregneted with cream | For the treatment of wound of different etiology | 31-3-2009 |
| 62. | Telmisartan 80 mg + Amlodipine 5 mg tablet | Additional higher strength | 31-3-2009 |
| 63. | Hyaluronic Acid Sodium Salt | For the treatment of wound of different etiology | 31-3-2009 |
| | (i) Each 15 gm tube contains: | | |
| | Hyaluronic Acid Sodium Salt 0.2 gm/100 gm | | |
| | (ii) Sterile Gauze Pads impregnated with cream: Each gauze pad contains Hyaluronic Acid Sodium Salt 50/100 mg | | |
| 64. | Abatacept Injection 250 mg/vial (Additional indication) | For reducing signs and symptoms of moderately to severly active polyarticular juvenile idiopathic arthritis/juvenile rheumatoid arthritis in pediatric patients 6 years of age and older who have had an inadequate response to one or more DMARD's such as MTX | 10-4-2009 |
| 65. | Cetrizine 10 mg + Ambroxol 75 mg SR Tablet | For symptomatic relief of productive cough associated with allergic rhinitis, when both anti-histamine and mucolytic agents are desired | 13-4-2009 |
| 66. | Ciclesonide 200 mcg + Formoterol 6 mcg + Tiotropium 9 mcg MDI | Indicated as third line treatment of severe cases of COPD, when monotherapy and second line therapy with two drugs do not respond adequately | 13-4-2009 |
| 67. | Each film coated bilayered tablet contains: Metoprolol 25 mg/50 mg + Chlorthalidone 12.5 mg | For the treatment of patients with mild to modrate essential hypertension | 13-4-2009 |
| 68. | Milnacipran 25/50 mg Caps (Addl. Indication) | For the management of fibromylagia | 15-4-2009 |
| 69. | Lornoxicam SR 16 mg Tabs | For the symptomatic treatment of pain and inflammation in patients of osteoarthritis & rheumatoid arthritis | 25-4-2009 |
| 70. | Palonosetron HCl 0.075 mg IV Inj (Addl. Strength) | Same as approved | 25-4-2009 |
| 71. | Each film coated bilayered tablet contains: Metoprolol 25 mg/50 mg + Chlorthalidone 6.25 mg/12.5 mg | Additional strength | 25-4-2009 |
| 72. | Albendazole 400 mg + Ivermectin 6 mg tablet | For the treatment of intestinal helminthes and suppression of microfilaraemia especially with bancrofti infections | 25-4-2009 |
| 73. | Decitabine Lyophilized powder for injection 50 mg/20 ml vial | For treatment of patients with myelodysplastic syndromes (MDS) including previously treated and untreated, de novo and secondary MDS of all French-American-British subtypes(refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, refractory | 25-4-2009 |

| | | anemia with excess blasts in transformation, and chronic myelomonocytic leukemia) and intermediate-1, intermediate-2, and high-risk international Prognostic Scoring System groups | |
|-----|---|--|-----------|
| 74. | Retapamulin Ointment 1% | For the topical treatment of bacterial skin and skin structure infections:— Primary impetigo, secondarily infected traumatic lesions, secondarily infected dermatoses | 25-4-2009 |
| 75. | Each ml contains Calamine 15% w/v + Zinc Oxide 5% w/v + Bentonite 3% w/v+ Glycerin 5% w/v + Sodium Citrate 0.05 % w/v lotion | For the treatment of pruritis and prickly heat | 8-5-2009 |
| 76. | Each 1000 ml of emulsion contains: Medium chain Triglycerides 100 gm + Soya oil 80 gm + Omega-3 Acid Triglycerides 20 gm + Egg Lecithin 12 gm + Glycerl 25 gm + Sodium Oleate 0.30 gm + Alpha-tocopherol 0.20 gm + Ascorbypalmitate 0.30 gm + Sodium Hydroxide max. 0.06 gm + water for injection 784 gm emulsion for infusion | For the supply of fats, including essential Omega-6 fatty acids and Omega-3 fatty acids, as part of a parenteral nutrition regimen in adults if oral or enteral nutrition is impossible, inadequte or contraindicated | 14-5-2009 |
| 77. | Sertaconazole Nitrate Powder 2% w/w & Lotion 2% w/v | For the topical treatment of the superficial fungal infections of the skin | 2-5-2009 |
| 78. | Amlodipine Tablets 5 mg/10 mg (Additional Indication) | To reduce of fatal coronary heart disease and non-fatal myocardial infarction, and to reduce the risk of stroke. To reduce the risk of coronary revascularization procedures and the need for hospitalization due to angina in patients with coronary artery disease | 2-5-2009 |
| 79. | Clindamycin Phosphate Inj. 150 mg/ml & Clindamycin HCl Capsules 150 mg/300 mg (Additional Indication) | Prophylaxis of infection in neck and head surgery | 2-5-2009 |
| 80. | Each 15 gm sachet contains: Lactulose 10 gm + Ispaghula husk 3.50 gm granules | For the treatment of chronic idiopathic constipation in adult patients only | 16-5-2009 |
| 81. | Sucralose | For use as sweetner in drug formulations | 9-5-2009 |
| 82. | Carvedilol Tablets 3.125/6.25/12.5/25 mg (Additional Indication) | To reduce cardiovascular mortality in clinically stable patients who have survived the acute phase of a myocardial infarction and have a left ventricular ejection fraction of \leq 40% (with or without symptomatic heart failure) | 9-5-2009 |
| 83. | Dexketoprofen 12.5 mg/25 mg + Paracetamol 500 mg film coated tablets | For the treatment of Acute musculoskeletal pain | 18-5-2009 |
| 84. | Pregabalin SR Tablet 150/300 mg | For the treatment of peripheral neuropathic pain in adults | 16-5-2009 |
| 85. | Guaiphenesin ER Tablet 1200 mg | For the treatment of productive cough accompanied by Phlegm | 16-5-2009 |
| 86. | Everolimus 5/10 mg Tablets (Addl. Indication) | Treatment of advance renal cell carcinoma | 16-5-2009 |
| 87. | Nimesulide ER 200 mg (Additional Indication) | Acute pain in Orthopaedic, ENT, Dental and Post-Operative condition | 16-5-2009 |
| 88. | Palonesetron Injection 0.25 mg/5 ml (Additional Indication) | For treatment of Post-Operative Nausea & vomiting | 16-5-2009 |
| 89. | Bendamustine Hydrochloride Lyophilized powder Injection 100 mg/vial | For the treatment of patients with chronic lymphocytic leukemia | 16-5-2009 |
| 90. | Caffeine (as Citrate) 10 mg/ml Injection | For the short term treatment of apnoea of prematurity of infants between 28 and < 33 weeks gestational age | 16-5-2009 |
| 91. | Perindopril Arginine 2.5 mg/ 5 mg/10 mg film coated tablet | For the treatment of arterial hypertension, congestive heart failure and coronary artery disease | 16-5-2009 |

| 92. | Each gm contains: Halobetasol 0.05% w/w + Mupirocin 2.0 % w/w gel | For the topical treatment of inflammatory dermatitis with secondry bacterial infection | 16-5-2009 |
|------|--|--|-----------|
| 93. | Mometasone 0.1 % + Fusidic acid 2% cream/ointment | For the treatment of dermatoses where secondry bacterial and/or candidial infection is present | 25-4-2009 |
| 94. | Thiocolchicoside 8 mg + Etoricoxib 60 mg tablets | Additional strength | 25-4-2009 |
| 95. | Methadone Hydrochloride 5 mg/ 10 mg per ml syrup; 5 mg/10 mg/20 mg/40 mg tablet | For the treatment of opiod dependence and in maintainence treatment for opioid dependence | 28-5-2009 |
| 96. | Trospium Chloride film coated tablet 20 mg | For the treatment of overactive bladder with the symptoms of urge urinary incontinence, urgency, and urinary frequency | 28-5-2009 |
| 97. | Arformoterol (as Tartrate) 15 mcg/2 ml Inhalation Solution | For the long term, twice daily (morning and evening) maintainence treatment of bronchoconstriction in patients with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema | 29-5-2009 |
| 98. | Zolmitriptan5.0% w/v Benzalkonium Chloride NF0.01% w/v Nasal spray | For the acute treatment of migraine with or without aura in adults | 29-5-2009 |
| 99. | Dexmedetomidine (as HCl) 100 mcg/ml Injection | For sedation of initially and mechanically ventilated patients during treatment in an intensive care setting | 29-5-2009 |
| 100. | Levetiracetam SR Tablet 500/750/1000 mg | Same as approved | 2-5-2009 |
| 101. | Glatiramer Acetate Injection (20 mg/ml) & Pre-filled syringe (20 mg/ml) | For the reduction of the frequency of relapses in patients with Relapsing-Remitting Multiple Sclerosis | 3-6-2009 |
| 102. | Yttrium-90 Microspheres Injection (Yttrium-903GBq (at the time of calibration) per vial) | For the treatment of unresectable metastatic liver tumors from primary colorectal cancer with intra-hepatic artery chemotherapy (IHAC) of FUDR (Floxuridine) | 3-6-2009 |
| 103. | Luliconazole Cream 1% | For the treatment of cutaneous mycosis viz. Tinea pedis, Tinea corporis, Tinea cruris | 3-6-2009 |
| 104. | Magnesium Valporate Prolonged release tablet 600 mg | (1) For the treatment of generalized or partial seizures as monotherapy or adjunctive threapy. | 4-6-2009 |
| | | (2) For prophylaxis of migraine headache in adults. | |
| | | (3) For the treatment of manic episodes associated with bipolar disorders | |
| 105. | Tenofovir Disoproxil Fumerate 300 mg | For the treatment of chronic Hepatitis B in adults | 19.6.2009 |
| 106. | Bosentan (as monohydrate) Tablets 62.5/125 mg | For the treatment of pulmonary arterial hypertension (WHO Group I) in patients with WHO Class III or IV symptoms, to improve excercise ability and decrease the rate of clinical worsening | 23.6.2009 |
| 107. | Lornoxicam 12 mg SR (Additional Strength) | Same as approved | 26.6.2009 |
| 108. | Azithromycin SR Granules for Oral Suspension 2000 mg | For the treatment for mild to moderate susceptible infection including RTI, uncomplicated skin/skin structure, non-gonococcal urethritis, cervicitis | 26.6.2009 |
| 109. | Metoprolol 50/50/25 mg + Atorvastatin 20/10/10 mg + Ramipril 10/5/2.5 mg capsules | As 2nd line treatment of patients with both essential hypertension and hypercholesterolemia who are not adequately controlled by combination of an antihypertensive with lipid lowering agent | 1-7-2009 |
| 110. | Methyl Phenidate ER Tablet 10/20/30 mg | For the treatment of Attention Deficit Hyperactivity Disorder (ADHD) | 10-7-2009 |
| 111. | Dasatinib Tablets 100 mg (Addl. Strength) | Same as approved | 10-7-2009 |
| 112. | Voglibose 0.2 mg + Metformin 500 mg Tablets | As 2nd line treatment of Type II Diabetes mellitus when diet, exercise and the single agent do not result in adequate glycemic control | 13-7-2009 |

| 113. | Combipack of Artesunate tablet and FDC Tablet of Sulphadoxine + Pyrimethamine Composition: | For the treatment of P. falciparum resistant malaria patients | 15-7-2009 |
|--|---|---|---|
| | A. Combipack of 1 strip of 3 tablets of artesunate 25 mg and 1 strip of 1 tablet of sulphadoxine 125 mg + pyrimethamine 6.25 mg | | |
| | B. Combipack of 1 strip of 3 tablets of artesunate 50 mg and 1 strip of 1 tablet of sulphadoxine 500 mg + pyrimethamine 25 mg | | |
| | C. Combipack of 1 strip of 3 tablets of artesunate 100 mg and 1 strip of 1 tablet of sulphadoxine 500 mg + pyrimethamine 25 mg | | |
| | D. Combipack of 1 strip of 3 tablets of artesunate 200 mg and 1 strip of 1 tablet of sulphadoxine 500 mg + pyrimethamine 25 mg | | |
| 114. | Milnacipran 100 mg (Additional Strength) | Same as approved | 16-7-2009 |
| 115. | Rifaximin 400 mg (Additional Indication) | For the treatment of hepatic encephalopathy | 18-7-2009 |
| 116. | Nevirapine tablet 50 mg for oral suspension mg | For the treatment of HIV infection in combination with other anti-retro viral agents | 18-7-2009 |
| 117. | Desvenlafaxine (as Succinate) ER Tablets 50 mg/100 mg | For the treatment of major depressive disorder (MDD) | 18-7-2009 |
| 118. | Posaconazole Oral Suspension 40 mg/ml | For the treatment of oropharyngeal candidiasis, including oropharyngeal candidiasis refractory to itraconazole and/or fluconazole. For prophylaxis of invasive Aspergillus and Candida infections in patients, 13 years of age and older, who are at high risk of developing these infections due to being severly immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy | 18-7-2009 |
| 119. | Caffeine Citrate 20 mg/ml (equivalent to 10 mg of Caffeine base) Oral Solution (Addl. Dosage form) | Same as approved | 18-7-2009 |
| 120. | Amphotericin B Emulsion 5 mg/ml (Addl. Indication) | Febrile Neutropenia in cancer patients | 18-7-2009 |
| | | | |
| 121. | Hydroxy ethyl starch 10% solution for infusion | For the treatment of imminent or manifest hypovolemia and shock | 18-7-2009 |
| 121. 122. | Hydroxy ethyl starch 10% solution for infusion Naratriptan (as HCI) Tablets 1/2.5 mg | | 18-7-2009 18-7-2009 |
| | | shock For the acute treatment of migraine with or without aura in | |
| 122. | Naratriptan (as HCl) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/ | shock For the acute treatment of migraine with or without aura in adults | 18-7-2009 |
| 122. 123. | Naratriptan (as HCl) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in | 18-7-2009 20-7-2009 |
| 122. 123. 124. | Naratriptan (as HCl) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung | 18-7-2009 20-7-2009 21-7-2009 |
| 122. 123. 124. | Naratriptan (as HCI) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation 500000IU/1000000IU/2000000IU | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung infection in patients with cystic fibrosis For the treatment of hypertension/ cardiac failure and | 18-7-2009 20-7-2009 21-7-2009 28-7-2009 |
| 122. 123. 124. 125. | Naratriptan (as HCI) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation 500000IU/1000000IU/2000000IU Ramipril MR capsule 10 mg | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung infection in patients with cystic fibrosis For the treatment of hypertension/ cardiac failure and myocardial infarction For the treatment of HIV-1 infection in combination with | 18-7-2009 20-7-2009 21-7-2009 28-7-2009 |
| 122. 123. 124. 125. 126. | Naratriptan (as HCI) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation 500000IU/1000000IU/2000000IU Ramipril MR capsule 10 mg Abacavir Sulfate Tablet 60 mg | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung infection in patients with cystic fibrosis For the treatment of hypertension/ cardiac failure and myocardial infarction For the treatment of HIV-1 infection in combination with other antiretroviral agents Prevention of clinical fractures after hip fracture in men and | 18-7-2009 20-7-2009 21-7-2009 28-7-2009 30-7-2009 |
| 122. 123. 124. 125. 126. 127. | Naratriptan (as HCl) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation 500000IU/1000000IU/2000000IU Ramipril MR capsule 10 mg Abacavir Sulfate Tablet 60 mg Zoledronic Acid Solution for Infusion (Addl. Indication) | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung infection in patients with cystic fibrosis For the treatment of hypertension/ cardiac failure and myocardial infarction For the treatment of HIV-1 infection in combination with other antiretroviral agents Prevention of clinical fractures after hip fracture in men and women | 18-7-2009 20-7-2009 21-7-2009 28-7-2009 30-7-2009 1-8-2009 |
| 122. 123. 124. 125. 126. 127. 128. | Naratriptan (as HCI) Tablets 1/2.5 mg Colistimethate Sodium powder for solution for injection/infusion 500000IU and 2000000 IU (Additional Strength) Simvastatin 20 mg + Ramipril 5 mg + Atenolol 50 mg + HCTZ 12.5 mg + Aspirin 100 mg (EC Pellets) Capsules Colistimethate Sodium powder for solution for inhalation 500000IU/1000000IU/2000000IU Ramipril MR capsule 10 mg Abacavir Sulfate Tablet 60 mg Zoledronic Acid Solution for Infusion (Addl. Indication) Sumatriptan 85 mg + Naproxen 500 mg Tablets | shock For the acute treatment of migraine with or without aura in adults Same as approved For secondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate Treatment by inhalation of Pseudomonas aeruginosa lung infection in patients with cystic fibrosis For the treatment of hypertension/ cardiac failure and myocardial infarction For the treatment of HIV-1 infection in combination with other antiretroviral agents Prevention of clinical fractures after hip fracture in men and women Additional strength for the treatment of migraine For the treatment of acute inflammation conditions | 18-7-2009 20-7-2009 21-7-2009 28-7-2009 30-7-2009 1-8-2009 3-8-2009 |

| 132. | Ofloxacin 2 mg + Ornidazole 5 mg per ml Infusion | For the treatment of diarrhoea of mixed infection in adult | 17-8-2009 |
|------|---|---|-----------|
| | | patients | |
| 133. | Clozapine Tablet 200 mg (Addl. Strength) | Indicated in the management of Schizophrenic patients | 19-8-2009 |
| 134. | Oxymetazoline Hydrochloride Nasal Spray 50 mcg/0.05%w/w | Is indicated for the relief of stuffy, running nose associated with colds and influenza, also post nasal drip, sinusitis | 19-8-2009 |
| 135. | Enoxaparin Sodium Injection (Addl. Indication) | Treatment of acute ST-segment elevation myocardial infarction inclu-ding patients to be managed medically or with subsequent percuaneous coronary intervention (PCI) | 19-8-2009 |
| 136. | Fluvoxamine Maleate 100/150 mg ER Tablets | For the treatment of obsessive compulsive disorder (OCD) and depression | 19-8-2009 |
| 137. | Lidocaine Lozenges 200/300 mg (Addl. Indication) | For single agent anesthesia in procedure involving the introduction of instrument & tubes into respiratory tract | 19-8-2009 |
| 138. | Risperidone long acting injection (Addl. Indication) | Indicated as monotherapy or as adjunctive therapy in lithium or valproate for the maintenance treatment of Bipolar-1 disorder | 29-8-2009 |
| 139. | Exemestane 25 mg Tablet (Addl. Indication) | Indicated for adjuvant treatment of Post Menopausal Women with estrogen receptor positive early breast cancer who have received two-three year of tamoxifen and are switched to exemestane for completion of total of five years of adjuvant hormonal therapy | 29-8-2009 |
| 140. | Artesunate (25/50/100 mg) + Amodiaquine (67.5 mg/135/270 mg) bilayerd Tablets | For the treatment of uncomplicated malaria due to plasmodium falciparum strain in various age groups | 1-9-2009 |
| 141. | Ezetimibe 10 mg + Fenofibrate 145 mg film coated tablets | For the treatment of combined hyperlipidemia in patients with normal hepatic and renal function | 1-9-2009 |
| 142. | Buprenorphine Transdermal Patch each Transdermal Patch having an area of 6.25cm², 12.5cm², and 25cm² contains: Buprenorphine5 mg/10 mg/20 mg having release rate of 5 mcg, 10 mcg, 20 mcg Buprenorphine per hour respectively (over a period of 7 days). | For the treatment of severe opioid responsive pain conditions which are not adequately responding to non-opioid analgesics. For Hospital supply only | 3-9-2009 |
| 143. | Trospium Chloride ER Capsules 60 mg | For the treatment of overactive bladder with the symptoms of urge urinary incontinence, urgency and urinary frequency | 4-9-2009 |
| 144. | Metoclopramide (as HCl) 200 mg/400 mg per ml Intranasal Solution | For the treatment of nausea and vomiting | 9-9-2009 |
| 145. | Isosorbibe 20 mg + Hydralazine 37.5 mg tablets | For the treatment of heart failure as an adjunct to standard therapy in patients to improve survival, to prolong time to hospitalization for heart failure and to improve patients reported functional status | 18-9-2009 |
| 146. | Cefixime 200 mg + Dicloxacillin ER 500 mg Tablets | For the treatment of adult patients with Upper and Lower respiratory tract infections, skin and soft tissue infections | 22-9-2009 |
| 147. | Chlorthalidone Tablet 6.25 mg | For the treatment of mild to moderate hypetension | 3-10-2009 |
| 148. | Cefovecin (as sodium) 80 mg/ ml Lyophilised powder for injection (Veterinary) | For the treatment of skin bacterial infections in dogs and cats | 3-10-2009 |
| 149. | Imidacloprid 10% + Moxidectin 1.0% & 2.5% Topical solution (Veterinary) each pipette contains: | For the treatment and prevention of flea infestations, prevention of heartworm disease and treatment of infections with gastrointestinal nematodes in dogs and cats | 3-10-2009 |
| | (a) 0.4 ml, 1.0 ml, 2.5 ml, 4 ml of 10% Imidacloprid and 1.0% moxidectin | | |
| | | ı | |

| 150. | Isoflupredone Acetate Suspension for Injection (2 mg per ml) (For Veterinary) | For the glucocorticoid, anti-inflammatory or supportive therapy to cattle suffering from conditions such as: | 3-10-2009 |
|------|--|---|------------|
| | (1 of veterinary) | (1) Overwhelming infections with severe toxicity | |
| | | (2) Bovine ketosis | |
| | | (3) Musculoskeletal conditions | |
| | | (4) Shock | |
| | | (5) Parturient paresis (supportive stress therapy for milk | |
| | | fever) | |
| 151. | Saxagliptin (as HCl) 2.5 mg/5 mg Tablets | As an adjunct to diet and exercise to improve glycemic control in adults with Type 2 Diabetes mellitus. (It should not be used for the treatment of Type 1 Diabetes mellitus or Diabetic ketoacidosis, as it would not be effective in these settings | 24-10-2009 |
| 152. | Calcipotriol Gel 0.005% (Additional dosage form) | For the treatment of moderate plaque psoriasis in adults | 3-10-2009 |
| 153. | Lamotrigine MR Tablets 150 mg (Addl. strength) | Same as approved | 3-10-2009 |
| 154. | Metoprolol Tartrate ER Tablet 75 mg (Additional strength) | Same as approved | 3-10-2009 |
| 155. | Valganciclovir Powder for Oral Solution 50 mg/ml | For the treatment of cytomegalovirus (CMV) retinitis in AIDS patients and for prevention of CMV disease in solid organ transplant patients at high risk | 9-10-2009 |
| 156. | Dexketoprofen Trometamol Gel 12.5 mg/gm | For the painful and inflammatory conditions associated with joints, tendons, ligaments and muscle | 13-10-2009 |
| 157. | Donepezil Hydrochloride Orodispersible Tablet 5 mg/10 mg | For the treatment of mild to moderate severe Alzheimer's Dementia | 15-10-2009 |
| 158. | Dexketoprofen Trometamol Injection 50 mg/2 ml | Symptomatic treatment of acute pain of moderate to severe intensity, when oral administration is not appropriate such as post-operative pain, renal colic and low back pain | 22-10-2009 |
| 159. | Bromocriptine Mesylate Tablet IP 0.8 mg (Additional Indication) | Indicated as an adjunct to diet and exercise to improve glycemic control in adults with Type 2 Diabetes mellitus | 22-10-2009 |
| 160. | Levosulpride ER Tablet 100/150/200 mg | For the treatment of depression and schizophrenia in adults | 27-10-2009 |
| 161. | Ambroxol 30 mg + Terbutaline 2.5 mg + Guaiphenesin 100 mg + Menthol 2.5 mg per 5 ml syrup | Additional strength | 27-10-2009 |
| 162. | Propofol 10 mg + Lignocaine 2.5 mg per ml injection | Additional strength | 27-10-2009 |
| 163. | Mecobalamin 750 mg + Pyri-doxine Hydrochloride 1.5 mg + Nicotinamide 45 mg tablets | For the treatment of Diabetic neuropathy | 27-10-2009 |
| 164. | Olmesartan (20/40 mg) + Atorvastatin 5/10/20 mg film coated tablets | For the treatment of co-existing essential hypertension and hyperlipidemia in adult patients | 27-10-2009 |
| 165. | Deltamethrin (0.50% m/v) and Piperonyl Butoxide (2.0% m/v) | Ready to use pour-on for cattle, sheep and goats. | 29-10-2009 |
| | per ml Pour-On Solution | 1. For use in cattle to control ticks, stable flies, horn flies, cattle louse flies and nuisance flies e.g. house flies, to kill lice and protection against blackflies | |
| | | 2. For use in sheep and goats for controlling Karoo paralysis, bont-legged and red-legged ticks | |
| 166. | Glucosamine sulphate potassium chloride 500 mg + Ibuprofen IP 200 mg film coated Tablets | For the treatment of adult patients with osteoarthritis of knee and/or hip joints | 3-11-2009 |
| 167. | Methylcobalamin 1500 mcg + Gabapentin SR 600 mg film coated tablets | Additional strength | 4-11-2009 |

| 168. | Propofol 1% for injection (Additional Indication) | For conscious sedation for surgical and diagnostic procedures | 5-11-2009 |
|------|--|---|------------|
| | | 2. For sedation of ventilated patients receiving intensive care | |
| 169. | Rupatadine 10 mg + Montelukast 10 mg Tablets | For the management of seasonal allergic rhinitis (SAR) | 10-11-2009 |
| 170. | Eperisone 50 mg + Paracetamol 500 mg Tablets | For improvement of mytonic conditions caused by neck shoulder arm syndrome, scapulhumeral periarthritis and low back pain in adult patients only | 10-11-2009 |
| 171. | Lafutidine Tablets5 mg/10 mg | For the treatment of: | 11-11-2009 |
| | | Gastric ulcers, duodenal ulcers and stromal ulcers | |
| | | Gastric mucosal lesions (erosion, hemorrhage, redness or edema) associated with acute gastritis and acute exacerbation of chronic gastritis | |
| 172. | Ramosetron Hydrochloride Tablets 2.5 mcg, 5 mcg, 0.1 mg & Injection 0.3 mg/2 ml | 1. Tablets 0.1 mg & Injection 0.3 — for the prevention and treatment of gastrointestinal symptoms (nausea and vomiting) associated with emetogenic cancer chemotherapy. | 11-11-2009 |
| | | 2. Tablets 2.5 mcg & 5 mcg — for the treatment of diarrhoea predominant irritable bowel syndrome | |
| 173. | Tadalafil tablet 20 mg (Additional Indication) | For the treatment of pulmonary arterial hypertension | 14-11-2009 |
| 174. | Flurbiprofen Sodium 0.03% Ophthalmic solution USP | For the reduction of post-operative inflammatory condition | 14-11-2009 |
| 175. | Dexmedetomidine HCl Injection (Additional Indication) | Sedation of non-intubated patients prior to and/or during surgical and other procedure | 14-11-2009 |
| 176. | Paliperidone ER tablet 12 mg (Additional Strength) | Same as approved | 14-11-2009 |
| 177. | (1) Triticum Vulgare Topical Cream each 100gm of cream contains: | (1) Triticum Vulgare Cream is indicated for the treatment of ulcers and bed sores. | 14-11-2009 |
| | Aqueous extract of Triticum Vulgare (dry residue 200 mg/ 100 ml) 15 gm | (2) Triticum Vulgare Vaginal Pessary is indicated for the treatment of phlogistic and dystrophic disorders of the female genital tract | |
| | 2-phenoxyethanol Eur. Ph 1 gm | | |
| | (2) Triticum Vulgare Vaginal Pessary each 3.5 gm of vaginal pessary contains: | | |
| | Aqueous extract of Triticum Vulgare (dry residue 200 mg/ 100 ml) 0.60 gm | | |
| | 2-phenoxyethanol Eur. Ph 0.04 gm | | |
| 178. | Febuxostat Tablets 80 mg/120 mg | For treatment of chronic hyperuricemia in conditions where urate depression has already occurred (including a history, or presence of tophus and/or gouty arthritis) | 14-11-2009 |
| 179. | Euphorbia Prostrata Dry Extract : 100 mg tablet (Addl. Indication) | For the treatment of 3° and 4° haemorrhoids (Addl. Indication) | 14-11-2009 |
| 180. | Euphorbia Prostrata dry extract ethanolic 80%v/v [(35-70):1] cream 1% w/w | For the treatment of 1° and 2° internal haemorrhoids | 14-11-2009 |
| 181. | Lactobacillus Rhammnosus 1 billion cells + Saccharomyces Boulardi 125 mg + Zinc Enriched yeast 20 mg sachets & capsules | For the treatment of acute diarrhoea in children | 4-11-2009 |
| 182. | Nimesulide 1.0% w/w + Ethanol 31% w/w topical solution | For the local application in relieving pain in acute low backache, tendonitis, tenosinovitis and sprains | 17-11-2009 |
| 183. | Diacerein Modified Release Capsule 100 mg | Symptomatic treatment of osteoarthritis of knee & hip | 20-11-2009 |
| 184. | Amlodipine 5 mg/10 mg/5 mg/ 10 mg/10 mg + Valsartan 160 mg/160 mg/160 mg/160 mg/320 mg + Hydrochlorothizide 12.5 mg/12.5 mg/25 | For the treatment of essential hypertension | 24-11-2009 |

| 185. | Maraviroc Tablets 150 mg/300 mg | For the treatment of patients infected with CCR 5-trophic HIV-1 | 25-11-2009 |
|------|--|--|------------|
| 186. | S (+) Etodolac 150/200/300 mg + Paracetamol 500/500/500 mg Tablets | For the symptomatic treatment of acute pain and inflammation in patients with osteoarthritis, rheumatoid arthritis and ankylosing spondylitis | 25-11-2009 |
| 187. | Tacrolimus Granules for oral suspension 0.2/1 mg per sachet | Prophylaxis of transplant rejection in kidney, liver or heart allograft recipient | 26-11-2009 |
| 188. | Rifaximin Dispersible Tablet 200 mg | For infections diarrhoea in adults | 26-11-2009 |
| 189. | Prednisolone Sodium Phosphate Orally Disintegrating Tablet 10/15/30 mg | Indicating for condition required Glucocorticoids activity including collagen disease, allergic disease and hypersensitive reaction | 26-11-2009 |
| 190. | Cinitapride Hydrogen Tartrate ER Tablet 3 mg | For the treatment of gastroesophageal reflux disease (GERD) and functional inconveniences of the gastrointestinal motility | 26-11-2009 |
| 191. | Sevelamer Carbonate (Additional Indication) | For control of serum phosphorus in patients with chronic kidney disease on haemodialysis or peritoneal dialysis and patients with CKD not on dialysis with serum phosphate > 5.5 mg/dl | 26-11-2009 |
| 192. | Oxcarbazepine Tablets (Additional Indication) 150/300/600 mg | Monotherapy in the treatment of partial seizures in children ages 4-16 | 26-11-2009 |
| 193. | Ketoprofen plaster 30 mg (7cm X 10 cm) | For the relief of musculoskeletal pain & inflammation | 11-12-2009 |
| 194. | Acipimox Capsules 250 mg | As adjunctive therapy to diet and weight loss in the treatment of Type II-a, II-b and IV lipid disorders | 26-11-2009 |
| 195. | Amisulpiride SR Tablet 100 mg | Same as approved | 27-11-2009 |
| 196. | Mycophenolate Sodium Enteric coated tablet 540 mg (Additional Strength) | Same as approved | 29-10-2009 |
| 197. | Loteprednol etabonate 0.5% w/v + Moxifloxacin 0.5% w/v opthalmic Suspension | For the treatment of post operative steriod responsive inflammatory ocular infections or risk of bacterial ocular infections exists | 1-12-2009 |
| 198. | Ropinirole ER tablet 2/4/8 mg | For the treatment of signs & symptoms of idiopathic Parkinson's disease | 3-12-2009 |
| 199. | Diclofenac Diethyl amine transdermal patch 100 mg/37.5 sq cm (Additional Strength) | Same as approved | 4-12-2009 |
| 200. | Lornoxicam ER Tablet 8 mg (Additional Strength) | Same as approved | 11-12-2009 |
| 201. | Flurbiprofen plaster 20 mg (7 cm x 10 cm) | For the relief of pain & inflammation due to rheumatoid arthritis, osteoarthritis and allied disorder | 11-12-2009 |
| 202. | Carvedilol Phosphate ER tablet 10/20/40/80 mg | 1. Mild to severe chronic heart failure. | 11-12-2009 |
| | | Left ventricular dysfunction following myocardial infarction in clinically stable patients. hypertension | |
| 203. | Halometasone monohydrate 0.05% w/w + Fusidic acid 2.00% w/w cream | For the treatment of acute and chronic infected eczematous dermatitis | 15-12-2009 |
| 204. | Glimepiride 1/2 mg + Metformin 500 mg + Atorvastatin 10 mg tablets | For the treatment of type II diabetes mellitus associated with dyslipidemia | 17-12-2009 |
| 205. | Betahistine Di hydrochloride CR capsule 24/32/48 mg | Indicated in the treatment of Menier's syndrome, characterized by unilateral or bilateral tinnitus, vertigo, sensorineural hearing loss | 17-12-2009 |
| 206. | Gefitinib tab 250 mg (Additional Indication) | Indicated for the first line treatment of patients with locally advanced or metastatic non small cell lung cancer (NSLC) who have activating mutation of the EGFRTK | 18-12-2009 |

| 207. | Midazolam HCl Syrup 2 mg/ml | For selection, anxiolysis and amnesia prior to diagnostic, therapeutic or endoscopic procedure or before induction of anesthesia | 18-12-2009 |
|------|--|--|------------|
| 208. | Guaiphenasin Syrup 133 mg/5 ml | For the treatment of productive cough accompanied by phlegm | 18-12-2009 |
| 209. | Balofloxacin Tablet 100 mg | For the treatment of uncomplicated urinary tract infections (such as cystitis and urethritis) | 22-12-2009 |
| 210. | Xylometazoline 0.5 mg + Ipratropium 0.6 mg Nasal spray | For symptomatic treatment of nasal congestion with running nose (rhinorrhea) | 23-12-2009 |
| 211. | Doxofylline 400 mg + Ambroxol 30 mg Tablets | For the treatment of patients with bronchial asthma and COPD, when use of both the drug is appropriate | 23-12-2009 |
| 212. | Ribavirin Capsules 200 mg (Addl. Strength) | Indicated in combination with peg interferon alpha-2b for the treatment of naïve adult patients with chronic Hepatitis C with clinically stable HIV co-infection | 24-12-2009 |
| 213. | Formoterol 6 mcg + Mometa-sone furoate 100/200/400 mcg Metered dose Inhaler & Dry Powder Inhalation | For the treatment of patients with bronchial asthma and COPD, where the use of combination is found appropriate | 30-12-2009 |
| 214. | Chlorohexidine Gluconate Gel 0.2% | For the prevention of Plaque and treatment of gengivitis | 30-12-2009 |
| 215. | Chlorohexidine Gluconate Mouth wash 0.1% | For the prevention of Plaque and treatment of gengivitis | 30-12-2009 |

| Sl. No. | Product Name | Indication | Date of Approval |
|---------|---|--|------------------|
| 1. | Ferrous ascorbate 100 mg + Folic acid 1.1 mg tablet and Omega-3-fatty acid 200 mg capsules combikit | For iron, folic acid & omega-3-fatty acid deficiency | 8-1-2010 |
| 2. | Sphaeranthus Indicus extract Tablets 700 mg | For the management of psoriasis | 8-1-2010 |
| 3. | Febuxostat 40 mg (Additional Strength) | Same as approved | 16-1-2010 |
| 4. | Dexamethasone Intravitreal Implant 0.7 mg (In solid polymer drug delivery system) | For the treatment of macular edema following branch retinal vein occlusion (CRVO) | 16-1-2010 |
| 5. | Buclizine HCl 25 mg Tablet (Additional Indication) | For the Symptomatic treatment of various allergic conditions (rhinitis, conjunctivitis and urticaria) and for prevention and treatment of motion sickness | 16-1-2010 |
| 6. | Pemetrexed Disodium 500 mg/ 100 mg Powder for Injection (Additional indication) | Indicated as a monotherapy for the maintenance treatment of locally advanced or metastatic Non Small Cell Lung Cancer (NSCLC) other than predominantly squamous cell histology in patients whose disease has not progressed immediately following platinum-based chemotherapy. First line treatment should be a platinum doublet with gemcitabine, paclitaxel or docetaxel | 16-1-2010 |
| 7. | Levobetaxolol HCl Opthalmic Suspension 0.5% | For lowering of intraoccular pressure in patients with chronic open-angle glaucoma or occular hypertension | 16-1-2010 |
| 8. | Sarpogrelate Hydrochloride film coated Tablets 100 mg | For the improvement of ischemic symptoms including ulcer, associated with chronic arterial occlusion | 16-1-2010 |
| 9. | Propanol 1-ol 18 gm + Ethanol 100% 45 gm/100 ml solution | For hygenic and surgical hand disinfection | 18-1-2010 |
| 10. | Aliskiren 150/160 mg + Valsartan 300/320 mg tablets | For the treatment of hypertension | 19-1-2010 |
| 11. | Piroxicam 20 mg tab & Paracetamol 500 mg tab combikit | For treatment of fever and pain associated with acute upper respiratory tract inflammation, acute musculoskeletal disorders, pain after operative intervention and following trauma, chronic condition like rheumatoid arthritis, osteoarthritis etc for short term management of acute painful episodes | 19-1-2010 |

| 12. | Soybean oil 60 gm + Triglycerides (medium chain) 50 gm + olive oil purified fish oil 30 gm + dl-alpha-tocopherol 163-225 mg + glycerol (anhydrous) 25 gm + Egg lecithin 12 gm + sodium hydroxide (to approx pH 8) + sodium pleate 0.3 mg emulsion for infusion in 100 ml | thoracic or major urological surgery | 19-1-2010 |
|-----|--|--|-----------|
| 13. | Raltegravir (as Potassium) film coated Tablets 400 mg | In combination with other anti-retroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection in treatment experienced patients with evidence of HIV-1 replication despite ongoing retroviral therapy | 27-1-2010 |
| 14. | Deflazacort Tablet 18 mg (Additional Strength) | Same as approved | 30-1-2010 |
| 15. | Pregabalin SR Tablet 75 mg (Additional Strength) | Same as approved | 30-1-2010 |
| 16. | Rivaroxaban film coated Tablets 10 mg | For the prevention of venous thromboembolism in patients (VTE) in adult patients undergoing hip or knee replacement surgery. | 30-1-2010 |
| 17. | Cinnarizine 20 mg + Dimenhydrinate 40 mg tablets | For the treatment of vertigo | 3-2-2010 |
| 18. | Lipid based Amphotericin-B Gel 0.1% | Indicated in the treatment of cutaneous and mucocutaneous mycotic infections caused by candida (Monilia) species | 5-2-2010 |
| 19. | Sodium Hyaluronate Injectable Gel 14 mg/ml, 18 mg/ml, 25.5 mg/ml (Additional Strength) | Same as approved | 6-2-2010 |
| 20. | Milnacipran HCl 12.5mg (Additional Strength) | Same as approved | 6-2-2010 |
| 21. | Ropinirole 6mg ER Tablets (Additional Strength) | Same as approved | 6-2-2010 |
| 22. | Coseal Surgical Sealent 2/4/8 ml (Additional Indication) | (1) Enforcement of suture and staple lines in lung resection procedures. | 6-2-2010 |
| | | (2) Patients undergoing cardiac surgery to prevent or reduce the incidence, severity and extend of post-surgical adhesion formation. | |
| | | (3) Patients undergoing laparotomy or laproscopic abdomino- pelvic surgery as an adjunct to good surgical technique intended to reduce the incidence, severity and extent of post- surgical adhesion formation | |
| 23. | Pregabalin SR 75/150/300 mg + Methylcobalamin 1500 mcg tablets | For the treatment of adult patients with peripheral neuropathy | 9-2-2010 |
| 24. | Ivabradine HCl Tablets 5/7.5 mg (Additional Indication) | For Symptomatic Treatment of chronic stable angina pectoris in coronary artery disease patients with normal sinus rhythm, indicated in combination with beta-blockers in patients inadequately controlled with an optimal beta-blocker dose and whose heart rate is > 60 bpm | 10-2-2010 |
| 25. | Paracetamol Solution for Infusion (10mg/ml) in 50ml (Additional Packsize) | Same as approved | 10-2-2010 |
| 26. | Ketorolac Tromethamine Ophthalmic Solution 0.45% | Indicated for the treatment of pain and inflammation following cataract surgery | 10-2-2010 |
| 27. | Cytarabine Injection 100 mg Additional Packsize (20 ml) | In combination with other approved anti-cancer drugs is indicated for remission induction in acute non-lymphocytic leukemia of adults and children. It has also been found useful in the treatment of acute lymphocytic leukemia and the blast phase of chronic myelocytic leukemia. Intrathecal administration of cytarabine is indicated in the prophylaxis and treatment of meningal leukemia | 10-2-2010 |
| 28. | Olopatadine HCl Nasal Spray 0.6% w/w (Additional Indication) | For the relief of the symptoms of seasonal allergic rhinitis in patients 6 years of age and older | 10-2-2010 |

| 29. | Sodium chloride 6.80 gm + Potassium chloride 0.30 gm + Calcium chloride dihydrate 0.37gm + Magnesium chloride hexahydrate 0.20 gm + Sodium acetate trihydrate 3.27 gm + L-malic acid 0.67 gm in 1000 ml solution for infusion | dehydration, especially if acidosis is being imminent or | 11-2-2010 |
|-----|---|---|-----------|
| 30. | Chemically Modified Sodium Hyaluronate and Caboxymethylcellulose Absorbable Adhesion Barrier | As an adjunct in abdominal or pelvic surgery for reducing the incidence, extent and severity of post-operative adhesions at the site of placement | 11-2-2010 |
| 31. | Lornoxicam 4 mg/4 mg + Thiocolchicoside 4 mg/8 mg tablets | For the treatment of patients with acute painful musculoskeletal conditions | 12-2-2010 |
| 32. | Cefixime 100 mg/200 mg + Cloxacillin 500 mg/500 mg tablets | For the treatment of adult patients with upper & lower respiratory tract infections, skin and soft tissue infections | 12-2-2010 |
| 33. | Celecoxib Mouth Dissolving Tablet 50/100/200 mg | For the treatment of Osteoarthritis and Rheumatoid Arthritis | 25-2-2010 |
| 34. | Propofol 2% w/v Injection (Additional Strength) | Same as Approved | 27-2-2010 |
| 35. | Spironolactone 25mg + Furosemide 20mg tablets | For treatment of resistant oedema associated with secondary hyperaldosteronism; resistant hypertension, chronic cardic failure and hepatic cirrhosis | 5-3-2010 |
| 36. | Olopatadine 5mg + Ambroxol 30mg tablets | For the management of cough in adult patients only | 5-3-2010 |
| 37. | Methylcobalamin 750mcg + Pregabalin 75/150mg + Vitamin B 61.5mg + Folic Acid 0.75mg + Benfothiamine 7.5mg capsules | For the treaetment of painful diabetic neuropathy in adults only | 5-3-2010 |
| 38. | Calcium Acetate 435mg + Magnesium Carbonate 235mg tablets | For the management of hyperphosphatemia in adults patients with renal failure being treated with dialysis (hemodialysis or peritoneal dialysis | 5-3-2010 |
| 39. | Dicyclomine 2.5mg + Dried Aluminium Hydroxide 200mg + Light Magnesium Oxide 100mg + Simethicone 20mg per 5ml suspension | For the treatment of the symptoms of functional gastrointestinal disorders like irritable bowel syndrome, functional dyspepsia, peptic ulcers, GERD which includes smooth muscle spasm, flatulence, abdominal distension, hyperacdity, gastric distress, bloting etc. | 5-3-2010 |
| 40. | Sertaconazole 2% w/v+ Zinc Pyrithione 1% w/v Shampoo | For the treatment in adults patients with dandruff/seborrheic dermatitis of the scalp | 5-3-2010 |
| 41. | Faropenam Sodium Extended Release Tablet 225/300/450 mg | For the treatment of respiratory tract infections, urinary tract infection, skin and skin structure infections and gynaecological infections | 8-3-2010 |
| 42. | Amlodipine 5mg + Ramipril 10mg/5mg Tablets (Additional Strength) | For the treatment of hypertension only | 10-3-2010 |
| 43. | Doxofylline MR Tablet 650 mg | For the treatment of Bronchial Asthma and COPD in Adults | 11-3-2010 |
| 44. | Lidocaine 7.5mg + Prilocaine 2.5mg per acutation spray | For the treatment of adult male patients with premature ejaculation | 12-3-2010 |
| 45. | Pramipexole Di HCl Monohydrate ER Tablet 0.375/ 0.75/1.5/3/4.5 mg | For the treatment of the sign and symptoms of Idiopathic Parkinsons disease | 13-3-2010 |
| 46. | Drosperinone 3mg + Ethinyl Estradiol 0.02mg tablets | For the treatment of symptoms of premenstrual dysphoric disorders (PMDD) in women (Additional Indication) | 16-3-2010 |
| 47. | Miglitol 25 mg + Metformin 500 mg tablets | Additional Strength | 16-3-2010 |
| 48. | Propranolol 40mg (SR Pellets) + Fluarizine 5mg/10mg capsules | For the prophylaxis of migraine | 16-3-2010 |
| 49. | Citicholine 500mg + Piracetam 400mg tablets | For the treatment of acute stroke | 16-3-2010 |
| 50. | Diclofenac 50mg + Chlorzoxazone 500mg tablets | For the treatment of acute inflammation conditions associated with spasm in adults only | 16-3-2010 |

| 51. | Mecobalamin 750 mcg + Pyridoxine 1.5 mg + Nicotinamide 45 mg + Benfotiamine 150 mg tablets | For the treatment of diabetic neuropathy in adults | 16-3-2010 |
|-----|--|---|-----------|
| 52. | Etoricoxib 1%+ Menthol 5% Spray | For the topical treatment of acute musculoskeletal pain | 16-3-2010 |
| 53. | Pantoprazole 40mg + Cinitapride (ER) 3mg capsules | For the treatment of patients suffering from non-ulcer dyspepsia (NUD) or gasteroesophageal reflux disease (GERD) | 17-3-2010 |
| 54. | Atorvastatin 20mg + Ramipril 2.5mg/5mg capsules | Additional higher strength | 17-3-2010 |
| 55. | Lamotrigine Orodispersible Tablet 12.5 mg (Additional Strength) | Same as Approved | 20-3-2010 |
| 56. | Trabectedin powder for conc. solution for infusion (Additional Indication) | In combination with pegylated liposomal doxorubicin, it is indicated for the treatment of patients with relapsed platinum-sensitive ovarian cancer | 20-3-2010 |
| 57. | Lornoxicam 8mg/8mg + Thiocolchicoside 4mg/8mg tablets | Additional higher strength | 23-3-2010 |
| 58. | Diclofenac Sodium SR Tablet 150 mg | For the treatment of rheumatoid arthritis, osteoarthritis, ankylosing spondilytis, gout, painful post-perative pain following dental surgery etc. | 26-3-2010 |
| 59. | Magnesium Valproate CR Tablet 200/ 300/400/ 500 mg (Additonal Strength) | Same as approved | 26-3-2010 |
| 60. | Lidocaine 2.5% w/w+Prilocaine 2.5% w/w Gel | Additional higher strength | 26-3-2010 |
| 61. | Nicotinic Acid and Laropiprant Modified Release Tablets: Nicotinic Acid1000 mg Laropiprant20mg | 1. For the treatment of dyslipidaemia, particularly in patients with combined mixed dyslipidaemia (characterised by elevated levels of LDL-cholesterol and triglycerides and low HDl-cholesterol) and in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial). 2. In combination with HMG-CoA reductase inhibitors (statins), when cholesterol lowering effect of HMG-CoA reductase inhibitor monotherapy is inadequate. It can be used as monotherapy only in patients in whom HMG-CoA reductase inhibitors are considered inappropriate or not tolerated. Diet or other non-pharmacological treatments (e.g. exercise, weight reduction) should be continued during therapy. | 26-3-2010 |
| 62. | Pantoprazole Mouth Dissolving Tablet 20/40 mg | For the treatment of gastric ulcer, duodenal ulcer & gastroesophageal reflux disease (GERD) | 30-3-2010 |
| 63. | Praziquntel 175mg + Pyrantel 504mg + Febental 525mg Tablets | For the treatment of mixed infections with roundworms and tapeworms in dogs | 30-3-2010 |
| 64. | Difluprednate Opthalmic Emulsion 0.05% w/v | For the treatment of inflammation and pain associated with ocular surgery | 30-3-2010 |
| 65. | Tolperisone HCl SR 450mg Tablets | For the relief of painful muscle spasms of the skeletal musculature | 30-3-2010 |
| 66. | S(-)Metoprolol 25mg/50mg (ER) + Atorvastatin (IR)10mg tablets | For the treatment of patients with both essential hypertension and hypercholesterolemia | 4-1-2010 |
| 67. | Atorvastatin10mg + Ramipril 5mg + Aspirin (EC pellets) 75mg/150mg + Metoprolol (ER tablets) 25mg Capsules | For secondary prophylaxis of ischemic heart disease in patients where use of such combinations in appropriate | 5-4-2010 |
| 68. | Lamotrigine 25mg Modified Release Tablet (Additional Strength) | Same as approved | 5-4-2010 |
| | | | |

| 70. | Azithromycin SR Granules for Oral Suspension 1gm (Additional Strength) | Same as Approved | 5-4-2010 |
|-----|--|--|-----------|
| 71. | Gemfibrozil Capsule USP & Gemfibrozil Tablet USP 200mg/600mg (Additional Indication) | For the treatment of other dyslipidemias: (a) Fredrickson Types III and V (b) Associated with diabetes | 5-4-2010 |
| 72. | Ibuprofen 400mg + Chlorzoxazone 500mg tablets | (c) Associated with xanthomata For the short term treatment of musculoskeletal pain associated with inflammation and spasm in adults only | 7-4-2010 |
| 73. | Doxophylline (SR) 400mg/ 800mg + Montelukast 10mg/ 10mg Tablets | For the treatment of bronchial asthma in adults | 7-4-2010 |
| 74. | Dronedarone(as Hydrochloride) film coated tablet400 mg. | To reduce the risk of cardiovascular hospitalization in patients with paroxysmal or persistent atrial fibrillation (AF) or atrial flutter (AFL), with a recent episode of AF/AFL and associated cardio-vascular risk factors (i.e., age > 70, hypertension, diabetes, prior cerebro-vascular accident, left atrial diameter > 50mm or left ventricular ejection fraction [LVEF] < 40%), who are in sinus rhythm or who will be cardioverted. | 8-4-2010 |
| 75. | Dosulepin HCl Tablet 25mg/75mg (Additional Indication) | For the treatment of chronic pain. | 8-4-2010 |
| 76. | Prasugrel (as Hydrochloride) film coated tablet 5 mg/ 10mg | To reduce the rate of thrombotic cardiovascular (CV) events (including stent thrombosis) in patients with acute coronary syndrome (ACS) who are to be managed with percutaneous coronary intervention (PCI) as follows: | 13-4-2010 |
| | | Patients with unstable Angina (UA) or non-ST-Elevation Myocardial Infarction (NSTEMI). | |
| | | Patients with ST-Elevation Myocardial Infarction (STEMI) when managed with primary or delayed PCI. | |
| 77. | Tofisopam uncoated tablet JP 50/100mg | For the treatment of anxiety and depression. | 13-4-2010 |
| 78. | Zotepine tablet 25/50/100mg | For the treatment of schizophrenia | 15-4-2010 |
| 79. | Armodafinil uncoated tablet 50/150/250mg | To improve wakefulness in patients with excessive sleepiness associated with treated obstructive sleep apnea (OSA), shift work sleep disorder (SWD) or narcolepsy. | 23-4-2010 |
| 80. | Indacaterol (as Maleate) Inhalation Powder Hard Capsules | For maintenance bronchodilator treatment of airflow obstruction in adult patients with chronic obstructive pulmonary disease (COPD). | 24-4-2010 |
| 81. | Ranolazine ER Tablets 750mg (Additional Strength) | Same as Approved | 24-4-2010 |
| 82. | Tacrolimus Capsules 3mg (Additional Strength) | Same as approved | 24-4-2010 |
| 83. | Cefixime 100mg/200mg + Ofloxacin 100mg/200mg tablets | For the treatment of patients with typhoid fever and urinary tract infection in adults | 26-4-2010 |
| 84. | Liquid paraffin 6% + White soft paraffin 15% w/v Cream | For symptomatic relief of dermatological condition associated with dry skin including atopic dermatitis, eczema, pruritus, ichthyosis, xerosis, psoriasis | 29-4-2010 |
| 85. | Levosimendan Injection 2.5mg/ ml | For the short term treatment of acutely decompensated chronic heart failure (ADHF) in situations where conventional therapy is not sufficient and in cases where ionotropic support is considered appropriate. | 30-4-2010 |
| 86. | Thiocholchicoside SR Tablet 16 mg | As an adjuvant treatment in painful spasms associated with degenerative vertebral disorders and vertebral static problems, torticollis dorsal pain, low back pain, traumatological and neurological disorders. | 31-4-2010 |

| | T | | |
|------|---|---|-----------|
| 87. | Telmisartan 40/80 mg+ Indapamide SR 1.5/1.5 mg tablet | For the treatment essential hypertension | 5-4-2010 |
| 88. | One strip of 3 uncoated tablets of artesunate 150mg & one strip of 2 uncoated tablets of FDC of Sulphadoxine IP 500mg + Pyrimethamine IP 25mg | Additional strength | 5-4-2010 |
| 89. | Lapatinib Ditosylate Tablets 250mg (Additional Indication) | In combination with letrozole for the treatment of postmenopausal women with hormone receptor positive metastatic breast cancer that overexpresses the HER2 receptor for whom hormonal therapy is indicated. | 10-5-2010 |
| 90. | Pentosan Polysulfate Sodium Capsules 100mg | For the relief of bladder pain or discomfort associated with interstitial cystitis. | 11-5-2010 |
| 91. | Indapamide (SR) 1.5mg/1.5mg +Nebivolol 2.5mg/5mg tablet | For the treatment of hypertension not controlled with monotherapy | 12-5-2010 |
| 92. | Cinitapride (ER) 3mg + Omeprazole (EC) 20mg Capsules | For the treatment of gastirc ulcer,gastroesophageal reflux disease(GERD) & Dyspepsia not responding to omeprazole alone | 12-5-2010 |
| 93. | Timolol Maleate Long Acting Ophthalamic Solution 5mg/ml | For the treatment of elevated intraocular pressure in patient with ocular hypertension or open angle glaucoma. | 14-5-2010 |
| 94. | Telmisartan tablet 20/40/80 mg(Additional Indication) | For the prevention of cardiovascular morbidity and mortality in patient 55 years older at high risk of cardiovascular disease | 14-5-2010 |
| 95. | Dexketoprofen Trometamol 25mg + Thiocolchicoside 4mg tablets | For the treatment of acute low back pain and other musculoskeletal painful & inflammatory conditions in adult patients | 20-5-2010 |
| 96. | Troxipide film coated Tablets 50mg/100mg | For the treatment of Gastric ulcers and for amelioration of gastric mucosal lesions (erosion, hemorrhage, redness and edema) in the diseases such as acute gastritis, acute exacerbation stage of chronic gastritis | 20-5-2010 |
| 97. | Ambrisentan film coated Tablets 5 mg/10mg | For the treatment of pulmonary arterial hypertension (WHO Group I) in patients with WHO Class II or III symptoms to improve exercise capacity and delay clinical worsening | 24-5-2010 |
| 98. | Rosuvastatin Calcium Tablet (Additional Indication) | Risk reduction of MI stroke and arterial evascularisation procedure in patients without clinically evident CHD but with multiple risk factor | 24-5-2010 |
| 99. | Paracetamol SR 1000mg Bilayered tablet | For symptomatic treatment of pain and fever | 24-5-2010 |
| 100. | Tacrolimus PR Caps 0.5/1/5 mg | For Prophylaxis of transplant rejection in adult kidney or liver allograft rejection | 24-5-2010 |
| 101. | Rizatriptan Benzoate orally disintegrating tablet 10 mg(Additional Dosage Form) | Same as Already approved | 24-5-2010 |
| 102. | Ramipril 1.25/2.5/5/10 mg (Additional Indication) | For the treatment of non-diabetic overt glomerular or incipient nephropathy | 24-5-2010 |
| 103. | Exenatide Inj. (Additional Indication) | Use in patients with type-2 diabetes mellitus who are using a thiazolidinedione alone or in combination with metformilin but have not achieved adequate glycemic control | 24-5-2010 |
| 104. | Temozolomide Lyophilised powder 100mg/vial | For the treatment of | 24-5-2010 |
| | | "Adult patients with newly-diagnosed glioblastoma multiforme concomitantly with radiotherapy (RT) and subsequently as monotherapy treatment." | |
| | | Children from the age of three years, adolescents and Adult patients with malignant giloma, such as glioblastoma multiforme or anaplastic astrocytoma, showing recurrence or progression after standard therapy. | |

| 105 Pulicissore Projektines Storage 125mag/25mag + Salmaterol For the maintenance treatment of radiumand for the treatment 25.5-2010 | | | | |
|--|------|---|---|-----------|
| dermis injection as well as for tip definition. The presence of lidocaine is meant to reduce the patients pain during the treatment of lidocaine is meant to reduce the patients pain during the treatment of lidocaine is meant to reduce the patients pain during the treatment of lidocaine is meant to reduce painful inflammatory conditions in adults only 108. Metformin SR 500mg/500mg + GliclazideSR 60mg/30mg + As 3rd line treatment of Type II diabetes mellitus when diet, exercise and the single agents and the second line therapy with two drugs do not restalf and elequate glycemic control 3 alabets of Artesunate 100mg + 3 tablets of Artesunate 50mg a 2 tablets of FDC of Pyrimethamine 25mg + Sulphadoxime 50mg combipack 110. Culcium Carbanate 600 mg + Simethicume 80mg Chewable 111. Activated Dimethicone 50mg + Magnesium Hydroxide 250mg + Dried Aluminium Hydroxide250 mg + Sorbiol (70%)12.5 mg/5 mf 50 totion of secondary disconstitution of coronory heart of flattisence 112. Aspirin 75mg + Simvastatin 10mg/20mg + Lisinopril 5mg/ 10mg + Aceolod 25mg/50mg tablets 113. Ilsogladine Maleate Tablet 2mg/4mg For gastric ulexus, improvement of gastric mucosal lesion (crosion, hemorrhage, redness and clean) caused by acute sestifists and cance exacerbation stage of chronic gastritis. 114. Lacosamide film coated tablet 50mg/100mg/150mg/200mg 115. Carmustine for inj. 100 mg Indicated as pallifactive therapy as a single agent or in established combination therapy with other approved demotherapeatic agents in the following: 116. Licosamide film coated tablet 50mg/100mg/150mg/200mg 117. Ilsogladine Maleate Tablet 2mg/4 mg Indicated as pallifactive therapy as a single agent or in established combination therapy with other approved demotherapeatic agents in the following: 118 Insin tumors—glioblistoma, brainstern glioma, medalloblastoma, asstruytoma, epachymoma, and metastatic brain tumors. 2 Militipe myeloma—in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who | 105. | | | 25-5-2010 |
| adults only As a flue treatment of Type II diabetes mellitus when diet, exercise and the single agents and the second line therapy with two drugs do not result in adequate glycemic control 3 tablets of Artesunate 100mg + 3 tablets of Artesunante 50mg & 2 tablets of PIOC of Pyrimethamine 25mg + Sulphadoxine 500mg combipace. 110. Calcium Curbonate 600 mg + Simethicone 80mg Chewable tablet 111. Activated Dimethicone 50mg + Magnesium Hydroxide tablet 220mg- Dried Alominium Hydroxide:230 mg + Sorbiological for the tablet 230mg- Dried Alominium Hydroxide:230 mg + Sorbiological for the tablet 112. Aspirin 75mg + Simvastatin 10mg/20mg + Lisinopril 5mg/ 10mg + Atenold Zsmg/Smg/Smg tablets 113. Irsogladine Maleate Tablet 2mg/4mg 114. I accsamide film caated tablet. Somg /100mg/150mg/200mg 115. Camustine for inj. 100 mg 116. Camustine for inj. 100 mg 117. Brain tumors glioblastoma, satrocytoma, ependymoma, and metastatic brain tumors. 28.5-2010 28.5-2 | 106. | | dermis injection as well as for lip definition. The presence of lidocaine is meant to reduce the patients pain during the | 28-5-2010 |
| Pioglitazone 15mg/15mg tablets exercise and the single agents and the second line therapy with twodugs do not result in adequate glycemic control Additional Strength 2 subsets of FTDC of Pyrimethamine 25mg + Sulphadoxine 50mg combipace 110. Calcium Carbonate 600 mg + Simethicone 80mg Chewable tablet 250mg + Did on the subset of FTDC of Pyrimethamine 25mg + Sulphadoxine 250mg + Dried Aluminum Hydroxide 250mg + Sorbiol (70%)1.25 gm/5ml Solution 28.5-2010 epigastric discomfort, or their equivalents) that are often associated with dyspepsia, peptic ulcers, gastritis, peptic esphagists and indicated for relief of flatulence 25mg/5mg tablets 112. Aspirin 75mg + Simvastatin 10mg/20mg + Lisinopril 5mg/ 10mg + Atended 25mg/5mg tablets 113. Insegladine Maleate Tablet 2mg/4mg For secondary prevention of coronory heart discuse/stroke in patients where use of such combination is appropriate 114. Lacosamide film coated tablet 50mg/100mg/150mg/200mg 115. Carmustine for inj. 100 mg Indicated as palliative therapy as a single agent or in established combination therapy with other approved chemocheraptetic agents in the following: 1. Brain tumors—glioblastoma, astroxytoma, ependymoma, and metastatic brain tumors. 2. Multiple myelona—in combination with prednisone. 3. Hodglor's Disease—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodglin's lymphomas—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodglin's lymphomas—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodglin's lymphomas—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond | 107. | Aceclofenac 1% w/w + Menthol 5% Spray | - | 28-5-2010 |
| 8. 2 tablets of FDC of Pyrimethamine 25mg + Sulphadoxine S00mg combipack. 110. Calcium Carbonate 600 mg + Simethicone 80mg Chewable tablet 111. Activated Dimethicone 50mg + Magnesium Hydroxide Por treatment of symptoms of hyperacidity (e.g. hearburn, 28-5-2010 epigsatric discomfort, or their equivalents) that are often associated with dyspepsia, peptic ulcers, gastritis, peptic esophagitis and indicated for relief of flatulence 112. Aspirin 75mg + Simwastatin 10mg/20mg + Lisimopril 5mg/ 10mg + Atenokol 25mg/50mg tablets 113. Irsogladine Maleate Tablet 2mg/4mg For geacondary prevention of coronory heart disease/stroke in patients where use of such combination is appropriate 114. Lacusamide film coated tablet 50mg/100mg/150mg/200mg As an adjunctive treatment of partial onset seizures in patients 115. Carmustine for inj. 100 mg Indicated as pallitative therapy as a single agent or in established combination therapy with other approved chemotherapetic agents in the following: 1. Brain tumors—glioblastoma, astrocytoma, ependymoma, and metastatic brain tumors. 2. Multiple myeloma—in combination with prednisone. 3. Hodgkin's Disease—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's Iymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's Iymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy. 4. Non-Hodgkin's Iymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy. 4. Non-Hodgkin's Iymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy. 5. For the relief of symptoms of seasonal alle | 108. | 1 | exercise and the single agents and the second line therapy with | 28-5-2010 |
| Activated Dimethicone 50mg + Magnesium Hydroxide 250 mg + Sorbitol 250mg+ Dried Aluminium Hydroxide 250 mg + Sorbitol 250mg+ Ministry 250mg+ Dried Aluminium Hydroxide 250 mg + Sorbitol 250mg+ Ministry 250mg+ Min | 109. | & 2 tablets of FDC of Pyrimethamine 25mg + Sulphadoxine | Additional Strength | 28-5-2010 |
| 250mg+ Dried Aluminium Hydroxide250 mg + Sorbitol (70%)1.25 gm/5ml Solution associated with dyspepsia, peptic ulcers, gastritis, peptic esophagitis and indicated for relief of flatulence sophagitis and indicated for relief of suprovement of gastric microscopic spatial sophagitis and indicated for relief of suprovement of gastric microscopic spatial sophagitis and indicated for relief of suprovement of gastrointestinal symptoms (nause and vomitting) associated with ematogenic cancer chemotherapy | 110. | _ | | 28-5-2010 |
| 10mg + Atenolol 25mg/50mg tablets patients where use of such combination is appropriate 113. Irsogladine Maleate Tablet 2mg/4mg For gastric ulcers, improvement of gastric mucosal lesion (crosion, hemorrhage, redness and edema) caused by acute gastritis and acute exacerbation stage of chronic gastritis. 114. Lacosamide film coated tablet 50mg/100mg/150mg/200mg As an adjunctive treatment of partial onset seizures in patients > 10-6-2010 > 17 years of age. 115. Carmustine for inj. 100 mg Indicated as palliative therapy as a single agent or in established combination therapy with other approved chemotherapeutic agents in the following: 1. Brain tumors—glioblastoma, brainstem glioma, medulloblastoma, astrocytoma, ependymoma, and metastatic brain tumors. 2. Multiple myeloma—in combination with prednisone. 3. Hodgkin's Disease—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's lymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg, Monteleukast Sodium For the relief of symptoms of seasonal allergic rhinitis in adults and pediatric patients 2 years of age and older. granule(Additional Indication) 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | 111. | 250mg+ Dried Aluminium Hydroxide250 mg + Sorbitol | epigastric discomfort, or their equivalents) that are often associated with dyspepsia, peptic ulcers, gastritis, peptic | 28-5-2010 |
| (erosion, hemorrhage, redness and edema) caused by acute gastritis and acute exacerbation stage of chronic gastritis. 114. Lacosamide film coated tablet 50mg/100mg/150mg/200mg As an adjunctive treatment of partial onset seizures in patients 517 years of age. 10-6-2010 | 112. | | | 1-6-2010 |
| Solution | 113. | Irsogladine Maleate Tablet 2mg/ 4mg | (erosion, hemorrhage, redness and edema) caused by acute | 5-6-2010 |
| established combination therapy with other approved chemotherapeutic agents in the following: 1. Brain tumors—glioblastoma, brainstem glioma, medulloblastoma, astrocytoma, ependymoma, and metastatic brain tumors. 2. Multiple myeloma—in combination with prednisone. 3. Hodgkin's Disease—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's lymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg; Monteleukast Sodium oral granule(Additional Indication) 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | 114. | Lacosamide film coated tablet 50mg/100mg/150mg/200mg | | 10-6-2010 |
| medulloblastoma, astrocytoma, ependymoma, and metastatic brain tumors. 2. Multiple myeloma—in combination with prednisone. 3. Hodgkin's Disease—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's lymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg; Monteleukast Sodium oral granule(Additional Indication) For the relief of symptoms of seasonal allergic rhinitis in adults and pediatric patients 2 years of age and older. 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | 115. | Carmustine for inj. 100 mg | established combination therapy with other approved | 10-6-2010 |
| 3. Hodgkin's Disease—as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's lymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg; Monteleukast Sodium For the relief of symptoms of seasonal allergic rhinitis in adults and pediatric patients 2 years of age and older. 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | | | medulloblastoma, astrocytoma, ependymoma, and metastatic | |
| with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 4. Non-Hodgkin's lymphomas—as secondary therapy in combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg; Monteleukast Sodium 4/5mg Chewable Tablet; Monteleukast Sodium Oral granule(Additional Indication) For the relief of symptoms of seasonal allergic rhinitis in adults and pediatric patients 2 years of age and older. 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | | | 2. Multiple myeloma—in combination with prednisone. | |
| combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. 116. Monteleukast Sodium Tablet 10mg; Monteleukast Sodium 4/5mg Chewable Tablet; Monteleukast Sodium Oral granule(Additional Indication) For the relief of symptoms of seasonal allergic rhinitis in adults and pediatric patients 2 years of age and older. For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | | | with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to | |
| 4/5mg Chewable Tablet; Monteleukast Sodium Oral granule(Additional Indication) 117. Ramosetron Dispersible Tablet 0.1 mg For the prevention and treatment of gastrointestinal symptoms (nausea and vomitting) associated with ematogenic cancer chemotherapy | | | combination with other approved drugs for patients who relapse while being treated with primary therapy, or who fail | |
| (nausea and vomitting) associated with ematogenic cancer chemotherapy | 116. | 4/5mg Chewable Tablet; Monteleukast Sodium Oral | | 12-6-2010 |
| 118. Amitraz 2% w/v pour on solution For the control of ticks on cattle and ostriches 12-6-2010 | 117. | Ramosetron Dispersible Tablet 0.1 mg | (nausea and vomitting) associated with ematogenic cancer | 12-6-2010 |
| | 118. | Amitraz 2%w/v pour on solution | For the control of ticks on cattle and ostriches | 12-6-2010 |

| 119. | Flupirtine Maleate Capsules 100mg | For treatment of acute and chronic pain, i.e, for painful increased muscle tone of the posture and motor muscles, primary headache, tumor pain, dysmenorrhea and pain after traumatologic/orthopaedic operations and injuries. | 12-6-2010 |
|------|---|--|-----------|
| 120. | Lidocaine Patch 350mg (Additional Pack Size) | For the relief of pain associated with post-herpetic neuralgia. It should be applied only to intact skin. | 17-6-2010 |
| 121. | Oseltamivir Phosphate (Additional Indication) | In the treatment of children 6 to 12 months of age during a pandemic influenza outbreak. | 19-6-2010 |
| 122. | Methylcobalamin 1500 mcg + Folic Acid 0.7 mg+Niacinamide 12mg injection | For the treatment of diabetic neuropathy in adults | 23-6-2010 |
| 123. | Methylcobalamin 1500mcg + Pyridoxine 100mg + Nicotinamide 100mg injection | For the treatment of diabetic neuropathy in adults | 23-6-2010 |
| 124. | 1 tablet of Tenofovir Disoproxil 300mg + Lamivudine 300mg & 2 tablet of Nevirapine 200mg combipack | For the treatment of HIV infection in adults | 23-6-2010 |
| 125. | Vitamin A (oily form) as Palmiate 2000IU + Vitamin B1 1mg + Vitamin B2 1mg + Vitamin B6 1.34mg + Vitamin B12 1mcg + Ascorbic Acid 33.3mg + Vitamin D3 200IU + Folic Acid 0.3mg + Calcium Pantothenate 1mg + Niacinamide 15mg + Dibasic Calcium Phosphate 100mg + Phosphorus 77mg+ Ferrous Fumarate 50mg + Copper 0.01mg+ Potassium Iodide 0.0015mg + Heavy Magnesium Oxide 1mg + Manganse Sulphate 0.01mg + Potassium Sulphate 1mg + Zinc Sulphate 0.15mg Capsule | | 23-6-2010 |
| 126. | Ramipril MR Capsules 2.5/5 mg (Additional Strength) | Same as Approved | 3-7-2010 |
| 127. | Sitagliptin Phosphate 25/50/100 mg Tablet (Additional Indication) | (i) Use of Sitagliptin Phosphate in combination with Metformin and a PPARγ agonist as an adjunct to diet & exercise in adult patients with Type-2 Diabetes mellitus who are inadequately controlled on combination therapy with Metformin and a PPARγ agonist. (ii) Use of Sitagliptin Phosphate in combination with insulin, alone or in combination with Metformin. | 3-7-2010 |
| 128. | Erlotinib HCl 25/100 mg Tablet (Additional Indication) | As monotherapy for maintenance treatment in patients with locally advanced or metastatic non-small cell lung cancer with stable disease after 4 cycles of standard platinum based first line chemotherapy. | 3-7-2010 |
| 129. | Gemcitabine HCl injection 2gm/vial (Additional Strength) | Same as Approved | 3-7-2010 |
| 130. | Etizolam 0.25 mg tablet (Additional Strength) | Same As Approved | 3-7-2010 |
| 131. | Dexketoprofen 1.25% w/w + Linseed Oil 3.0%w/w +Menthol 5%w/w + Methyl Salicylate 10%w/w (12.5mg) | For the topical treatment of acute musculoskeletal pain | 6-7-2010 |
| 132. | Cefuroxime Sodium 750/ 1500mg +Sulbactam Sodium 375/750mg | For the treatment of mild to moderate lower respiratory tract infection | 6-7-2010 |
| 133. | HPMC 0.3% +Dextran 0.1% + Glycerin 0.2% | For the temporary relief of burning and irritation due to dryness of the eye and for use as a protection against further irritation. For the temporary relief of discomfort due to minorirritations of the eyes or to exposure to wind or sun | 6-7-2010 |
| 134. | Cyanocobalamin 15mcg + Chromium Picolinate USP 250mcg + Folic Acid USP 1500 mcg+Nicotinamlide IP100 mg + Pyridoxine 3mg +Selenius Acid USP 100mcg+Zinc Sulfate Monohydrate USP 61.8 mg eq. to 22.5 mg | For Mineral & Vitamins deficiency states in adult patients | 6-7-2010 |

| 135. | Methylparaben 0.15%w/w + Propyl Paraben 0.05%w/w + Glycerin 20%w/w +CMC 1.74%w/w +Allantoin 0.5%w/w +Benzyl Alcohol 0.5%w/w. | For short term treatment of chronic wounds including Stage III & IV pressure sores, leg ulcers | 6-7-2010 |
|------|--|--|-----------|
| 136. | Clonazepam IR 0.5mg/0.5mg+ Paroxetine CR 12.5/25mg | For the treatment of patients with co-morbid depression and axiety | 8-7-2010 |
| 137. | Nilotinib Capsules 200mg | For the treatment of chronic and accelerated phase Philadelphia chromosome positive chronic myelogenous leukemia (CML) in adult patients resistant to or intolerant to prior therapy that included imatinib. | 10-7-2010 |
| 138. | Imipenem 50mg + Clistatin 50mg | Treatment of infection caused by susceptible bacteria for the following indication as in neonates and infants (below 3 months of age) | 14-7-2010 |
| | | 1. Lower respiratory tract infections | |
| | | 2. Urinary tract infections | |
| | | 3. Intra-abdominal infections | |
| | | 4. Bacterial septicaemia | |
| | | 5. Bone and joint infections | |
| | | 6. Skin and Skin structure infections | |
| | | 7. Bacterial endocarditis | |
| 139. | Lidocaine Lozenges 100mg (Additional Strength) | Same as approved | 15-7-2010 |
| 140. | Metoprolol Succinate ER 25/50 mg+Telmisartan 20/40mg | For the treatment of essential hypertension | 16-7-2010 |
| 141. | Etodolac 300/300mg + Thiocolchicoside 4/8mg | For the treatment of patients with acute painful musculoskeletal conditions | 19-7-2010 |
| 142. | Celecoxib 100/200mg + Diacerein 50/50mg | For the treatment of adult patients with osteoarthritis of knee and/or hip joints | 19-7-2010 |
| 143. | Metoprolol succinate ER 25mg+50mg+Olmesartan 20/20mg | For the treatment of essential hypertension | 19-7-2010 |
| 144. | 1 tablet of Atazanavir 300mg & 1 tablet of Ritonavir 100mg & FDC of Tenofovir Disoproxil Fumarate 300mg + Lamivudine 300mg (Each Co-pack contains:— 1 capsule of Atazanavir 300mg & 1 Tablet of Ritonavir 100mg & 1 Tablet of FDC of Tenofovir Disoproxil Fumarate 300mg + Lamivudine 300mg) | For the treatment of HIV infections in adults | 19-7-2010 |
| 145. | Gliclazide 40/80mg+Metformin 500mg+Pioglitazone15mg | Additional strength | 19-7-2010 |
| 146. | Lidocaine HCl Ophthalmic Sterile Gel 35 mg/ml | Indicated for ocular surface anaesthesia during ophthalmologic procedures. | 30-7-2010 |
| 147. | Naftopidil Tablets 25/50/75mg | For the treatment of dysuria associated with benign prostratic hyperplasia. | 30-7-2010 |
| 148. | Metformin 500/850/1000mg+ Sitagliptin Phosphate 50/50/50 mg (Additional Indication) | (A) FDC is indicated as triple combination therapy with a PPARy agonist (i.e a thiazolidinedione) as an adjunct to diet and exercise in patients inadequately controlled on their maximal tolerated dose of Metfomin and a PPARy agonist. | 2-8-2010 |
| | | (B) FDC is also indicated as add-on to insulin(i.e triple combination therapy) as an adjunct to diet and exercise to improve glycemic control in patients when stable dosage of insulin and metformin alone do not provide adequte control | |
| 149. | Bimatoprost Eye Drops 0.1mg/ ml | For the treatment of hypotrichosis of the eye-lashes by increasing their growth including length, thickness & darkness | 5-8-2010 |
| 150. | S(+) Etodolac 5/10%w/w + Camphor 4%w/w + Menthol 10%w/w + Methyl Salicylate 5%w/w + Linseed oil 3%w/w | For the painful & inflammatory conditions associated with joint, tendons, ligaments and muscles | 5-8-2010 |

| | T | | |
|------|---|--|-----------|
| 151. | Cefpodoxime 100/200mg+Dicloxacillin ER 500/500mg | For the treatment of lower RI, Skin and soft tissue, bone and joint infections and ENT infections in adult patients prone to antibiotic induced diarrhoea | 6-8-2010 |
| 152. | Voglibose0.3 + Metformin (SR) 500mg | Additional higher strength | 6-8-2010 |
| 153. | Amoxycillin Oral Suspension 200/400 mg per 5ml | In the treatment of infections due to susceptible strains of the designated micro-organisms in the condition of E.N.T infection, skin & skin structure infection, lower respiratory tract infection, genito-urinary tract infection, gonorrhea, acute uncomplicated (anogenital & urethral) infections | 9-8-2010 |
| 154. | Olmesartan 40mg + Amlodipine 5mg (Additional Strength) | Additional indication: | 9-8-2010 |
| | | For the treatmenmt of hypertension alone or with other antihypertensive agents | |
| | | 2. To use as initial therapy in patients who are likely to need multiple antihypertensive agents to achieve their blood pressure goals | |
| 155. | Doripenem 500 mg powder for injection (Additional Indication) | Complicated intra-abdominal infections and complicated urinary tract infections, including complicated and uncomplicated pyelonephritis and cases with concurrent bacteremia | 10-8-2010 |
| 156. | Nadolol tablet USP 20/40/80 mg | In the treatment of Angina-pectoris, Cardiac Arrythmias & essential hypertension | 20-8-2010 |
| 157. | Fluconazole for Oral Suspension 50/200 mg per 5ml (Pack Size : 35ml) | For the treatment of systemic candidiasis, mucosal candidiasis, prevention of fungal infections in patients with malignancy | 20-8-2010 |
| 158. | Metformin 500/500mg + Repaglinide 1/2mg | Indicated as an adjunct to diet and exercise to improve glycemic control in adults with Type 2 Diabetes mellitus who are already treated with a Meglitinide & Metformin or who have inadequte glycemic control on a Meglotinide alone or Metformin | 20-8-2010 |
| 159. | Hydroxyzine Hydrochloride SR Tablets 25/50/75mg | For management of pruritus due to allergic conditions such as chronic urticaria and atopic contact dermatoses, and in histamine - mediated pruritus | 20-8-2010 |
| 160. | Rosuvastatin 5/10/20mg+ Fenofibrate 67/145/160mg | For the treatment of combined hyperlipidemia in patients with normal hepatic and renal function | 23-8-2010 |
| 161. | Fexofenadine 120mg + Montelukast 10mg | For treatment of allergic rhinitis in adults only | 31-8-2010 |
| 162. | Irinotecan HCl Trihydrate Injection 20 mg/ml (Pack Size 25ml) | For the treatment of metastatic colorectal cancer. | 31-8-2010 |
| 163. | Cefixime SR 200mg/400mg + Ofloxacin SR 200mg/400mg Tablet | Additional strength | 31-8-2010 |
| 164. | Ademetionine enteric coated Tablets200mg/400mg | For the management of intrahepatic cholestasis and liver disease | 1-9-2010 |
| 165. | Memantine HCl ER Tablet 28mg | in the treatment of moderate to severe Alzheimer's disease. | 6-9-2010 |
| 166. | Buserelin Acetate injection 1mg/ml (Additional Indication) | indicated for the palliative (initial & maintenance treatment) of patients with hormone-dependent advanced carcinoma of the Prostate gland (Stage D). | 8-9-2010 |
| 167. | Etodolac 400mg/400mg + Thiocolchicoside 4mg/8mg tablet | Additional higher strength | 9-9-2010 |
| 168. | Montelukast Sodium 10 mg Tablet, Montelukast Sodium 4/5 mg Chewable Tablet and Montelukast Sodium Oral Granules (Additional Indication) | For the relief of perennial allergic rhinitis in adults & pediatrics patients 6 months of age and older | 11-9-2010 |

| 169. | Acyclovir SR Tablet 1200 mg | For the treatment of uncomplicated Herbis Zoster in immunocompetent patients | 11-9-2010 |
|------|---|---|------------|
| 170. | Lornoxicam 4mg + Eperisone 50mg tablet | For the improvement of myotonic conditions caused by neck shoulder arm syndrome, scapulumeral periarthiritis and low back pain in adult patients only | 13-9-2010 |
| 171. | Pirfenidone film coated tablet200mg | For the treatment of idiopathic pulmonary fibrosis. | 14-9-2010 |
| 172. | Beclomethasone Dipropionate IP 0.025% w/w + Neomycin Sulphate IP 0.5% w/w cream | Additional lower strenth | 15-9-2010 |
| 173. | Amlodipine 5mg/2.5mg + Valsartan 320mg/80mg + HCTZ 12.5mg/12.5mg tablet | Additional strength | 21-9-2010 |
| 174. | Paracetamol 500mg/500mg +Thioclchicoside 4mg/8mg tablet | For the treatment of acute inflammation conditions associated with spasm in adults only | 21-9-2010 |
| 175. | Ketorolac Tromethamine 0.4% w/v +Moxifloxacin 0.5% w/v Opthalmic solution | Additional lower strenth | 21-9-2010 |
| 176. | Nilotinib Capsules 150 mg. | Treatment of newly diagnosed adult patients with Philadelphia chromosome positive chronic Myeloid Leukemia (Ph+CML) in chronic phase | 29-9-2010 |
| 177. | Levocloperastine (as Fendizoate) Suspension eq. to Levocloperastine HCl 20mg/5ml | For the treatment of non-productive cough in adult patients | 29-9-2010 |
| 178. | Zileuton uncoated bilayered extended release tablet600mg | For the prophylaxis and chronic treatment of asthma in adults | 30-9-2010 |
| 179. | Etodolac 400mg + Paracetamol IP 500mg tablet | Additional higher strength | 1-10-2010 |
| 180. | Dried Aluminium Hydroxide Gel IP 600mg +Magnesium Hydroxide IP 300mg +Simethicone IP 25mg +Oxetacaine BP 10mg per 5ml syrup | For the treatment of symptoms of functional gastrointestinal disorders like irritable bowel syndrome, functional dyspepsia, peptic ulcer, gasteroesophageal reflux disease which include smooth muscle spasms, flatulence, abdominal distension, hyperacidity, gastric distress, bloting etc. | 5-10-2010 |
| 181. | S(-) Metoprolol Succinate ER eq. to S (-) Metoprolol Tartarate 25mg/50mg + Clopidogrel (IR) 75mg/75mg tablet | For the treatment of essential hypertension who need antiplatelet therapy especially for prophylaxis of atherombotic event | 7-10-2010 |
| 182. | Exenatide Injection (Additional Indication) | Exenatide is a glucagon like peptide-1 (GLP-1) receptor agonist indicated as an adjunct to diet and exercise to improve glycemic control in adults with Type 2 Diabetes mellitus | 9-10-2010 |
| 183. | Bimatoprost Ophthalmic Solution 0.01% w/v (Additional Strength) | Same as approved | 9-10-2010 |
| 184. | Drosperinone + Ethinylestradiol tablets Same as approved | For the treatment of moderate Acne vulgaris in adult females, who have no known contra-indications to oral contraceptive therapy, and have achieved menarche. It should be used for the treatment of acne only if the patient desires an oral contraceptive for birth control | 9-10-2010 |
| 185. | Etodolac USP 400mg + Tolperisone HCl 150mg tablet | For the treatment of acute inflammation conditions associated with spasm in adults only | 11-10-2010 |
| 186. | Citicoline 500mg +Piracetam 400mg per 5 ml syrup | Additional dosage form | 11-10-2010 |
| 187. | Thiocolchicoside 0.25% +Methyl Salicylate IP 100mg +Menthol IP 50mg +Absolute alcohol IP 50mg per gm gel | For the topical treatment of painful muscles spasms of skeletal musculature | 11-10-2010 |
| 188. | Amlodipine Besylate IP Eq. to Amlodipine 10mg/5mg/2.5mg +Telmisartan 80mg/40mg/20mg + HCTZ 25mg/12.5mg/12.5mg tablet | For the treatment of essential hypertension | 12-10-2010 |
| 189. | Pregabalin 150mg/75mg + Methylcobalamin 750mcg + Alpha Lipoic Acid 100mg + Pyridoxine 3mg +Folic Acid 1.5mg capsule | For the treatment of painful diabetic neuropathy in adults only | 12-10-2010 |

| 190. | Ferrous Bisglycinate 60mg + Zinc Sulphate monohydrate Eq. to Zinc 14mg + Copper as Copper Sulphate 1 mg + Selenium as Selenomethionine 70mcg + Vitamin B6 IP 10mg + Methylcobalamin 1500mcg + Folic Acid 5mg + Fish oil 10/40 (DHA 200mg) 500mg capsule. | For Vitamin and mineral deficiency | 13-10-2010 |
|------|--|---|------------|
| 191. | Combipack of 21 tablets of FDC of Ethinylestradiol IP 0.03 mg + Levonorgestrel IP 0.15 mg and 7 tablets of Ferrous Fumarate IP 75mg (Eq. to 24.375 mg of Ferrous iron) | As contraceptive pill for women | 13-10-2010 |
| 192. | Amitriptyline Hydrochloride IP 4%w/w + Ketamine HCL eq. to Ketamine 2% w/w per gm cream. | For the treatment of patients suffering from neuralgia | 13-10-2010 |
| 193. | Paracetamol IP 500mg + Phenylephrine HCL IP 5mg +Caffeine (anhydrous) IP 30mg + Diphenhydramine HCL IP 25mg tablet. | For the treatment of fever associated with cold, nasal congestion in adults | 13-10-2010 |
| 194. | Levocetirizine Dihydrochloride 2.5mg +Montelukast 5mg dispersible tablet | Additional strength | 13-10-2010 |
| 195. | Calcium Citrate Maleate eq. to elemental Calcium 250mg + Vitamin D3 IP 200 IU tablet | For the treatment of calcium and Vitamin D deficiency states | 13-10-2010 |
| 196. | Levonorgesterel 100mcg + Ethniyl Estradiol IP 20mcg tablet | (Additional indication) For the treatment of moderate Acne Vulgaris in women more than 18 years of age who have no therapy contraindications to oral contraceptive therapy, desire contraception, and have achieved menarche | 13-10-2010 |
| 197. | Eberconazole 1% w/w + Mometasone Furate USP 0.1% w/w cream | For the treatment of mild to moderate inflammed cutaneous mycoses | 13-10-2010 |
| 198. | Lubiprostone Soft Gelatin Capsules8mcg/24mcg | For the treatment of chronic constipation in adults. For the treatment of irritable bowel syndrome with constipation (IBS-C) in women > 18 years old | 16-10-2010 |
| 199. | Pazopanib (as Hydrochloride) film coated Tablets200mg/400mg | For the treatment of patients with advanced renal cell carcinoma | 16-10-2010 |
| 200. | Eltrombopag (as Olamine) Tablets25mg / 50mg | For the treatment of thrombocytopenia in patients with chronic immune (idiopathic) thrombocytopenic purpura who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy | 16-10-2010 |
| | | (It should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding. It should not be used in an attempt to normalize platelet counts) | |
| 201. | Tolperisone HCl 150mg + Paracetamol IP 500mg tablet | For the treatment of patients with acute painful musculoskeletal conditions | 21-10-2010 |
| 202. | Zaltoprofen film coated tablet80mg JP | For anti-inflammatory and analgesic activity in rheumatoid arthritis, osteoarthritis, low back pain, periarthritis of shoulder, cervicobrachial syndrome, postoperative, post-trauma, and post-dental extraction | 31-10-2010 |
| 203. | Hyoscine Butylbromide 20 mg tablet (Additional Strength) | Same as approved | 4-11-2010 |
| 204. | Cerebroprotein Hydrolysate injection (Lyophilised)1050 mg/2100mg (approx) equivalent to nitrogen30mg/ 60mg respectively | For amelioration of cranial injury, cerebrovascular pathological sequelae and aprosexia in dementia | 4-11-2010 |
| 205. | Clebopride (as Malate) (1) Tablets. 0.5 mg Ph. Eur. (2) Syrup 0.1 mg/ml Ph. Eur. (3) Injection 1 mg/2ml Ph. Eur. | For the treatment of various disorders associated with altered gastrointestinal motility like non-ulcer dyspepsia (NUD) and / or chronic gastritis, gastroesophageal reflux disease (GERD), nausea and vomiting of organic or functional origin | 4-11-2010 |

| 206. | Tolperisone HCl Mouth Dissolving tablet 50/100 mg (Additional Dosage Form) | Same as Approved | 4-11-2010 |
|------|--|---|------------|
| 207. | Oxybutynin Gel 100 mg | For the treatment of overactive bladder with the symptoms of urge urinary incontinence, urgency and urinary frequency | 4-11-2010 |
| 208. | Dapoxetine (as Hydrochloride) Tablets 30mg/60mg | For the treatment of premature ejaculation (PE) in men 18 to 64 years of age | 13-11-2010 |
| 209. | Opipramol dihydrochloride Tablet 50mg/100mg | For the treatment of Generalised Anxiety Disorder (GAD) and Somatic Disturbances | 13-11-2010 |
| 210. | Ondansetron Rapid Film Oral Strip 4/8 mg | For chemotherapeutic induced nausea and vomiting | 16-11-2010 |
| 211. | Acebrophylline SR Tablet 200 mg | For the treatment of adult patients with Chronic Obstructive Pulmonary Disease (COPD) and bronchial asthma | 23-11-2010 |
| 212. | Ammonium Lactate Cream 12% w/w | Indicated for the treatment of Dry, scaly skin(xerosis) and ichthyosis vulgaris and for temporary relief of itching associated with these condition. | 3-12-2010 |
| 213. | Rabeprazole Sodium Powder oral suspension (sachet of 20 mg) (Additional Dosage Form) | Same as approved | 3-12-2010 |
| 214. | Temsirolimus Concentrate for Infusion for Injection 25mg/ml | For the treatment of advanced renal cell carcinoma | 11-12-2010 |
| 215. | Acetyl Cysteine Eye Drops 5% w/v | Indicated the relief of Dry eye symptoms associated with defecient tear secretion, impaired or abnormal mucus production | 14-12-2010 |
| 216. | Clindamycin Palmitate HCl for Oral Solution 75mg/5ml | Indicated for the Prophylaxis against alpha-hemolytic(viridans group) sreptococci before dental, oral and upper respiratory tract surgery | 14-12-2010 |
| 217. | Deferasirox Tablet for Oral Suspension (Additional Dosage Form) | Same as approved | 14-12-2010 |
| 218. | Ramelteon film coated tablets 8 mg | For the treatment of insomnia characterized by difficulty with sleep onset | 15-12-2010 |
| 219. | Memantine ER Tablet 7mg/14 mg/21mg (Additional Strength) | Same as approved | 17-12-2010 |
| 220. | Solution for haemofiltration and haemodialysis (Each unit contains two compartments A & B. Before reconstitution: | For treatment of acute kidney disease (renal failure), as substitution solution in continuous haemofiltration and haemodiafiltration and as a dialysis solution in continuous | 23-12-2010 |
| | • 1000 ml of electrolyte solution in Compartment A contains: | haemodialysis | |
| | Calcium Chloride 2H2O 5.145gm | | |
| | Magnesium Chloride 2H2O 2.033gm Lactic Acid 5.4gm • 1000ml of buffer solution in Compartment B contains: | | |
| | Sodium Chloride 6.45gm | | |
| | Sodium Bicarbonate 3.090gm) | | |
| 221. | Pyridoxine HCl IP 3mg + Nicotinamide IP 100mg + Cyanocobalamin IP 15mcg + Folic Acid IP 500mcg + Chromium Picolinate USP 250mcg Tablet | For Mineral & Vitamins deficiency states in adult patients | 23-12-2010 |
| 222. | Rosuvastatin Calcium IP 5mg/10mg + Fenofibrate BP 160mg/160mg tablet | Additional strength | 29-12-2010 |
| 223. | Chlorhexidine Gluconate IP 1%w/v + Ethyl Alcohol IP 61%w/v solution | Surgical hand scrub & healthcare personnel hand wash | 29-12-2010 |

Page **49** of **96**

| Sl. No. | Name of Drug | Indication | Date of issue |
|---------|---|---|---------------|
| 1. | Alpha Lipoic Acid USP 100mg + Methylcobalamin 1500mcg + Vitamin B6 IP 3mg + Folic Acid IP 1.5mg + Benfotiamine 50mg + Biotin USP 5mg + Chromium Picolinate USP Eq. to Chromuim 200mcg Capsule | For the treatment of diabetic neuropathy | 3-1-2011 |
| 2. | Ropinirole ER Tablet 1 mg. (Additional Strength) | Same as approved | 4-1-2011 |
| 3. | Erlotinib HCl Tablet 150 mg (Additional Indication) | Monotherapy for the Maintenance Treatment of Patients with locally advanced metastatic non-small lung cancer whose disease has not progressed after four cycles of Platinum based First Line Chemotherapy | 4-1-2011 |
| 4. | Moxifloxacin HCl BP 0.5% w/v + Bromfenac Sodium 0.09% w/v Eye drop | For the reduction of post operative inflammatory conditions of the eye | 5-1-2011 |
| 5. | Ferrous Ascorbate 100mg + Folic Acid IP 1.5mg + Cyanocobalamin IP 15mcg + Zinc Sulphate Monohydrate Eq. to Elemental Zinc 22.5mg Tablet | For the treatment of iron deficiency anaemia | 5-1-2011 |
| 6. | S (+) Etodolac 300mg +Thiocolchicoside 8mg Tablet | For the treatment of patients with acute painful musculoskeletal conditions | 5-1-2011 |
| 7. | Beclomethasone Dipropionate IP 100mcg + Formoterol Fumarate Dihydrate BP Eq. to Formoterol Fumarate 6 mcg Metered Dose Inhaler | For the treatment of bronchial asthma where use of inhaled corticosteroid therapy found appropriate | 5-1-2011 |
| 8. | Amlodipine Besilate IP Eq. to Amlodipine 5mg + Indapamide USP SR 1.5mg Tablet | For the treatment of mild to moderate hypertension | 5-1-2011 |
| 9. | Metformin HCl IP 500mg + Alpha Lipoic Acid USP 200mg Tablet | For the treatment of patients with diabetic polyneuropathy | 5-1-2011 |
| 10. | Spironolactone IP 25mg + Furosemide IP 40mg Tablet | Additional strength | 5-1-2011 |
| 11. | Bicalutamide Tablet 150 mg.(Additional Strength and Indication) | Indicated either alone or as adjuvant to radical Prostatetomy or radiotherapy in Patients with Locally advanced Prostate cancer at High risk for disease Progression | 5-1-2011 |
| 12. | Paracetamol Chewable Tablet 500 mg. (Additional Dosage Form) | Same as approved | 5-1-2011 |
| 13. | Methylcobalamin 500mcg + Folic Acid IP 1.5mg + Zinc Sulphate IP 25mg + Chromium Polynicotinate 200mcg + Selenium as Sodium Selenite USP 65 mcg + Inositol 100mg + Alpha Lipoic Acid 50mg + Streptococcus faecalis 30 Millions (30 lakh) + Clostridium butyricum 2 Million (20 lakh) + Bacillus mesentericus 1 Million (10 lakh) + Lactic acid bacillus 50 Million (500lakh) Capsule | I | 11-1-2011 |
| 14. | Dextromethorphan Extended Release Oral Suspension 30 mg per 5 ml | For the Relief of Persistent Dry irritating cough in adult patients Only | 17-1-2011 |
| 15. | Hydroquinone USP 2% w/w + Tretinoin USP 0.025% w/w + Allantoin USP 1%w/w Gel | For treatment of hyper pigmentry disorders | 31-1-2011 |
| 16. | Paracetamol Fast Release Tablet 500mg (Additional Dosage Form) | Same as approved | 10-2-2011 |
| 17. | Each bowel preparatory kit contains: One enteric coated tablet Biscodyl IP 5mg & One sachet of powder for reconstitution contains: Polyethylene glycol 3350NF: 210gm + Sodium Bicarbonate IP 2.86gm + Sodium Chloride IP 5.6gm + Potassium Chloride IP | For bowel cleansing prior to colonoscopy in adult patients only | 11-2-2011 |

| | 0.74gm | | |
|-----|--|---|-----------|
| 18. | Ferric Carboxymaltose equivalent to elemental Iron 50mg/ml Injection | For the treatment of iron deficiency in adults when oral iron preparations are ineffective or cannot be used | 11-2-2011 |
| 19. | Pidotimod Tablet 400 mg/800 mg & Oral Solution 400 mg/800 mg per 7ml | For infections of the respiratory system in secondary and primary immunodeficiency with alteration in maturation of T cells in adults only | 11-2-2011 |
| 20. | Ursodeoxycholic Acid Suspension 125mg/5ml and 250 mg /5ml. | For the treatment of patients with chronic cholestatic liver disease | 11-2-2011 |
| 21. | Iloperidone Tablets 1mg/2mg/4mg/6mg/8mg/10mg/12mg | For the acute treatment of adults with schizophrenia | 17-2-2011 |
| 22. | Calf lung surfactant Intratracheal Suspension containing Phospholipids35mg (containing 26 mg phosphatidylcholine of which 16mg is disaturated phosphatidylcholine) Proteins .0.65mg (including 0.26mg of surfactant associated proteins B) per ml | (a) As a prophylaxis therapy for premature infants <29 weeks of gestational age at significant risk for Respiratory Distress Syndrome (RDS) (b) For the treatment of infants <72 hours of age with RDS (confirmed by clinical and radiologic findings) and requiring endotrachteal intubation | 17-2-2011 |
| 23. | Baclofen ER Capsule 10 mg/50 mg. (Additional Strength) | Same as approved | 17-2-2011 |
| 24. | Dexketoprofen Trometamol ER Tablet 50 mg (Additional Strength) | Same as approved | 21-2-2011 |
| 25. | Gatifloxacin Eye drop 5 mg/ml | For the treatment of bacterial conjunctivitis caused by susceptible strains | 21-2-2011 |
| 26. | Paclitaxel Injection 30mg/100 mg/300 mg (Additional Indication) | Advanced squmous cell carcinoma of Head and Neck SCCHN: | 24-2-2011 |
| | | 1. Diagnosed but unrectable locally advanced SCCHN | |
| | | 2. Recurrent SCCHN, if surgery or radiation is not feasible | |
| | | 3. Palliative treatment of metastatic SCCHN | |
| 27. | Eslicarbazepine Acetate Tablets 200mg/400mg/600mg/800mg | As an adjunctive therapy in adults with partial-onset seizures with or without secondary generalisation | 7-3-2011 |
| 28. | Ketoprofen Plaster (Size - 7cm x 10 cm) | For the Relief of Musculoskeletal pain and Inflammation | 9-3-2011 |
| 29. | Meropenem IP 1000mg + Sulbactam sodium USP 500mg injection | For the treatment of lower respiratory tract infection caused by gram negative bacteria in adults only | 15-3-2011 |
| 30. | Dasatinib 20mg/50mg/70mg (Additional Indication) | Treatment of newly Diagnosed Adults with Chronic Myeloid Leukaemia (CML) in Chronic Phase | 16-3-2011 |
| 31. | Adenosylcobalamin 15mcg + Ferrous Ascorbate 100mg + Folic Acid IP 1.5mg + Zinc Sulphate USP Eq. to Elemental Zinc 22.5mg tablet | For the treatment of iron deficiency anaemia in adults only | 21-3-2011 |
| 32. | Tioconazole Vaginal Gel 65mg per gm | For the local treatment of vulvo vaginal candidiasis in adult patients | 22-3-2011 |
| 33. | Aspirin (EC Tablets) 75mg/ 75mg/150mg + Rosuvastatin (Granules) 5mg/10mg/10mg Capsules | For the treatment of dyslipidemia associated with arthersclerotic arterial disease with risk of Myocardial infarction, stroke or peripheral vascular disease | 30-3-2011 |
| 34. | Ilaprazole Tablets 5mg/10mg | For the treatment of duodenal ulcer in adults only | 5-4-2011 |
| 35. | Asenapine Maleate Sublingual Tablets 5mg/10mg | For acute treatment of schizophrenia in adults only | 7-4-2011 |
| 36. | Cefpodoxime IP 100mg/100mg/ 200mg/200mg + Ofloxacin IP 100mg/200mg/100mg/200mg Tablets | For the Treatment of Lower and upper respiratory tract infection and Typhoid fever in adults only | 16-4-2011 |
| 37. | Tapentadol Hydrochloride Tablets 50mg/75mg/100mg | For relief of moderate to severe acute pain in adults 18 years of age or older | 18-4-2011 |

| 38. | Cefixime Trihydrate IP Eq. to Anhydrous Cefixime 100mg/200mg+Ofloxacin IP 100mg/200mg Dispersible Tablet | Same as approved | 20-4-2011 |
|-----|--|---|-----------|
| | (Additional dosage form) | | |
| 39. | S-Bupivacaine Hydrochloride Injection 2.5mg/5mg/7.5mg per ml | For surgical anaesthesia in adults for epidural (including caesarean section), intrathecal, peripheral nerve block | 28-4-2011 |
| 40. | Dexketoprofen 50mg + Dicyclomine 20mg per 2ml Injection | For the treatment of acute colicky pain in adult patients | 28-4-2011 |
| 41. | Gatifloxacin 0.3% + Loteprednol Etabonate 0.5% + Benzalkonium Chloride 0.01% w/v Opthalmic suspension | For Post-Operative Steroid repulsive inflammatory Ocular conditions when Ocular bacterial infections or a risk of bacterial infection exists | 19-5-2011 |
| 42. | Diclofenac Colestyramine 145.6 mg corresponding to 75mg Diclofenac Sodium Capsules. | For the treatment of acute arthritis (including attacks of gout), chronic arthritis especially rheumatoid arthritis (chronic polyarthritis), ankylosing spondylitis (Morbus Bechterew) and other inflammatory, rheumatoid syndromes of vertebral column | 19-5-2011 |
| 43. | Brinzolamide Ophthalmic Suspension 1%w/v | For the treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma | 26-5-2011 |
| 44. | Besifloxacin Ophthalmic Suspension 0.6% w/v | For the treatment of bacterial conjunctivitis | 31-5-2011 |
| 45. | Diclofenac (as EC Tablets) 25mg + Paracetamol 325 mg Capsule | Same as approved | 31-5-2011 |
| | (Additional Strength) | | |
| 46. | Amlodipine 10mg/10mg+ Telmisartan 40mg/80mg Tablets (Additional Strength) | Same as approved | 31-5-2011 |
| 47. | 1. PROBIO: | For the treatment of rotavirus diarrhoea | 9-6-2011 |
| | Each sachet (1g) contains: | | |
| | Total Probiotic Not less than 1.25 Billion CFU (Lactobacillus acidophilus, Lactobacillus rhamnosus, Bifidobacterium longum, Saccharomyces boulardii) | | |
| | 2. PROBIO & PREBIO (F)s | | |
| | Each sachet (1g) contains: | | |
| | Probio (CMP 5) | | |
| | Total ProbioticNot less than 1.25 Billion CFU (Lactobacillus acidophilus, Lactobacillus rhamnosus, Bifidobacterium longum, Saccharomyces boulardii) | | |
| | Fructo Oligo Saccharides 100mg | | |
| | 3. PROBIO & PREBIO (D) | | |
| | Each sachet (1.3g) contains: | | |
| | Probio (CMP 5) | | |
| | Total probiotic not less than 1.25 billion CFU (Lactobacillus acidophilus, Lactobacillus rhamnosus, Bifidobacterium longum, Saccharomyces boulardii) | | |
| | Fructo Oligo Saccharides1000mg | | |
| 48. | Paracetamol 325mg + Tramadol Hydrochloride 37.5 mg Tablets | Additional indication. (For Symptomatic Treatment of Moderate To Severe Pain.) | 21-6-2011 |
| 49. | Silodosin Capsules 2 mg/4mg/ 8mg | For the treatment of signs and symptoms of benign prostatic hyperplasia (BPH) in adults only. | 23-6-2011 |
| 50. | Tiapride Hydrochloride Tablets 25mg/50mg/100mg | For the treatment of agitation and aggressiveness in adult patients with cognitive impairment. | 23-6-2011 |
| 51. | Fexofenadine 120mg + Montelukast 10mg Chewable tablet | Additional dosage form | 7-7-2011 |

| 52. | Each Combipack of 21 tablets contains: | As contraceptive pill for women | 27-7-2011 |
|-----|---|---|-----------|
| | (A) FDC of Norgestrel & Ethinyl estradiol tablets IP Each sugar coated white tablet contains: | | |
| | Norgestrel 0.30mg & Ethinyl estradiol 0.03mg | | |
| | (B) 7 tablets of Ferrous Fumarate: | | |
| | Each sugar coated tablet contains : - Ferrous Fumarate 75mg eq. to 24.375mg of ferrous iron | | |
| 53. | Cefixime (SR) 400mg + Moxifloxacin (SR) 400 mg Tablets | For the treatment of lower respiratory tract infections in adults only | 8-8-2011 |
| 54. | Alpha Lipoic acid 100mg + Folic acid 1.5mg + Methylcobalamin 1500mcg + Pyridoxine Hydrochloride 3mg + Vitamin D3 1000 IU Tablets | For the treatment of Diabetic neuropathy | 10-8-2011 |
| 55. | Levosulpride (SR) 75 mg + Rabeprazole sodium IP (EC Pellets) 20mg Capsule | For the treatment of Gastro-oesophageal Reflux Disease (GERD) | 19-8-2011 |
| 56. | Each Combikit contains | For the treatment of HIV infection in adults. | 23-8-2011 |
| | Part A - One film coated tablet of Tenofovir Disoproxil Fumarate 300mg eq. to Tenofovir 245mg + Emtricitabine 200mg. | | |
| | Part B - One film coated tablet of Atazanavir sulphate eq. to Atazanavir 300mg + Ritonavir 100mg | | |
| 57. | Diclofenac sodium (SR) 100mg + Eperisone Hydrochloride (SR) 150mg Capsules | For the treatment of patient with acute musculoskeletal spasm associated with low back pain. | 24-8-2011 |
| 58. | Rosuvastatin Calcium eq. to Rosuvastatin 20mg/20mg + Fenofibrate BP 67.5mg/145mg Tablets | Additional Strength | 24-8-2011 |
| 59. | Eletriptan Hydrobromide Tablets 20mg/40mg | For the treatments of Migraine with or without aura in adults | 25-8-2011 |
| 60. | Acyclovir 50g + Hydrocortisone 10mg Cream | For the early treatment of recurrent cold sores in adult patients to reduce likelihood of ulcerative cold sores and to shorten the lesion healing time | 30-8-2011 |
| 61. | Clindamycin Phosphate eq. to Clindamycin 1% w/w + Hydrous Benzoyl Peroxide eq. to Anhydrous Benzoyl peroxide 2.5% w/w topical gel | For topical treatment of inflammatory acne vulgaris (Additional lower strength) | 30-8-2011 |
| 62. | Polyethylene glycol 0.4% w/v + Propylene Glycol 0.3% w/v Eye Drops | For temporary relief of burning and irritation due to dryness of the eye | 13-9-2011 |
| 63. | Glutathione for Injection 600mg | For the treatment of alcoholic liver diseases like alcoholic fatty liver, alcoholic liver fibrosis, alcoholic liver cirrhosis, and hepatitis | 14-9-2011 |
| 64. | Udenafil Tablet 100mg/200mg | For the treatment of erectile dysfunction | 15-9-2011 |
| 65. | Metformin (ER) 1000mg/ 500mg +Saxagliptin 2.5 mg/5mg Tablets | As an Adjunct to diet and exercise to improve glycemic control in adults patients with type 2 diabetes mellitus when treatment with both Saxagliptin and Metformin is appropriate | 19-9-2011 |
| 66. | Ceftriaxone Sodium eq. to Ceftriaxone Sodium 1000mg/2000mg + Sulbactam Sodium eq. Sulbactam 500mg/1000mg.injection | Additional Indication - For the treatment of urinary tract infections and lower respiratory tract infections, various bacteria | 21-9-2011 |
| 67. | Difluprednate 0.05% w/v + Moxifloxacin 0.5% w/v Eye Drops | For steroid responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists | 22-9-2011 |
| | • | , | |

| 60 | S() Amindining 2 5mg/5mg (Classide and 75 175 | For the treatment of high right house to be a set of the | 22.0.2011 |
|-----|--|---|------------|
| 68. | S(-) Amlodipine 2.5mg/5mg + Clopidogrel 75mg/75mg Tablets | For the treatment of high risk hypertensives who need anti- platelet therapy especially for prophylaxis of artherothrombotic events in patients with a history of recent stroke or established peripheral arterial disease | 22-9-2011 |
| 69. | Acrivastine Capsules 8mg | For the symptomatic relief of allergic rhinitis. | 26-9-2011 |
| 70. | Tacalcitol ointment 4mcg | For the treatment of psoriasis vulgaris in adults only. | 26-9-2011 |
| 71. | Each Combipack contains: | For the treatment of Osteoporosis in post menopausal women | 30-9-2011 |
| | (A) One tablet of Risedronate Sodium - Each film coated tablet contains: | | |
| | Risedronate Sodium (as Risedronate Sodium Hemi- Pentahydrate) - 35 mg | | |
| | (B) Six Capsules of FDC of Calcitriol +Calcium carbonate + Zinc | | |
| | Each soft gelatin capsule contains:- | | |
| | Calcitriol IP - 0.25mcg | | |
| | Calcium Carbonate IP - 500mg (eq. to elemental calcium 200mg) | | |
| | Zinc (as Zinc Sulphate USP) 7.5 mg | | |
| 72. | Akiskiren Hemifumarate eq. to Aliskiren 150mg/150mg/300mg/300mg +Amlodipine Besilate IP eq. to Amlodipine 5mg/10mg/5mg/10mg Tablets | | 12-10-2011 |
| 73. | Ferrous Ascorbate eq. to elemental iron 100mg + Folic acid IP 1.1mg + Methylcobalamin 1.5mg +Zinc Sulphate monohydrate IP eq to elemental Zinc 22.5mg Tablets. (Additional strength) | For the treatment of Iron deficiency anaemia | 19-10-2011 |
| 74. | Arterolane Maleate 150mg + Piperaquine Phosphate 750mg Tablet | Treatment of acute uncomplicated malaria infection due to Plasmodium falciparum in adults. | 19-10-2011 |
| 75. | Tafluprost Eye Drops 0.0015% w/v | For the reduction of elevated intraocular pressure in open angle glaucoma and ocular hypertension. | 21-10-2011 |
| 76. | Gatifloxacin IP eq. to Gatifloxacin (Anhydrous) 3.0mg + Difluprednate 0.5mg + Benzalkonium Chloride IP (As preservative) 0.01% w/v Eye Drops | For the treatment in the patients with inflammatory occular conditions where bacterial infection or a risk of bacterial occular infection exists | 24-10-2011 |
| 77. | Diazinon Flea Drops | For dogs infected with fleas. | 11-11-2011 |
| | Spot on Dogs 30.0% w/w | | |
| 78. | Diclofenac Sodium IP (as sustained release Pellets) 100mg + Thiocolchicoside (as sustained release pellets) 16mg Capsule. (New Dosage Form) | For the treatment of acute inflammation conditions associated with spasm in adults only. | 15-11-2011 |
| 79. | Epalrestat 50mg + Methylcobalamin 500mcg Tablets | For the treatment of Diabetic Neuropathy | 16-11-2011 |
| 80. | Cabazitaxel Injection 60mg/1.5ml | In combination with prednisone for treatment of patients with hormone-refractory metastatic prostrate cancer previously treated with a docetaxel-containing treatment regimen. | 16-11-2011 |
| 81. | Ketorolac Tromethamine IP 4.00mg + Olopatadine Hcl eq. to Olopatadine 1.00mg + Benzalkonium Chloride IP 0.05mg Ophthalmic solution | - · · | 21-11-2011 |

| 82. | Tramadol Hcl IP (As Immediate Release Layer) 50mg + Diclofenac Sodium IP (As Sustained Release Layer) 75mg Tablets | For the symptomatic treatment of moderate to severe pain in adult patients | 21-11-2011 |
|-----|--|--|------------|
| 83. | Budesonide IP 1mg/0.5mg + Formoterol Fumarate Dihydrate IP eq to Formoterol Fumarate 20mcg/20mcg Respirator suspension. (Additional Strength) | For the treatment of asthma where the use of corticosteroid and long acting beta agonist in combination is appropriate | 21-11-2011 |
| 84. | Amlodipine Besilate IP eq.to Amlodipine 10mg/5mg/5mg/5.5mg + Olmesartan Medoxomil 40mg/40mg/20mg/5mg/5mg + Hydrochlorothiazide IP 25mg/12.5mg/12.5mg/12.5mg/12.5mg/12.5mg | For the treatment of essential hypertension | 23-11-2011 |
| 85. | Each Combipack contains Part (A) One Strip of FDC of Ramipril + Amlodipine Tablets;- Each uncoated tablet contains Ramipril IP - 5mg, Amlodipine besylate IP eq to Amlodipine 5mg B) Each Film coated tablet contains ;- Atorvastatin Calcium IP eq to Atorvastatin 10mg/20mg. | For the treatment of hypertension and as an adjunct to diet in patients to reduce elevated total cholesterol and triglyceride levels hypertension and dyslipidemia. The polypack should be used in patients for whom treatment with all of the three i.e Ramipril, Amlodipine and Atorvastatin is appropriate. | 28-11-2011 |
| 86. | Rosuvastatin Calcium IP eq to Rosuvastatin 5mg + Ezetimibe 10mg. (Additional Strength) | For the treatment of patients with primary hypercholesterolemia. | 5-12-2011 |
| 87. | Tolperisone Hcl IP 150mg + Diclofenac IP 50mg Tablets | For the treatment of Patients with acute muscle/musculoskeletal spasm in adult | 7-12-2011 |
| 88. | Paracetamol IP 325mg + Pamabrom USP 25mg Tablets | For the treatment of patient with pre-menstrual syndrome in adults only | 7-12-2011 |
| 89. | Dosulepine Hcl 50mg/75mg + Methylcobalamin 1500mcg/1500mcg Tablets | For the treatment of neuropathic pain. | 8-12-2011 |
| 90. | Trifluridine Eye Drops 1% w/v | For the treatment of primary keratoconjunctivitis and recurrent epithelial keratitis due to herpes simplex viruses type 1 and 2. | 8-12-2011 |
| 91. | Dabigatran Etexilate(as Mesilate) hard gelatin Capsules 75mg/110mg/150mg | For prevention of stroke, systemic embolism and reduction of vascular mortality in adult patients with atrial fibrillation. | 12-12-2011 |
| 92. | Montelukast Sodium IP eq.to Montelukast 10mg +Olopatadine Hcl 5mg Tablets | For the treatment of relief of symptoms of allergic rhinitis | 16-12-2011 |
| 93. | Fosaprepitant (as Dimeglumine) for Injection 150mg. | Prevention of acute and delayed nausea and vomiting associated with highly emetogenic cisplatin based cancer chemotherapy in adults. Prevention of nausea and vomiting associated with | 16-12-2011 |
| | | moderately emetogenic cancer chemotherapy in adults. | |
| 94. | Abiraterone Acetate Tablets 250mg | In combination with prednisone for the treatment of patients with metastatic castration-resistant prostrate cancer who have received prior chemotherapy containing docetaxel. | 16-12-2011 |
| 95. | Crizotinib hard gelatin Capsules 200mg/250mg | For the treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) - positive as detected by an FDA-approved test. | 16-12-2011 |
| 96. | Amlodipine Besilate IP eq. to Amlodipine 5mg + Perindopril Erbumine BP 8mg Tablets. (Additional Strength). | For the treatment of mild to moderate hypertension in adults only and or treatment of stable coronary artery disease in patients already controlled with Perindopril & Amlodipine given concurrently at same dose level | 20-12-2011 |
| 97. | Iron Amino Acid Chelate as Ferrous Bisglycinate (75mg) eq. to Elemental Iron 15mg + Folic acid IP 0.25mg + Vitamin B 12 IP 3.75mcg +Zinc Picolinate (26mg) eq. to Elemental Zinc 5.5mg Syrup | For the treatment of Iron Deficiency Anaemia | 22-12-2011 |

| Sl. No | Name of Drug | Indication | Date of issue |
|--------|---|---|---------------|
| 1. | Ornidazole IP 1 % w/w + Povidone Iodine IP (0.5% w/w available Iodine) 5% w/w Ointment | For the prevention and treatment of surgical wound infections in adult patients undergoing surgery | 3-1-2012 |
| 2. | Aceclofenac IP (Sustained released pellets) 200mg/200mg + Thiocolchicoside IP (Sustained released pellets) 8mg/16mg Capsules. (New Dosage Form) | For the treatment of acute inflammation conditions associated with spasm in adults only | 4-1-2012 |
| 3. | Budesonide IP 1mg/0.5mg + Formoterol Fumarate Dihydrate IP eq to Formoterol Fumarate 20mcg/20mcg Respirator suspension. (Additional Indication) | In the regular treatment of patients with moderate to severe Chronic Obstructive Pulmonary Disease (COPD) with frequent symptoms and a history of exacerbations | 4-1-2012 |
| 4. | Diclofenac Sodium IP (Enteric coated granules) 50mg + Eperisone Hcl IP (Tablet) 50mg Hard Gelatin Capsules. (New Dosage Form) | For the treatment of Patients with acute musculoskeletal spasm associated with low back pain | 6-1-2012 |
| 5. | Degarelix for Injection 80mg/120mg per vial | For the treatment of adult patients with advanced hormone dependent prostate cancer | 19-1-2012 |
| 6. | Linagliptin film coated Tablet 5mg | As an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus Important limitations of use: 1. Should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis 2. Has not been studied in combination with insulin | 23-1-2012 |
| 7. | Eribulin mesylate solution for injection 0.88mg / 2ml vial | For the treatment of patients with locally advanced or metastatic breast cancer who have progressed after at least two chemotherapeutic regimens for advanced disease. Prior therapy should have included an anthracycline and a taxane unless patients were not suitable for these treatments | 27-1-2012 |
| 8. | Plerixafor Injection 20mg/ml | In combination with Granulocyte Colony Stimulating Factor (G-CSF) to enhance mobilisation of hematopoietic stem cells to the peripheral blood for collection and subsequent autologous hematopoeitic stem cells transplantation in patients with non-Hodgkin's lymphoma and multiple myeloma whose cells mobilise poorly | 31-1-2012 |
| 9. | Cephalexin IP (as monohydrate) 200mg + Kanamycin IP (as monosulphate) 100,000 IU Suspension per 10 gm Intramammary Suspension | | 7-3-2012 |
| 10. | Mucotrol Concentrated Oral Gel Wafer | For the management of pain and relief of pain, by adhering to the mucosal surface of the mouth, soothing oral lesions of various etiologies, including oral mucositis/stomatitis (may be caused by chemotherapy or radiotherapy). Irritation due to oral surgery and traumatic ulcers caused by braces or ill fitting dentures of diseases and for diffuse apthous ulcers | 7-3-2012 |
| 11. | Ulinastatin for Injection 50000 IU/100000 IU | For the treatment of severe sepsis | 3-4-2012 |
| 12. | Sterile Solution for Peritoneal Dialysis Each 1000 ml contains: Calcium chloride dihydrate 0.184 g/l + Glycine 0.51 g/l + L- Alanine 0.951 g/l + L- Arginine 1.071 g/l + L- Isoleucine 0.85 g/l + L- Histidine 0.714 g/l + L- Leucine 1.02 g/l + L- Lysine (as Hcl) 0.955 g/l + L-Methionine 0.85 g/l + L- Phenylalanine 0.57 g/l + L-Proline 0.595 g/l + L- Serine 0.51 g/l + L- Threonine 0.646 g/l + L-Tryptophan 0.27 g/l + L-Tyrosine 0.3 g/l + L-Valine 1.393 g/l + Magnesium chloride hexahydrate | As Nutritional supplements for malnourished renal failure patients (Albumin concentration lower than 35 g/litre) being maintained on peritoneal dialysis | 18-4-2012 |

| | 0.051 g/l + Sodium chloride 5.38 g/l + Sodium lactate 4.48 g/l | | |
|-----|---|--|-----------|
| 13. | Blonanserin Tablet 2mg/4mg/8mg | For the treatment of Schizophrenia in adult patients when first line treatment is ineffective | 24-4-2012 |
| 14. | Etravirine Tablet 100mg | In combination with other antiretroviral agents for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in antiretroviral treatment-experienced adult patients who have evidence of viral replication and HIV-1 strains resistant to an NNRTI and other antiretroviral agents. It should be used only in situations where other treatments have failed or cannot be used because of side effects | 26-4-2012 |
| 15. | Ticagrelor Tablet 90mg | For the prevention of thrombiotic events (cardiovascular death, myocardial infarction and stroke) in patients with Acute coronary syndromes (ACS) unstable angina, non ST elevation Myocardial infarction (STEMI) including patients managed medically, and those who are managed with percutaneous those who are managed with percutaneous coronary intervention (PCI) or coronary artery by-pass grafting (CABG) | 3-5-2012 |
| 16. | Clenbuterol Hydrochloride Solution for injection 30mcg/ml | For the treatment of equine respiratory diseases in horses where airway obstruction due to bronchospasm and/or accumulation of mucus is contributing factor and improved muco-ciliary clearance is desirable. The product should be used alone or as an adjuvant therapy. The product is of particular value in cases of: - Acute, sub-acute and chronic infections where the presence of mucus and/or microorganisms may stimulate bronchospam e.g. Bronchitis, broncholitis and Bronchopneumonia | 1-7-2012 |
| | | — Acute, sub-acute and chronic respiratory allergies. | |
| 17. | Apixaban Tablets 2.5mg | — Chronic obstructive pulmonary disease (COPD) Prevention of venous thromboembolic events (VTE) in adult patients who have undergone elective hip or knee replacement surgery | 3-8-2012 |
| 18. | Ethyl esters of iodized fatty acids of poppy seed oil solution for injection 48%. | In Diagnostic Radiology: Lymphography and Diagnosis of liver lesions. (Diagnosis of the spread of malignant lesions, whether hepatic or not, by selective hepatic arterial injection). In Interventional Radiology: Embolization with surgical glues. (In association with surgical glues during vascular embolizations). | 22-8-2012 |
| | | In Endocrinology: Prevention of iodine deficiency disorders. (This treatment should only be used when other methods of supplementation, particularly iodization of salt and/or drinking water, cannot be undertaken) | |
| 19. | Conivaptan Hydrochloride Injection 20mg/4ml | For the treatment of euvolemic hyponatremia (e.g. the syndrome of inappropriate secretion of antidiuretic hormone, or in the setting of hypothyroidism, adrenal insufficiency, pulmonary disorders, etc.) in hospitalized patients | 21-8-2012 |

| 20. | Colesevelam Hydrochloride Tablets 625mg & Sachets for oral suspension 3.75gm/1.875 gm | Indicated as an adjunct to diet and exercise to reduce elevated low-density lipoprotein cholesterol (LDL-C) in adults with primary hyperlipidemia as monotherapy or in combination with an hydroxymethyl-glutaryl-coenzyme A (HMG CoA) reductase inhibitor (statin). Indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus | 29-8-2012 |
|-----|---|--|------------|
| 21. | Tolvaptan Tablets 15/30mg | For the treatment of clinically significant hypervolemic and euvolemic hyponatremia [serum sodium < 125mEq/L or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction], including patients with heart failure, cirhossis, and syndrome of inappropriate antidiuretic hormone (SIADH) | 6-9-2012 |
| 22. | Agomelatine Tablets 25mg | For the treatment of major depression in adult patients aged 18 –65 years only with normal liver functions | 10-9-2012 |
| 23. | Almotriptan Tablet 6.25mg/ 12.5mg | For acute treatment of migraine attacks with a history of migraine with or without aura in adults | 5-10-2012 |
| 24. | Choline Fenofibrate DR Capsule 45mg/135mg & DR Tablet 45mg/135mg. | For treatment of mixed lipidemia in combination with statins, severe hypertriglyceridemia | 1-11-2012 |
| 25. | Trimethoxysilyl Quaternary Ammonium Chloride solution for spray inanimate surfaces for disinfection 0.71% | Surface protectant and disinfectant for persistent disinfection of all inanimate surfaces. Effective up to 14 days | 16-11-2012 |
| 26. | Imipenem IP (Sterile) eq to anhydrous Imipenem 1 g + Cilastatin Sodium IP (sterile) eq to Cilastatin 1 g Powder for Injection (Additional strength) | | 28-6-2012 |
| 27. | Diclofenac Diethylamine BP 4.64 % w/v Eq. to Diclofenac Sodium IP 4.00 % + Absolute Alcohol IP 10.00 % v/v Topical Solution | | 21-12-2012 |

| Sl. No. | Name of Drug | Indication | Date of issue |
|---------|--|--|---------------|
| 1. | Diclofenac Diethylamine BP 2.32 % w/v Eq to Diclofenac Sodium 2.00 % w/v + Methyl Salicylate IP 10.00% w/v + Menthol 5.00 % w/v + Absolute Alcohol IP 10.00 % v/v Topical Solution (Additional Strength) | For the Treatment of Acute Lower Back Pain | 1-1-2013 |
| 2. | Telmisartan 20mg + Amlodipine Besylate Eq. to Amlodipine 2.5mg + Hydrochlorothizide 6.25mg Tablets (Additional Strength) | For the Treatment of Essential Hypertension | 3-1-2013 |
| 3. | Amlodipine besilate IP Eq. to Amlodipine 10mg + Indapamide SR 1.5 mg Tablet (Additional Strength) | For the treatment of mild to moderate Hypertension | 2-1-2013 |
| 4. | Nicoumalone Tablets 0.5mg/5mg/6mg | For the prevention and treatment of thromboembolic diseases such as venous thrombosis, atrial fibrillation with embolization, pulmonary embolism and as an adjunct in the treatment of coronary occulsion. | 18-1-2013 |
| 5. | Ruxolitinib Tablet 5mg/15mg/20mg | For the treatment of patients with myelofibrosis including primary myelofibrosis, post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis. | 28-1-2013 |
| 6. | Dabigatran Etexilate Mesilateeq to Dabigatran Etexilate 75mg/110mg/150mg | For the prevention of venous thromboembolic events in patients who have undergone orthopaedic surgery | 11-2-2013 |

| 7. | Dexamethasone Intravitreal Implant 0.7mg (in solid polymer drug delivery system) | For the treatment of non-infectious uveitis affecting the posterior segment of the eye. However it should not be used when there is gap in posterior capsule. | 20-2-2013 |
|-----|---|---|-----------|
| 8. | Cefixime Tablets 200mg & Cefixime Oral Suspension 100mg/5ml | For the treatment of enteric (typhpoid) fever. | 22-2-2013 |
| 9. | Diclofenac Colestyramine 145.6mg Capsule Corresponding to Diclofenac Sodium 75mg (Additional Indication) | For the treatment of post-traumatic or post-operative swelling and inflammation | 25-2-2013 |
| 10. | Saroglitazar 2mg/4mg Tablets | For the treatment of diabetic dyslipidemia and hypertriglyceridemia with type-2 diabetes mellitus not controlled by statin therapy | 25-2-2013 |
| 11. | Tadalafil Tablet 2.5mg/5mg | For the treatment of erectile dysfunction. For the treatment of erectile dysfunction with benign prostatic hyperplasia. | 27-2-2013 |
| 12. | Azithromycin Tablet 250mg/500mg | For the treatment in uncomplicated multi drug resistant enteric fever only | 6-3-2013 |
| 13. | Betahistine Dihydrochloride SR Tablets 24mg/32mg/48mg | For the symptomatic treatment of peripheral vertigo | 14-3-2013 |
| 14. | Palonosetron Melt in Mouth Tablets 0.5mg | For prevention of chemotherapy induced nausea and vomiting | 19-3-2013 |
| 15. | Iobitridol Solution for Injection Each 100mg solution contain 65.81g Iobitridolcorressponding mass of Iodine 30gm | By intracavity route for Arthrography and Hysterosalpingography. | 19-3-2013 |
| 16. | Everolimus 2.5mg/5mg/10mg Tablets | For the treatment of tuberous sclerosis complex (TSC) who have renal angiomyolipoma not requiring immediate surgery. For the treatment of postmenopausal women with hormone receptor positive advanced breast cancer in combination with an aromatase inhibitor. | 21-3-2013 |
| 17. | Each sustained release tablet contains: Thiocolchicoside IP 8/16mg + Lornoxicam 8/16mg (Additional Strength) | For the treatment of patients with acute painful musculoskeletal conditions | 8-4-2013 |
| 18. | Combination of Lactic Acid Bacteria (VSL#3) Sachet | For mild to moderately active ulcerative colitis as an adjunctive therapy subject to condition that it should not be used in patients who are on steroids or immunocompromised. | 8-4-2013 |
| 19. | Tolperisone HCl 450mg + Diclofenac Sodium IP100mg SR tablets (Additional Strength) | For the treatment of Patients with acute muscle/musculo skeletal spasm in adult | 12-4-2013 |
| 20. | Telmisartan IP 80mg + Hydrochlorothiazide IP 25mg uncoated bilayered tablet (Additional Strength) | For the treatment of hypertension as second line therapy | 14-5-2013 |
| 21. | Carboxy Methyl Cellulose Sodium Lubricant Eye Drops 0.5% w/v (Unit dose pack without preservative) | For temporary relief from burning irritation and discomfort due to dryness of the eye or due to exposure to wind or sun. May also be used as a protectant against further irritation. | 27-5-2013 |
| 22. | Garenoxacin Mesylate tablets 200mg | For the treatment of bacterial respiratory tract infection | 7-6-2013 |
| 23. | Trimethoxysilyl Quaternary Ammonium Chloride 0.40% hand protectant and sanitizer | Hand Protectant and Sanitizer for external use. Effective up to 24 hours. | 11-6-2013 |
| 24. | Sildenafil Citrate Orally Disintegrating Strips. 25mg/50mg (Additional Dosage Form) | For the treatment of erectile dysfunction. | 5-7-2013 |
| 25. | Dextromethorphan Hydrobromide Extended Release suspension 30mg/5ml (Additional Dosage Form) | For the relief of persistent dry irritating cough in adults patients only. | 1-8-2013. |
| | | | |

| 26. | Iron (III) isomaltoside 1000 solution for injection/infusion 100mg per ml | For the treatment of iron deficiency anaemia when oral iron preparations are ineffective or cannot be used and when there is a clinical need to deliver iron rapidly. | 20-8-2013 |
|-----|---|--|------------|
| 27. | Tramadol HCL 200mg Controlled Release tablet (Additional Dosage Form) | For use in in-patients under hospital settings for severe acute pain for a period not exceeding 5 days. | 12-9-2013 |
| 28. | FDC of Aeterolane Malate (150mg) + Piperaquine Phosphate (750mg) Tablet | Treatment of acute uncomplicated malaria infection due to Plasmodium vivax in adults | 11-10-2013 |
| 29. | Micronised Purified Flavonoid Fraction of Rutacea 1000 mg (MPFF) | Acute hemorrhoid piles. | 28-11-2013 |
| | Each film coated tablet contains "Diosmin 900mg and Flavonoids expressed as Hesperidine 100mg of Micronised Purified Flavonoid Fraction of Rutaceae 1000mg (MPFF) | | |
| | (Additional strength) | | |
| 30. | Artesunate injection. Each combi pack contains: | Promoted for use in severe malaria including cerebral | 13-11-2013 |
| | "Artesunate for injection 120mg | malaria and as a second line in chloroquine resistant malaria cases only. | |
| | (Each vial contains Artesunate IP 120mg), Sodium bicarbonate injection IP 5%w/v | cases only. | |
| | (Each 2ml ampoule contains: | | |
| | Sodium Bicarbonate IP 5% w/v), Sodium chloride Injection IP 0.9% w/v | | |
| | (Each 10ml ampoule contains: | | |
| | Sodium chloride IP 0.9% w/v)" | | |
| | (Additional strength) | | |
| 31. | Tapentadol Extended release Tablet 50mg (Additional Strength) | For use in in-patients for dose titration under hospital settings for severe acute pain for a period not exceeding 5 days. | 12-11-2013 |
| 32. | Divalproex Sodium Oral Solution 500mg/5ml (Additional Dosage Form) | For monotherapy and adjunctive therapy in the treatment of patients with complex partial seizures that occur either in isolation or in association with types of seizures for adult patients only. | 23-12-2013 |
| 33. | Diclofenac Sodium Sustained Release Tablet (Two layered tablets) 150mg. | For symptomatic treatment of pain and inflammation in irritation in association with degenerative joint | 20-12-2013 |
| | Each tablet contains Diclofenac Sodium IP-125 mg Slow Release, Diclofenac Sodium IP-25 mg Immediate Release(Additional Dosage Form) | disease/osteoarthritis. | |
| 34. | Ivabradine HCL Tablets 5mg/7.5mg | Additional indication of "treatment of chronic heart failure in | 19-12-2013 |
| | (Additional indication) | chronic heart failure NYHA II to IV class with systolic dysfunction, in patients in sinus rhythm and whose heart rate is > 75 bpm, in combination with standard therapy including | |
| | | beta-blocker therapy or when beta-blocker therapy is contraindicated or not tolerated". | |
| 35. | Risperidone prolonged release suspension for injection (Deep Intramuscular Deltoid route using appropriate safety needle) | Same as already approved | 19-12-2013 |
| | (Additional route of administration) | | |

| Sl. No | Drug | Indication | Date |
|--------|--|---|------------|
| 1. | Tolterodine Tartrate Extended release tablets 2mg/4mg (Additional Dosage Form) | For the treatment of overactive bladder with symptoms of urge urinary incontinence, urinary incontinence, urgency and frequency. | 31-1-2014 |
| 2. | Bortezomib for injection 3.5mg (Sub-cutaneous route of administration as an alternate to intravenous route) (Additional route of administration) | Same as already approved | 28-1-2014 |
| 3. | Paclitaxel Injection Concentrate for Nanodispersion 100mg and 300mg (Additional Dosage Form) | For the treatment of Breast cancer after failure of combination therapy metastatic disease or relapse within six months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated. | 22-1-2014 |
| 4. | Heparin Sodium Topical Solution 1000 IU/ml (Additional Dosage Form) | For the management of post infusion superficial thrombophlebitis. | 16-1-2014 |
| 5. | Simethicone Orally disintegrating strip 62.5mg | For the treatment of flatulance and as adjuvant in hyperacidity. | 14-11-2010 |
| 6. | Eltrombopag Olamine tablet 25/50mg (Additional Indication) | Indicated in patients with chronic hepatitis C Virus (HCV) infection for the treatment of thrombocytopenia to: Enable the initiation of interferon based therapy, optimise interferon based therapy. | 7-4-2014 |
| 7. | Decitabine lyophilised powder for injection 50mg/20ml vial (additional indication) | For the treatment of adult patients aged 65 years and above with newly diagnosed de novo or secondary acute myeloid leukemia (AML), according to World Health Organisation (WHO) classification, who are not candidates for standard induction chemotherapy. | 9-4-2014 |
| 8. | Glycopyrronium bromide inhalation powder hard capsule with inhaler. Each capsule contains 63mcg Glycopyrronium bromide eq. to Glycopyrronium 50 mcg. | Indicated as a once daily maintenance bronchodilator treatment to relieve symptoms of patients with Chronic Obstructive Pulmonary Disease (COPD). | 1-4-2014 |
| 9. | Micafungin sodium for injection 100mg/vial (Additional Strength) | Treatment of patient with candidemia, acute Disseminated Candidiasis, Candida Peritonitis and abscesses. Treatment of patients with Esophageal Candidiasis, Prohylaxis of Candidia and Aspergillus Infections in patients undergoing Hemopoeitic stem cell transplantation. Treatment of patient with Fungemia, Respiratory mycosis, Gastrointestinal mycosis caused by Aspergillus sp | 2-5-2014 |
| 10. | Apixaban tablets 2.5mg/5mg (Additional Strength & Additional Indication) | Prevention of stroke and systemic embolism in adult patients with non-valvular artrial fibrillation (NVAF), including those with one or more risk factors, such as prior stroke or transient ischemic attack (TIA); Age≥75 years; hypertension; diabetes mellitus; symptomatic heart failure (NYHA Class≥II). Compared to warfarin apixaban also results in less bleeding, including intracranial hemorrhage. | 16-5-2014 |
| 11. | Mometasone furoate nasal spray 50 mcg/actuation (Additional Indication) | For the treatment of nasal symptoms of seasonal allergic and perennial allergic rhinitis, in adults and pediatric patients 2 years of age and older. | 21-5-2014 |
| 12. | Micronised Purified Flavonoid 1000mg (MPFF). "Each film coated tablet contains diosmin 900mg and flavonoids expressed as Hesperidine 100 mg of Micronised Purified Flavonoid fraction of Rutaceae 1000mg (MPFF)" | Acute hemorrhoid piles | 28-11-13 |

| 13. | Gemcitabine HCl injection ready to use infusion bags Gemcitabine HCL equivalent to Gemcitabine 10mg per mL(100,120,130,140,150,160,170,180,190 and 200mL) (Approved as infusion bags) | Ovarian cancer-Gemcitabine injection in combination with carboplatin is indicated for the treatment of patients with advanced ovarian cancer that has replaced at least 6 months after completion of platinum based therapy. Breast Cancer-Gemcitabine injection in combination with Paclitaxel is indicated for the first line treatment of patients with meta static breast cancer after failure of prior anthracycline containing adjuvant chemotherapy, unless anthracyclinees were clinically contraindicated. Non-small cell lung concer-Gemcitabine injection is indicated in combination with cisplatin for the first line treatment of patients with inoperable, locally advanced (Stage IIIA or IIIB), or meta static (stage IV) non- | 17-6-2014 |
|-----|--|---|------------|
| 14. | Tadalafil Orally Disintegrating Strip 10mg & 20mg (Additional dosage form) | For erectile dysfunction | 30-7-2014 |
| 15. | Hydroxychloroquine Sulphate USP 400 mg tablets (Additional Indication) | As an adjunct to diet and exercise to improve glycemic control of patients on metformine, sulfonylurea combination in patients with Type II Diabetes. | 28-7-2014 |
| 16. | Nevirapine Extented release tablet 400mg (Additional Strength) | For use in combination with other anti-retroviral agents for the treatment of HIV-1 infection in adults | 1-7-2014 |
| 17. | Artesunate injection. Each combipack contains: "Artesunate for injection 120mg (Each vial contains Artesunate IP 120mg), Sodium bicarbonate injection IP 5%w/v (Each 2ml ampoule contains: Sodium Bicarbonate IP 5% w/v), Sodium chloride Injection IP 0.9% w/v (Each 10ml ampoule contains: | Promoted for use in severe malaria including cerebral malaria as a second line in chloroquine resistant malaria cases only. | 13-11-2013 |
| | Sodium chloride IP 0.9% w/v)" | | |
| 18. | Cerebrolysin solution for injection. Each ml contains: Porcine brain derived peptide preparation (Cerebrolysine concentrate) 215.2mg | For amelioration of cranial injury, cerebrovascular pathological sequelae and aprosexia in dementia. | 25-7-2014 |
| 19. | Artesunate powder for injection 60mg/vial along with 6ml ampoule of phosphate buffer solution (pH 8.0;0.30M) | For the treatment of severe Falciparum malaria in areas where there is evidence if quinine resistance | 2-7-2014 |
| 20. | Montelukast sodium chewable tablet 4mg/5mg and Montelukast sodium film coated tablet 10mg. | Asaddon therapy in mild to moderate asthma in adequately controlled by inhaled cortico steroids and short active B2 agonist, Exercise induced broncho constriction. | 26-2-2002 |
| 21. | Lactobacillus brevis CD2 Lozenges 100mg (corresponding to not less than 1 billion) of live, lyophilised, lactic acid bacteria, Lactobacillus brevis CD2. (Additional indication) | Prevention of radiotherapy and chemotherapy induced oral mucositis in cancer patients. | 9-8-2014 |
| 22. | Rivaroxaban tablet 15/20mg (Additional strength/indication) | Treatment of deep vein thrombosis and for prevention of recurrent DVT and pulmonary embolism. For the prevention of stroke and systemic embolism in patient with non-valvular arterial fibrillation. | 2-9-14 |
| 23. | Hydroxychloroquine Sulphate Tablet 300mg (Additional strength/indication) | Indicated for the treatment of patients with lower body weight i.e 45 to 60kg in Rhemuatoid arthritis, Systemic Lupus Erythematosus & Polymorphic Light Eruption | 9-8-2014 |
| 24. | Ginkgo biloba extract Tablet 120mg (Additional strength) | For the treatment of dementia, vertigo and tinnitus in adult patients. | 9-12-2014 |
| 25. | Bendamustine Hydrochloride lyophilised injection 25mg (Additional strength) | Treatment of patients with chronic lymphocytic leukemia | 19-9-2014 |
| 26. | Metformin ER tablet 750mg (Additional strength) | As an adjunct to diet and exercise to improve glycemic control in adult patients with Type II Diabetes. | 25-9-14 |

| 27. | Sorafenib tosylate tablet 200mg (Additional indication) | For the treatment of patients with locally advanced or meta static differentiated thyroid carcinoma refractory to radioactive iodine. | 25-9-14 |
|-----|--|--|------------|
| 28. | Deferasirox Dispersible tablet 100/400mg (Additional indication) | Treatment of chronic iron overload in patients with non-transfusion dependent thalassemia (NTDT) syndromes aged 10 years and older. | 26-9-2014 |
| 29. | IMATINIB mesilate 100/400mg tablets and 100 mg capsules (Additional indication) | Treatment of Paediatric patients with newly diagnosed Philadelphia chromosome positive acute Lymphoblastic Leukaemia (Ph+ALL) integrated with chemotherapy | 9-9-2014 |
| 30. | Tapentadol Extended release tablet 100/150/200mg (Additional strength) | For use in in-patients under hospital setting for severe acute pain for a period not exceeding 5 days | 9-12-13 |
| 31. | Olanzapine pamoate Prolonged release powder for suspension for I.M injection (combipack) Each combipack contains a. Vial of Olanzapine pamoate Prolonged release powder for suspension. Each vial contains: Olanzapine pamoate monohydrate eq to Olanzapine | For the treatement of schizophrenia | 14-10-14 |
| | 210mg/300mg/405mg. After reconstitution each ml of suspension contains 150mg. b. Vial of vehicle: 3ml clear, colourless to slight yellow solution contains: | | |
| | Manitol IP 5.070%w/vSodium carboxymethyl cellulose IP 0.770% w/v Polysorbate 80 IP 0.025%w/vHCL IPq.s. NAOH IPq.s Water for injection q.s. to 100% | | |
| 32. | Rivastigmine transdermal patch. Each Transtermal patch of 15cm2 contains Rivastigmine 27mg (Additional indication) | For the treatment of patients with severe dementia of the Alzheimer's type | 28-10-2014 |
| 33. | Dabigatran Etexilate Mesilate hard gelatin capsule 75/110/150mg (Additional Indication) | Treatment of acute deep vein thrombosis (DVT) and /or pulmonary embolism (PE) and prevention of related death. Prevention of recurrent deep vein thrombosis (DVT) and | 17-11-14 |
| 34. | Pegasperagase 3,750 IU/5 mL | /or pulmonary embolism (PE) and related death Indicated as a component of multi agent chemotherapeutic regimen for the treatment of patients with acute lymphoblastic leukemia who are hypersensitive to asparaginase | 7-3-2014 |
| 35. | Tulobuterol Transdermal Patch 0.5 mg/ 1.0 mg/2.0 mg | For treatment of patients with Asthma and COPD without co-morbidity | 24-4-14 |
| 36. | Azacitidine 100 mg/vial | For the treatment of adult patients with all subtypes of Myelodysplastic Syndrome | 29-4-14 |
| 37. | Azacitidine | For the treatment of adult patients with all subtypes of Myelodysplastic Syndrome | 23-7-14 |
| 38. | Regorafenib film coated tablet 40 mg | For the "treatment of patients with metastatic colorectal cancer (CRC) who have been previously treated with fluoropyrimidine, oxaliplatin and irinotecan-base chemotherapy, ananti-VEGF therapy, and, if KRAS wild type, an anti-EGFR therapy". | 1-7-14 |
| 39. | Roflumilast Tab 500 mcg | For the treatment of severe chronic obstructive pulmonary disease (COPD) (FEV1 post-bronchodilator less than 50% predicted) associated with chronic bronchitis in adult patients with a history of frequent exacerbations as add on to bronchodilator treatment. | 17-7-14 |
| 40. | Dimethyl Fumarate delayed release 120 mg/240 mg Tablet | For "Relapsing remitting multiple sclerosis" | 27-8-14 |

| 41. | Alogliptin Tablet 6.25 mg/12.5 mg/ 25 mg | "To improve glycemic control in combination with other glucose lowering medicinial products including insulin when these together with diet and exercise, do not provide adequated glycemic control or as monotherapy as adjunct to diet and exercise to improve glycemic control in adults aged 18 years or older with type 2 Diabetic mellitus." | 27-8-14 |
|-----|--|--|-----------|
| 42. | Axitinib Tablet 1mg/5mg | For the "treatment of advanced renal cell carcinoma after failure of one prior systemic therapy" | 18-9-14 |
| 43. | Ulinastatin Injection | For the treatment of mild to severe acute pancreatitis. | 19-9-14 |
| 44. | Canagliflozin Tablet 100/300mg | "Indicated as an adjunct to diet and exercise to improve glycemic control in adults with Type-2 Diabetic Mellitus". | 17-11-14 |
| 45. | Levosulpiride 75/150/200 mg (add. Strength) | 75 mg tablet: For the treatment of gastro intestinal problem like fanatical dyspepsia, nausea, vomiting and diabetic gastroperisis. 150/200mg: For the treatment of depression and schizophrenia in adults. | 9-12-14 |
| 46. | Nitrofurantoin dispersible tablets 50 mg (add. Strength/dosage form) | for the treatment of acute uncomplicated urinary tract infections (acute cystitis) caused by susceptible strains of E coli or staphylococcus saprophyticus | 15-12-14 |
| 47. | Methylcobalamin orally disintegrating strips 1500mcg (add. Dosage form/indication) | for the treatment of peripheral neuropathy and diabetec neuropathy | 17-12-14 |
| 48. | Paclitaxel suspension for injection Conc. For nanodispersion 100/300mg (add. Indication) | management of endometriosis associated pain | 18-12-14 |
| 49. | Lapatanib tablet 250mg (Additional indication) | lapatinib in combination with trastuzumab, indicated for the treatment of patients with hormone receptor — negative meta static breast cancer whose tumours over express HER2/neu (ErbB2) and who have progressed on prior trastuzumab therapy in combination with chemotherapy in the meta static setting. | 22-12-14 |
| 50. | Medroxyprogesterone acetate suspension for injection s.c. Single dose container with 104mg Medroxyprogesterone acetate in 0.65 ml suspension for injection (SAYNA<< PRESS) (Additional route/ strength/indication) | Indicated for the management of endometriosis-associated pain. | 18-12-14 |
| 51. | Empty Hydroxy Propyl Methyl Celulose Capsules | NA | 11-1-2011 |

| Sl. No | Name Of Drug | Indication | Date of Approval |
|--------|---|--|------------------|
| 1 | Levocetirizine ODS 2.5mg/5mg (Additional dosage form) | For allergic rhinitis and chronic urticaria. | 14-1-2015 |
| 2 | Decitabine Injection 30mg/vial (Additional pack size) | 1. For treatment of patients with myelodysplastic syndromes (MDS) including previously treated and untreated, de novo and secondary MDS of all French- American-British subtypes(refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, refractory anemia with excess blasts in transformation, and chronic myelomonocytic leukemia) and intermediate-1, intermediate-2, and high-risk international Prognostic Scoring System groups. | 13-2-2015 |
| | | 2. For the treatment of adult patients aged 65 yrs and above with newly diagnosed de novo or secondary acute myeloid leukemia, according to WHO classification, who are not candidates for standard induction chemotherapy. | |

| 3 | Meloxicam ODT 7.5mg/15mg (Additional dosage form) | Non-steroidal anti-inflammatory. | 23-2-15 |
|----|--|---|-----------|
| 4 | Ruxolitinib Tablet 5mg/15mg/20mg (Additional indication) | Treatment of patients with polycythemia vera who are resistant to or intolerant of hydroxyurea. | 23-2-2015 |
| 5 | Gadopentetic acid dimeglumine salt injection 469mg/1ml (0.5mmol) (Additional indication) | For use in MRI in adult patients to facilitate visualization of lesions with abnormal vascularity in the body (excluding heart). | 23-2-2015 |
| 6 | Pazopanib Tablet 200mg/400mg (additional indication) | For the treatment of patients with advanced Soft Tissue | |
| | | Sarcoma (STS) who have received prior chemotherapy. | 23-2-2015 |
| 7 | Eribulin mesylate solution for injection 0.88mg in 2ml vial (Add. Indication) | For the treatment of patients with locally advanced or metastatic breast cancer who have progressed after at least one chemotherapeutic regimens for advanced disease. Prior therapy should have included an anthracycline and a taxane in either the adjuvant or meta static setting unless patients were not suitable for these treatments. | 18-2-2015 |
| 8 | Abiraterone acetate Tablet 250 mg (Additional Indication) | For the treatment of meta static castration resistant prostate cancer in adult men who are asymptomatic or mildly symptomatic after failure of androgen deprivation therapy in whom chemotherapy is not yet clinically indicated, with prednisone or prednisolone. | 27-3-2015 |
| 9 | Indomethacin Injection 1mg/vial (additional Dosage form) | To close a hemodynamically significant patent ductus arteriosus (PDA) in premature infants weighing between 500g and 1750g when after 48 hrs. usual medical management (e.g. fluid restriction, diuretics, digitalis, respiratory support, etc.) is ineffective. Clear-cut clinical evidence of hemodynamically significant patent ductus arteriosus should be present, such as respratory distress, a continuous murmur, a hyperactive precordium, cardiomegaly and pulmonary plethora on chest X-ray. | 30-4-2015 |
| 10 | Ketorolac ODS 10mg (Additional Dosage form) | For the treatment of short-term management of post- operative pain and musco-skeletal pain. | 29-4-2015 |
| 11 | Tiamulin hydrogen fumarate solution for injection 162 mg/ml (additional Dosage form) | For treatment and prevention of swine dysentery caused by Branchyspira hydodysenteriae. | 17-4-2015 |
| | | The product is not appropriate for use for the prevention of disease at the level of herd treatment but should only be used for prevention of swine dysentery in individual animals with a known history of exposure to diseased animals. | |
| 12 | Montelukast Orally Disintegrating Strips 4mg/5mg/10mg (additional Dosage form) | For the Relief of perennial allergic rhinitis in adults & pediatric patients 6 months of age and older. | 29-5-2015 |
| 13 | Apixaban Tablets 2.5/5mg (additional Indication) | Treatment of deep vein thrombosis (DVT) and pulmonary Embolism (PE), and prevention of Recurrent DVT and PE in adult patients. | 29-5-2015 |
| 14 | Ruxolitinib 10mg tablet (additional Strength) | For the treatment of patients with myelofibrosis including primary myelofibrosis, post-polycythemeia vera myelofibrosis or post-essential thrombocythemia myelofibrosis Treatment of patients with polycythemia vera who are resistant to or intolerant of hydroxyurea. | 29-5-2015 |
| 15 | Lorazepam Orodispersible Tablet 1mg/2mg (additional Dosage form) | For the management of anxiety disorders or for the short term relief of symptoms of anxiety or anxiety associated with depressive symptoms. | 29-5-2015 |

| 16 | Brimonidine topical gel 0.33% (Add. Dosage form/indication) | For the topical treatment of persistent (non-transient) facial erythema of rosacea in adults 18 years of age or older. | 18-6-2015 |
|----|---|---|--|
| 17 | Rivaroxaban tablet 2.5mg (add. strength) | Rivaroxaban 2.5mg tablet, co-administered with Acetylsalicylic acid (ASA) alone or with ASA plus Clopidogrel or Ticlopidine, is indicated for the prevention of atherothrombotic events in adult patients after an acute coronary syndrome (ACS) with elevated cardiac biomarkers. | 13-7-2015 |
| 18 | Lamotrigine ER Tablet 25mg/50mg/100mg/200mg | Add-on therapy for partial and secondary generalised tonic-clonic seizures. | 6-8-2012 |
| 19 | Feracrylum 3% solution | For the management of post-operative wounds cuts, Abrasions as a Haemostatic Agent cum anti-septic. | 13-2-2002 |
| 20 | Feracrylum 3% gel | For the management of post-operative wounds cuts, Abrasions as an Haemostatic Agent cum anti-septic. | 13-2-2002 |
| 21 | Arteether injection 150mg/mL | Indicated for the treatment of severe malaria including cerebral malaria and as a second line in chloroquine resistant malaria cases only. | 3-12-2010 |
| 22 | Everolimus dispersible Tablet 0.1mg/0.25mg (additional Indication) | For the prophylaxis of organ rejection in patients receiving a hepatic transplant. In liver transplantation, Everolimus should be used in combination with tacrolimus and corticosteriods. | 21-8-2015 (Condition: Phase-IV trial duly approved by CDSCO shall be conducted in statistically significant number of patients.) |
| 23 | Everolimus Tablet 0.25mg/0.5 mg/0.75mg/1.0mg (additional Indication) | For the prophylaxis of organ rejection in patients receiving a hepatic transplant. In liver transplantation, Everolimus should be used in combination with tacrolimus and corticosteriods. | 21-8-2015 (Condition: Phase-IV trial duly approved by CDSCO shall be conducted in statistically significant number of patients.) |
| 24 | Cetrorelix Acetate injection 0.25mg/0.5ml in 0.5 ml pre- filled syringe. (Additional pack size) | Prevention of premature ovulation in patients undergoing a controlled ovarian stimulation, followed by oocyte pick-up and assisted reproductive techniques. | 26-8-2015 |
| 25 | Exenatide 2 mg powder and solvent for prolonged release suspension for injection in a pre-filled pen (s.c. Injection) (Additional strength) | For treatment of type 2 diabetes mellitus in combination with Metformin Sulphonylurea Thiazolidinedione Metformin and Sulphonylurea Metformin and Thiazolidinedione In Adults who have not achieved adequate glycemic control on maximally tolerated doses of these oral therapies. | 9-9-2015 |
| 26 | Vitamin D3 (Cholecalciferol) Orally disintegrating strips 2000 IU (additional Dosage form) | For the treatment of Vitamin D3 deficiency. | 1-10-2015 |
| 27 | Dexamethasone Intravitreal implant 0.7mg (Additional Indication) | For the treatment of visual impairment due to Diabetic Macular Edema (DME) who are pseudophakic or who are | 12-10-2015 |
| | | | |

| | | considered insufficiently responsive to, or unsuitable for non-corticosteroid therapy. | |
|----|---|--|------------|
| 28 | Zolpidem Tartarate Orally disintegrating Tablet 5mg/10mg (Additional dosage form) | For insomnia. | 12-10-2015 |
| 29 | Etodolac i.m injection 400mg/2ml (Additional dosage form) | For the treatment of post operative orthopaedic pain. | 14-12-2015 |

| Sl. No. | Name of Drug | Indication | Date of issue |
|---------|--|---|---------------|
| 1 | Nintedanib soft Gelatin Capsule 100/150mg | For the treatment of Idiopathic Pulmonary Fibrosis (IPF). | 11-3-2016 |
| 2 | Cisatracurium Besylate Bulk & 2mg/ml injection | As an adjunct to general anaesthesia, to facilitate tracheal intubation and to Provide skeletal muscle relaxation during surgery. | 14-3-2016 |
| 3. | Fomepizole Bulk & 1.5gm/ Ampoule | As an antidote for Ethylene glycol or Methanol used poisoning or for use in suspected Ethylene glycol or Methanol ingestion, either alone or in combination with hemodialysis. | 14-3-2016 |
| 4 | Tofacitinib Tablets 5 mg(Tofacitinib Citrate) | "For the treatment of adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response or intolerance to Methotrexate. It may be used as monotherapy or in combination with methotrexate or other nonbiologic disease-modifying antirheumatic drugs (DMARDs)". | 1-4-2016 |
| 5 | Nintedanib Soft Gelatin Capsule 100/150mg (Nintedanib ethanesulfonate) | "In combination with docetaxel for the treatment of adult patients with locally advanced, metastatic or recurrent non-small cell lung cancer (NSCLC) of adenocarcinoma tumor histology after line chemotherapy". | 1-4-2016 |
| 6 | Ceftaroline Fosamil Injection 600 mg/Vial | "For the treatment of adult (≥ 18 years of age patients with community-acquired pneumonia". | 9-5-2016 |
| 7 | Panobinostat Hard Gelatin Capsules 10mg/15mg/20mg (Panobinostat lactate) | "In combination with bortezomib and dexamethasone, is indicated for the treatment of patients with multiple myeloma, who have received at least 1 prior therapy" | 27-5-2016 |
| 8 | Bepotastine Besilate Bulk & Bepotastine Besilate 1.5 %w/v Ophthalmic solution | "For the treatment of itching associated with allergic conjunctivitis" | 8-6-2016 |
| 9 | Ibutilide Bulk & Description (1.1 mg/ml) Ibutilide Fumarate Injection (1.1 mg/ml) | "For the rapid conversion of atrial fibrillation or atrial flutterof recent onset to sinus rhythm. Patient with atrial arrhythmias of longer duration are less likely to respond to Ibutilide Fumarate Injection. The effectiveness of Ibutilidehas not been determined in patients with arrhythmias of more than 90 days in duration". | 10-6-2016 |
| 10 | Acotiamide Hydrochloride Bulk & Tablet 100mg | "For the treatment of bloating after meals, epigastric bloating and early satiety in functional dyspepsia". | 6-7-2016 |
| 11 | Sacubutril+ Valsartan film coated tablets 50mg/100 mg/200mg | "To reduce the risk of cardiovascular death and hospitalization for heart failure in patients with chronic heart failure (NYHA Class II-IV) and reduced ejection fraction". | 14-7-2016 |
| 12 | Lurasidone Hydrochloride Bulk & Lurasidone Hydrochloride Tablets 40mg/80mg | "For the treatment of Patients with Schizophrenia" | 18.07.16 |
| 13 | Fenticonazole Nitrate Vaginal Capsule 600 mg | "For the treatment of vulvovaginal candidiasis" | 10-8-2016 |

| 14 | Palbociclib Capsules 75mg/100mg/125 mg | "Palbociclib is a kinase inhibitor indicated in combination with Letrozole for the treatment of postmenopausal women with estrogen receptor (ER)-Positive, human epidermal growth factor receptor 2 (HER2)-Negative advanced breast cancer as initial endocrine-based therapy for their metastatic disease". | 11-8-2016 |
|----|---|--|------------|
| 15 | Midodrine Hydrochloride 2.5 mg Tablet | "For the treatment of symptomatic orthostatic hypotension" | 2-9-2016 |
| 16 | Phospholipids Fraction from Bovine Lung (surfactant) 50 mg/vial | "Preventive use in premature neonates with a high risk of respiratory Distress Syndrome" | 12-9-2016 |
| 17 | Dolutegravir 50 mg Tablet & Bulk (Dolutegravir Sodium) | "Indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults weighing more than 40 kg" | 7-11-2016 |
| 18 | Alcaftadine Eye Drops 0.25% w/v & Bulk | "For the prevention of itching associated with allergic conjunctivitis in patients between the age group 10 to 60 years" | 21-11-2016 |
| 19 | Bepotastine Besilate Bulk | | 23-11-2016 |
| 20 | Lenvatinib 4mg/10mg Hard Gelatin Capsules (Lenvatinib Mesylate) | "For the treatment of patients with locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer" | 2-12-2016 |
| 21 | Perampanel 2mg/4mg/6mg/8mg/10mg/12mg Tablets | "The adjunctive treatment of partial-onset seizures with or without secondarily generalized seizures in patients with epilepsy aged 12 years and older" | 2-12-2016 |
| 22 | Azilsartan Medoxomil Bulk & 40mg/80mg Tablets | "Indicated for the treatment of hypertension in adults patients, either alone or in combination with other antihypertensive agents" | 9-12-2016 |

| Sl. No. | Name of Drug | Indication | Date of issue |
|---------|--|---|---------------|
| 1 | Hydrocortisone Aceponate 0.584mg/ml Cutaneous Spray Solution (Vet.) | For symptomatic treatment of inflammatory and pruritic dermatosis in dogs. | 6-1-2017 |
| 2 | Dexlansoprazole Delayed Release Capsule 30/60mg & Bulk | For the treatment of (i) Healing of all grades of erosive esophagitis (EE). (ii) Maintaining healing of EE and relief of heartburn. (iii) Treating heartburn associated with symptomatic nonerosive gastroesophageal reflux disease (GERD). | 16-1-2017 |
| 3 | Carfilzomib Sterile Lyophilized Powder for Injection 60mg/vial (50ml vial) | Relapsed or refractory multiple myeloma Carfilzomib for injection is indicated in combination with dexamethasone or with lenalidomide plus dexamethasone for the treatment of patients with relapsed or refractory multiple myeloma who have received one to three lines of therapy. Carfilzomib for injection is indicated as a single agent for the treatment of patients with relapsed or refractory multiple myeloma who have received one or more lines of therapy. | 17-1-2017 |
| 4 | Dabrafenib 50mg/75mg Capsules (Dabrafenib Mesylate) | As a single agent for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E mutation as detected by an appropriate test. In combination with Trametinib for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E mutation as detected by an appropriate test. | 18-1-2017 |

| _ | Transfinit () 5mg/m - T-L1-t- | As a manothermy and in anathropia. 24 Data 6 27 6 | 10 1 2017 | |
|----|--|---|------------------------------|--|
| 5 | Trametinib 0.5mg/2mg Tablets (Trametinib Dimethyl Sulfoxide) | As a monotherapy and in combination with Dabrafenib for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E mutation as detected by an appropriate test. | 18-1-2017 | |
| 6 | Alectinib 150mg Capsules (Alectinib Hydrochloride) | For the treatment of patients with anaplastic lymphoma kinase (ALK)-Positive, metastatic non-small cell lung cancer (NSCLC) who have progressed on or are intolerant to Crizotinib. | 23-1-2017 | |
| 7 | Eliglustat 84mg Capsules (Eliglustat Tartrate or Hemitartrate Salt) For the long term treatment of adult patients with Gaucher disease type 1 who are CYP2D6 extensive metabolizers (EMs), intermediate metabolizers (IMs), or poor metabolizers (PMs) as detected by an appropriate test. | | 31-1-2017 | |
| 8 | Pasireotide long-acting release 20/40/60mg powder for suspension for injection (Additional Indication) (Pasireotide pamoate) | For the treatment of patients with acromegaly who have had an inadequate response to surgery and/ or for whom surgery is not an option. | egaly who have had 14-2-2017 | |
| 9 | Etonogestrel 68 mg implant | For use by women to prevent pregnancy. | 15-2-2017 | |
| 10 | Bepotastine Besilate 1.5 %w/v Ophthalmic solution | For the treatment of itching associated with allergic conjunctivitis | 22-2-2017 | |
| 11 | Dienogest 2mg Tablets | For the management of Pelvic pain associated with Endometriosis | | |
| 12 | Eprinomectin 0.5% for beef and dairy cattle (vet.) | It is indicated for the treatment and control of gastrointestinal roundworm (including inhibited Ostertagia ostertagi), lungworm, grubs, sucking and biting lice, chlorioptic and sarcoptic-mange mites, and horn flies in beef and dairy cattle of all ages, including lactating dairy cattle. | 14-3-2017 | |
| 13 | Bepotastine Besilate 10 mg Tablets | For the treatment of allergic rhinitis | 27-3-2017 | |
| 14 | Prucalopride 1mg/2mg Tablet & Bulk (Prucalopride Succinate) | | | |
| 15 | Teriflunomide 14 mg tablets | For the treatment of patients with relapsing form of multiple sclerosis | 13-4-2017 | |
| 16 | Pomalidomide 1mg/2mg/3mg/4mg Capsules & Bulk | In combination with dexamethasone, for patient with patient multiple myeloma who have received at least two prior therapies including lenalidomide and a proteasome inhibitor and have demonstrated disease progression on or within 60 days of completion of the last therapy. | 1-5-2017 | |
| 17 | Sofosbuvir 400 mg + Velpatasvir 100 mg Tablet For the treatment of adult patients with - chronic Hepatitis C virus, Genotype 1, 2, 3, 4, 5 or 6 infection. - Without cirrhosis or with compensated cirrhosis - With decompensated with chronic for use in combination with Ribavirin. | | 4-5-2017 | |
| 18 | Velpatasvir Bulk | | 4-5-2017 | |
| 19 | Osimertinib 40 mg/80 mg Film coated Tablets (Osimertinib Mesylate) | For the treatment of patient with metastatic epidermal growth factor receptor (EGFR) T790 M mutation-positive non-small cell lung cancer (NSCLC), as detected by an appropriate test, whose disease has progressed on or after EGFR TKI therapy | 29-5-2017 | |

| 20 | Argatroban Hydrate Bulk & Injection 250 mg/2.5 ml | (1) For prophylaxis or treatment of thrombosis in adult patients with Heparin induced thrombocytopenia (HIT). | 30-5-2017 |
|----|--|---|-----------|
| | | (2) As an anticoagulant in adults patients with or at risk for heparin-induced thrombocytopenia undergoing percutaneous coronary intervention (PCI) | |
| 21 | Fluralaner chewable tablets 112.5 mg/250 mg/500mg/1000mg/1400 mg | For the treatment of tick and flea infestations on dogs for 12 weeks. This veterinary medicinal product is a systemic insecticide and acaricide with a long duration of action that provides immediate and persistent tick (adult and juvenile Ixodes ricinus, Ixodes hexagonus, Ixodes scapularis, Ixodes holocyclus, Dermacentor reticulates, Dermacentor variabilis and Rhipicephalus sanguineus) and flea (Ctenocephalides felis and Ctenocephalides canis) killing activity for 12 weeks. Fleas and ticks must attach to the host and commence feeding in order to be exposed to the active substance. the onset of effect is within 8 hours of attachment for fleas (C. felis) and 12 hours of attachment for ticks (I. ricinus) The product effectively controls environmental flea population in area to which treated dogs have access. The product can be used as part of an treatment strategy for the control of Flea Allergy Dermatitis (FAD) | 30-5-2017 |
| 22 | Fluticasone Furoate (100 mg mcg) and Vilanterol Trifenatate (25 mcg) powder for inhalation | Indicated for the maintenance treatment of the airflow obstruction in patients with chronic obstructive pulmonary disease (COPD) including chronic bronchitis and/ or emphysema and to reduce exacerbation of COPD in patients with an exacerbation history. | 29-6-2017 |
| 23 | Mirabegron Prolonged Release Tablet 25 mg/50 mg | Symptomatic treatment of urgency, increased micturition frequency and / or urgency incontinence as may occur in patients with over active bladder (OAB) Syndrome | 12-7-2017 |
| 24 | Delamanid 50 mg tablet | For the use as part of an appropriate combination regimen for pulmonary multi-drug resistant tuberculosis (MDR-TB) in adult patients when an effective treatment regimen cannot otherwise be composed for reasons of resistance or tolerability | 2-8-2017 |
| 25 | Mirabegron Prolonged Release Tablet 25 mg /50 mg & Bulk | Symptomatic treatment of urgency, increased micturition frequency and / or urgency incontinence as may occur in patients with over active bladder (OAB) Syndrome | 18-8-2017 |
| 26 | Efonidipine Hydrochloride Ethanolate Bulk & Tablets 10 mg/20mg/40mg | Indicated for the management of • Hypertension • Renal parenchymal hypertension • Angina | 28-8-2017 |
| 27 | Brivaracetam Film Coated Tablets 50mg/75mg/100mg | As adjunctive therapy in the treatment of partial -onset seizures in patients 16 years of age and older with epilepsy | 7-9-2017 |
| 28 | Treosulfan Bulk & injection 5g/vial | For the conditioning treatment prior to haematopoetic stem- cell transplantation | 26-9-2017 |
| 29 | Ribociclib 200 mg Film coated Tablets | In combination with an aromatase inhibitor as initial endocrine-based therapy for the treatment of post menopausal women with the hormone receptor (HR)-Positive, human epidermal growth factor receptor 2 (HER2)-Negative advanced or metastatic breast cancer | 27-9-2017 |

| 30 | Dienogest Bulk & 2mg Tablet | For the management of Pelvic pain associated with Endometriosis | 28-9-2017 |
|----|---|---|------------|
| 31 | Apremilast bulk & film coated tablets 10 mg/20 mg/30 mg | for treatment of patients with moderate to severe plaque psoriasis who are candidate for phototherapy or systemic therapy | 13-10-2017 |
| 32 | Arbekacin bulk & Injection 200mg/4ml Treatment of following infections caused by Methicillin resistant Staphylococcus aureus (MRSA)", sepsis pneumonia | | 20-10-2017 |
| 33 | Benestermycin Intra-mammary suspension (vet.) | For the treatment of subclinical mastitis at drying off, and the prevention of new bacterial infections of the udder during the dry period in dairy cows, caused by bacteria susceptible to penicillin and framycetin | 3-11-2017 |
| 34 | Midostaurin 25 mg Capsules | • In combination with standard induction and consolidation chemotherapy followed by single agent in maintenance of therapy for adult patients with newly diagnosed with acute myeloid leukemia (AML) who are FLT-3 Mutation positive. | 9-11-2017 |
| | | For the treatment of adult patients with advanced systemic mastocycosis (Advanced SM) | |
| 35 | Tenofovir Alafenamide Fumarate bulk & 25 mg Tablets | For the treatment of chronic Hepatitis B virus infection in adults with compensated liver disease | 10-11-2017 |
| 36 | Arteolane Maleate and Piperaquine phosphate Dispersible tablets (37.5 mg +187.5 mg | Indicated in children aged 6 months to 12 years for the treatment of: - Acute uncomplicated Plasmodium falciparum malaria infection - Acute uncomplicated Plasmodium Vivax malaria | 24-11-2017 |
| | | infection | |
| 37 | Macitentan Bulk and Tablets 10 mg | Macitentan is an endothelin receptor antagonist (ERA)l indicated for the treatment of Pulmonary arterial hypertension (PAH, WHO group I) to delay disease progression | 7-12-2017 |

$Manufacturing \ and \ Marketing \ permission \ is sued \ from \ SND \ Division \ from \ 01.01.2018 \ to \ 31.12.2018$

| Sl. No. | Drug Name | Composition | Indication | Date of Approval |
|---------|--|---|---|------------------|
| 1 | Pegaspargase injection 3750 IU per vial | Each vial contains: Pegaspargase 3750 IU per vial | As a component of multi agent chemotherapeutic regiment for the treatment of patients with acute lymphoblastic leukaemia who are hypersensitive to L-Asparaginase. | 04-01-2018 |
| 2 | , | Each 5ml reconstituted suspension contains: Cefditoren Pivoxil equivalent to Cefditoren 100mg | For the treatment of mild to moderate infection also in children's of (2 months to 12 years of age) which are caused by susceptible strains of the designated microorganisms in the condition listed below: • Acute Bacterial Exacerbation of Chronic Bronchitis • Community-Acquired pneumonia • Pharyngitis/Tonsillitis • Uncomplicated Skin and skin - structure infection | 04-01-2018 |
| 3 | Eltrombopagolamine tablets 12.5mg (additional strength & indication) | Entrollioopag olalilile equivalent | 1. Chronic ITP: for Adults For the treatment of thrombocytopenia in patients with chronic immune (idiopathic) Thrombocytopenic purpura (ITP) who have had an insufficient response to | 06-02-2018 |

| | | | corticosteroids, immunoglobulins, or splenectomy. (It should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding. It should not be used in an attempt to normalize platelet counts). For Paediatric patients aged 1 year and above: Thrombocytopenia in paediatric patients 1 year and older with immune (idiopathic) thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins or splenectomy (It should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding. It should not be used in an attempt to normalize platelet counts). 2. For the treatment of patients with chronic Hepatitis C virus (HCV) infection for the treatment of thrombocytopenia to: Enable the initiation of interferon based therapy, optimize interferon based therapy. With the condition: to be sold by retail on the prescription of Haematologist only | |
|---|---|---|---|------------|
| 4 | Alectinib Capsules 150mg (additional indication) | contains: | First-line treatment of patients with Anaplastic Lymphoma Kinase (ALK) - positive locally advanced or metastatic non-small cell lung cancer (NSCLC). With the condition: to be sold by retail on the prescription of Oncologist only. | 05-03-2018 |
| 5 | Fulvestrant 250mg/5ml injection (additional indication) | Each 5ml pre-filled syringe contains: Fulvestrant 250mg ethanol (96%)-10% w/v | For the treatment of estrogen receptor positive, locally advanced or metastatic breast cancer in postmenopausal women not previously treated with endocrine therapy With the condition: to be sold by retail on the prescription of Oncologist only. | 20-03-2018 |
| 6 | Bendamustine hydrochloride injection 90mg/ml (0.5ml vial & 2.0 ml vial) (additional indication) | Pandamustina hydrochlorida | Front line treatment of multiple myeloma (Durie-Salmon stage II with progress or stage III) in combination with prednisone for patients older than 65 years who are not eligible for autologous stem cell transplantation and who have clinical neuropathy at time of diagnosis precluding the use of thalidomide or bortezomib containing treatment. With the condition: to be sold by retail on the prescription of Oncologist/Specialist only. | 23-03-2018 |
| 7 | Methotrexate injection prefilled syringe 15mg/0.3ml, 20mg/0.4ml and 25mg/0.5ml (Additional strength & indication) | contains: Methotrexate 15mg | It is indicated in treatment of 1. Active rheumatoid arthritis in adult patients. 2. Polyarthritic forms of severe, active juvenile idiopathic arthritis, when the response to nonsteroidal anti-inflammatory drug (NSAIDs) has been inadequate, 3. Severe recalcitrant disabling psoriasis, which is not adequately responsive to other forms of therapy such as phototherapy, PUVA, and retinoids and severe psoriatic arthritis in adult patients. 4. Mild to moderate Crohn's disease either alone or in combination with corticosteroids in adult patients refractory or intolerant to thiopurines. | 09-03-2018 |

| 8 | Sildenafil power for oral | Each ml of reconstituted | Adults: | 20-03-2018 |
|----|--|--|---|------------|
| Ü | _ | suspension contains: Sildenafil citrate equivalent to 10mg, Sodium benzoate 4.375mg | Treatment of adult patients with pulmonary arterial hypertension classified as WHO functional class II and III, to improve exercise capacity. Efficacy has been shown in primary pulmonary hypertension and pulmonary hypertension associated with connective tissue disease. | |
| | | | Paediatric population: | |
| | | | Treatment of paediatric patients aged 1 year to 17 years old with pulmonary arterial hypertension. Efficacy in terms of improvement of exercise capacity or pulmonary haemodynamics has been shown in primary pulmonary hypertension and pulmonary hypertension associated with congenital heart disease. | |
| | | | With the condition: to be sold by retail on the prescription of Cardiologist only. | |
| 9 | Dydrogestero ne tablet 10mg (additional indication) | Each film coated tablet contains: Dydrogesterone 10mg | Luteal support as part of an Assisted Reproductive Technology (ART) treatment. | 06-04-2018 |
| | | | With the condition: to be sold by retail on the prescription of Gynaecologist only. | |
| 10 | Febuxostat tablet 60mg & 20mg (Additional Strength) | Each film coated tablet contains: Febuxostat 60mg & 20mg | For the treatment of chronic hyperuricemia in conditions where urate deposition has already occurred (including a history, or presence of tophus and/or gouty arthritis) | 06-04-2018 |
| 11 | | Each injection contains: Posaconazole 300mg/16.7ml | For prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy. With the condition: to be sold by retail on the prescription of Specialist only. | 13-04-2018 |
| 12 | Teriflunomide tablet 7mg (Add. Lower strength) | Each film coated tablet contains: Teriflunomide 7mg | For the treatment of patients with relapsing form of multiple sclerosis. With the condition: to be sold by retail on the prescription of Neurologist only. | 07.05.2018 |
| 13 | Apremilast film coated tablet 10/20/30mg (Additional Indication) | Each film coated tablet contains: Apremilast-10mg, 20mg,30mg | For the treatment of adult patients with Active psoriatic Arthritis. | 11.05.2018 |
| 14 | Aripiprazole tablet 2mg (additional strength) | Each uncoated tablet contains: Aripiprazole-2mg | For the treatment of Schizophrenia in adolescent patients (13 to 17 years of age) With the condition: to be sold by retail on the prescription of Psychiatrist only. | 11.05.2018 |
| 15 | Dextromethor phan HBr Lozenges 10mg (Add. dosage form & strength) | Each lozenges contains: Dextromethorpha n HBr-33.33% Complex 30mg eq. to Dextromethorpha n HBr 10mg | Cough suppressant for the relief of acute nonproductive (dry, tickly) cough associated with respiratory tract infection in adult patients only. | 11.05.2018 |
| 16 | Voglibose Orally Disintegrating Strips 0.2mg (Modified dosage form) | Each Orally Disintegrating Strips contains: Voglibose 0.2mg & 0.3mg | For improvement of parandial hyperglycaemia in diabetes mellitus, when only diet and/or exercise or oral hypoglycaemic drugs or insulin preparation in addition to diet and/or exercise do not result in adequate glycemic control. | 11.05.2018 |

| 17 | Tranexamic acid tablet 1000mg (Additional strength) | Each film coated tablet contains: Tranexamic acid-1000mg. | For the treatment of menorrhagia | 21.05.2018 |
|----|--|--|---|------------|
| 18 | Eltrombopag Olamine Tablet 75mg (Additional strength) | Each film coated tablet contains: Eltrombopag Olamine as free acid-75mg | 1. For the treatment of thrombocytopenia in patients with chronic immune (idiopathic) thrombocytopenic purpura (ITP) who have had an insufficient response to corticosteroids, immunoglobulins or splenectomy.(It should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding. It should not be used in an attempt to normalize platelet counts). | 29.05.2018 |
| | | | 2. Indicated in patients with chronic hepatitis C virus (HCV) infection for the treatment of thrombocytopenia to: a. Enable the intiation of interferon based therapy. b. Optimize interferon based therapy | |
| | | | 3. For the treatment of patients with severe aplastic anaemia (SAA) who have had an insufficient response to immunosuppressive therapy | |
| | | | 4. For the treatment of thrombocytopenia in paediatric patients 1 year and older with chronic immune (idiopathic) thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins or splenectomy. (It should be used only in patients with ITP whose degree of thrombocytopenia and clinical condition increase the risk for bleeding. It should not be used in an attempt to normalize platelet counts). With the condition: to be sold by retail on the prescription of Hematologist only. | |
| 19 | Thyroxine Sodium Tablet 37.5 mcg (Additional strength) | Each uncoated tablet contains: Thyroxine Sodium eq. to Anhydrous Thyroxine Sodium. 37.5 mcg. | 1. As replacement or supplemental therapy in patients of any age or state (including pregnancy) with hypothyroidism of any etiology except transient hypothyroidism during the recovery phase of sub-acute thyroidit is resulting from thyroid dysfunction, primary atrophy or partial or total absence of the thyroid gland or from the effects of surgery, radiation or drugs with or without the presence of goiter, including sub-clinical hypothyroidism; secondary (pituitary) hypothyroidism and tertiary (hypothalamic) hypothyroidism. | 08.06.2018 |
| | | | 2. As a pituitary TSH suppressant in the treatment or prevention of various types of euthyroidgoites, including thyroid nodules, sub-acute or chronic lymphocytic thyroiditis (Hashimoto's) multi-nodular goiter and in conjuction with surgery and radioactive iodine therapy in the management of thyrotropin-dependent well differentiated papillary or follicular carcinoma of the thyroid. | |
| 20 | Dabrafenib Hard Gelatin Capsule 50mg,&75mg | Each Hard Gelatin Capsule contains: | Dabrafenib in combination with Trametinib for the treatment of adult patients with advanced non-small cell lung cancer with BRAF V600 mutation. | 08.06.2018 |
| | (Additional indication) | Dabrafenib Mesylate eq. to Dabrafenib 50mg & 75mg | With the condition: to be sold by retail on the prescription of Oncologist only. | |
| 21 | Trametinib Tablet 0.5mg & 2mg (Additional indication) | Each Film coated tablet contains: Trametinib 0.5mg & 2mg | Trametinib in combination with Dabrafenib for the treatment of adult patients with advance non-small cell lung cancer with a BRAF V 600 mutation. With the condition: to be sold by ratail on the prescription of | 08.06.2018 |
| | | | With the condition: to be sold by retail on the prescription of Oncologist only. | _ |

| | I | | ı | 1 |
|----|---|--|--|------------|
| 22 | Azacitidine for injection 50mg/vial (Additional strength) | Azacitidine. 50mg Lyophilized | For the treatment of adult patients with all subtypes of Myelodysplastic Syndrome. With the condition: to be sold by retail on the prescription of Oncologist only. | 22.06.2018 |
| 23 | Nilotinib Capsules 50mg (Additional strength & add. indication) | Each Capsule contains: Nilotinib 50mg | Paediatric patients with newly diagnosed Philadelphia Chromosome Positive Chronic Myelogenous Leukaemia (CML) in the chronic phase. | 22.06.2018 |
| | | | 2. Paediatric patients with chronic phase Philadelphia Chromosome Positive (CML) with resistance or intolerance to prior therapy including imatinib. | |
| | | | With the condition: to be sold by retail on the prescription of Oncologist only. | |
| 24 | Fulvestrant 250mg/5ml injection (additional indication) (for IM use only) | Each 5ml pre-filled syringe contains: Fulvestrant USP 250mg Ethanol (96%) BP-10% w/v | In combination with Palbociclib for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer in women who have received prior endocrine therapy". | 13.07.2018 |
| | | | With the condition: to be sold by retail on the prescription of Oncologist only. | |
| 25 | Ganciclovir Sodium powder for Infusion 500mg (additional indication) (for IV use only) | | For the treatment of cytomegalovirus (CMV) disease in immunocompromised individuals and for the prevention of CMV disease in patients with drug-induced immunosuppression following organ transplantation or cancer chemotherapy. | 13.07.2018 |
| | | | With the condition: to be sold by retail on the prescription of a Specialist only. | |
| 26 | Nepafenac Ophthalmic Suspension 0.3%w/v (Additional strength) | Each ml contains: Nepafenac 3.0mg | For the treatment of pain and inflammation associated with cataract surgery. With the condition: to be sold by retail on the prescription of Ophthalmologist only. | 13.07.2018 |
| 27 | Tadalafil Orally Disintegrating Strip 5mg (Additional strength) | Each Orally Disintegrating Strip contains: Tadalafil 5mg | For the treatment of Erectile Dysfunction | 13.07.2018 |
| 28 | Delamanid Tablets 50mg (additional indication) | Each film coated tablet contains: Delamanid 50mg | In exceptional cases and after a careful benefit/risk assessment Delamanid treatment can be initiated in children >6 years of age, for use as part of an appropriate combination regimen for pulmonary multi-drug resistant tuberculosis (MDR-TB) when an effective treatment regimen cannot otherwise be composed for reasons of resistance or tolerability. With the condition: For the use in Revised National Tuberculosis Control Program (RNTCP). | 27.07.2018 |
| 29 | Osimertinib Film coated tablet 40mg and 80mg (Additional indication) | Each film coated tablet contains: Osimertinib 40mg, 80mg | As first-line treatment of patients with locally advanced or metastatic Non-Small Cell Lung Cancer (NSCLC) whose tumors have Epidermal Growth Factor Receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitutions mutations. With the condition: To be sold by retail on the prescription of Oncologist only. | 03.08.2018 |

| 30 | Nicotine Polacrilex Orally disintegrating strips 2mg & 4mg (New dosage form) | Each Orally disintegrating strips contains: Nicotine Polacrilex eq. to Nicotine 2mg, 4mg | To reduce the withdrawal symptoms, including nicotine craving, associated with quitting the smoking and quitting chewed tobacco and gutkha containing tobacco | 17.08.2018 |
|----|--|--|---|------------|
| 31 | Zolmitriptan Orally disintegrating strips 2.5mg & 5mg (Additional Dosage form) | Each Orally disintegrating strips contains: Zolmitriptan 2.5mg & 5mg | For Acute treatment of migraine with or without aura in adults. | 10.08.2018 |
| 32 | Ethyl esters of Iodidzed fatty acids of Poppy Seed Oil solution for injection (Lipodol Ultra Fluid 10ml) (Additional Indication) | Each ampoule (10ml) contains: Ethyl esters of Iodidzed fatty acids of Poppy Seed Oil Iodine content 48% i.e. 480mg iodine per ml | Conventional Trans-Arterial Chemo-Embolization (cTACE) of Hepatocellular Carcinoma (HCC) in adults. | 06.09.2018 |
| 33 | Methylcobala min nasal spray 250mcg/spray (New Dosage form) | Each 0.1ml contains: Methylcobalamin 500mcg Benzalkonium chloride (As preservative) 0.02% w/v | Treatment of Vitamin B12 deficiency in patients without having neurological involvement. | 28.09.2018 |
| 34 | Apomorphine Hydrochloride injection 10mg/ml (2ml and 5ml Ampoule) (subcutaneous use only) (Add. Dosage form and add. indication) | Each ml contains: Apomorphine hydrochloride 10mg | Treatment of disabling motor fluctuations ("on-off" phenomena) in patients with Parkinson's disease which persist despite individually titrated treatment with levodopa (with a peripheral decarboxylasee inhibitor) and/or other dopamine agonists. With the condition: to be sold by retail on the prescription of Neurologist only. | 28.09.2018 |
| 35 | Carfilzomib for Injection 30mg/vial (Lyophilized injection) (Additional Strength) | Each vial contains: Carfilzomib 30mg | Carfilzomib for injection is indicated in combination with dexamethasone or with lenalidomide plus dexamethasone for the treatment of patients with relapsed or refractory multiple myeloma who have received one to three lines of therapy. Carfilzomib for injection is indicated as a single agent for the treatment of patients with relapsed or refractory multiple myeloma who have received one or more lines of therapy. With the condition: to be sold by retail on the prescription of Oncologist only. | 28.09.2018 |
| 36 | Nilotinib hard Capsules 150mg & 200mg (Additional indication) | Each Capsule contains: Nilotinib 150mg & 200mg | Treatment of Paediatric patients with newly diagnosed Philadelphia Chromosome Positive Chronic Myelogenous Leukaemia (CML) in the chronic phase. Treatment of Paediatric patients with chronic phase Philadelphia Chromosome Positive (CML) with resistance or intolerance to prior therapy including imatinib. With the condition: to be sold by retail on the prescription of Oncologist only. | 18.10.2018 |
| 37 | Dasatinib film coated tablet 20mg, 50mg, 70mg and 100mg (Additional indication) | Each film coated tablet contains: Dasatinib 20mg, 50mg, 70mg and 100mg | Treatment of Paediatric patients with Philadelphia Chromosome Positive Chronic myeloid leukemia (Ph+CML) in chronic phase. With the condition: to be sold by retail on the prescription of Oncologist only | 18.10.2018 |

| 38 | _ | Each Extended release film | Vasospastic Angina, | 18.10.2018 |
|----|---|---|---|------------|
| | Tablet 60mg (Add. Strength and indication) | | Chronic stable Angina (Classical effort-Associated Angina), | |
| | | Nifedipine 60mg | • Hypertension. | |
| | | | With the condition: to be sold by retail on the prescription of Cardiologist only. | |
| 39 | 500mg/100ml, 1000 | contains: Levetiracetam in 0.82% sodium chloride (500mg/100ml) Levetiracetam in 0.75% sodium chloride | Partial onset seizures, Myoclonic seizures in patients with juvenile myoclonic epilepsy, Primary generalized tonic-clonic seizures in adults with | 27.10.2018 |
| 40 | | Each ml contains: Nepafenac 3.0mg | For the treatment of post-operative pain and inflammation associated with cataract surgery. With the condition: to be sold by retail on the prescription of Ophthalmologist only. | 09.11.2018 |
| 41 | Ibuprofen soft gelatine capsules 400mg. (New Dosage form) | Each soft gelatin capsule contains Ibuprofen 400mg | For chronic arthritic disorder and painful musculoskeletal conditions. | 20.11.2018 |
| 42 | Everolimus tablets 2.5 mg (Additional Indication) | Each tablets contains: Everolimus 2.5 mg | Treatment of patients with Progressive neuroendocrine tumors of pancreatic origin (PNET) with unresectable, locally advanced or metastatic disease, Treatment of patients with Advanced Renal Cell Carcinoma | 05.12.2018 |
| | | | (RCC). With the condition: To be sold by retail on the prescription of Oncologist only. | |
| 43 | Dissolving tablets 0.25/0.5/1.0/2. Omg (New | Each uncoated mouth dissolving tablet contains: Clonazepam 0.25/0.5/1.0/2.0 mg | Treatment of seizure disorders (Petit mal and its variant, akinetic and myoclonic seizures), Panic disorder & add-on therapy in acute mania. | 10.12.2018 |
| | Dosage Form) | | With the condition: To be sold by retail on the prescription of Neurologists or Psychiatrist only. | |
| 44 | Divalproex Sodium Oral Solution 250mg/5ml (Additional Strength) | Each 5 ml contains: Divalproex 250mg | Monotherapy and adjunctive therapy in the treatment of patients with 'complex partial seizures that occur either in isolation or in association with types of seizures for adult patients only. | 21.12.2018 |
| 45 | Ivermectin Cream 1.0% w/w (New Dosage form) | Each Gram contains: Ivermectin 1.0% w/w | Treatment of inflammatory lesions of rosacea. With the condition: To be sold by retail on the prescription of Dermatologist only. | 21.12.2018 |

List of drugs approved from SND Division from 01.01.2019 to 31.12.2019.

| Sl. No. | Drug Name | Composition | Indication | Date of Approval |
|---------|---|--|---|------------------|
| 1 | Donepezil Hydrochloride Orally Disintegrating Strips (Additional osage form) | contains: | For the treatment of mild to moderate severe Alzheimer's Dementia. With the condition: To be sold by retail on the prescription of Neurologists only. | 11.01.2019 |
| 2 | Tapentadol Hydrochloride capsule 50mg (New dosage form) | Each capsule contains: Tapentadol Hydrochloride 50mg | For relief of moderate to severe acute pain in adults 18 years of age or older. | 30.01.2019 |
| 3 | Midodrine Hydrochloride tablet 2.5mg, 5mg and 10mg (Additional strength) | Each tablet contains: Midodrine Hydrochloride 2.5mg, 5mg and 10mg | It is indicated in the treatment of symptomatic orthostatic hypotension. With the condition: To be sold by retail on the prescription of Cardiologists only. | 20.02.2019 |
| 4 | Ribociclib film coated tablet 200mg (Additional indication) | | Ribocicib in combination with an aromatase inhibitor for the treatment of pre/peri-menopausal or post-menopausal women, with hormone receptor (HR)-positive, human epidermal growth factor receptor 2(HER2)-negative advanced or metastatic breast cancer, as initial endocrine-based therapy. With the condition: To be sold by retail on the prescription of an Oncologist only. | 13.03.2019 |
| 5 | Azacitidine for injection 150mg/vial (Additional strength) | Azacitidine. 150mg Lyophilized | For the treatment of adult patients with all subtypes of Myelodysplastic Syndrome. With the condition: to be sold by retail on the prescription of an Oncologist only. | 25.03.2019 |
| 6 | Lenvatinib capsule 4mg/10mg (Additional Indication) | Each capsule contains: Lenvatinib mesylate 4.90mg eq. To Lenvatinib 4mg, Lenvatinib mesylate 12.25 mg eq. To Lenvatinib 10mg. | For the first line treatment of patients with unresectable hepatocellular carcinoma. With the condition: to be sold by retail on the prescription of an Oncologist only. | 26.03.2019 |
| 7 | Primidone tablet 50mg (Additional Strength) | Each uncoated tablet contains: Primidone USP 50mg | It is indicated for the management of grand mal and psychomotor (temporal lobe) epilepsy. It is also of value in the management of focal or jacksonian seizures, myoclonic jerks and akinetic attacks. Management of essential tremor. With the condition: To be sold by retail on the prescription of an Neurologist only. | 15.04.2019 |
| 8 | Amisulpride orodispersible tablet 50/100/200/300/4 00mg (New dosage form) | _ | For acute and chronic schizophrenic disorders, in which positive and negative symptoms are prominent, including patients characterised by predominant negative symptoms. With the condition: To be sold by retail on the prescription of a Psychiatrist only. | 03.05.2019 |
| 9 | Progesterone solution for Injection 25mg/1.119ml vial (Add. Dosage form and add. route of administration) | Progesterone 25mg (22.35mg/ml) | It is indicated in adults for luteal support as part of an Assisted Reproductive Technology (ART) treatment program in infertile women who are unable to use or tolerate vaginal preparations. With the condition: To be sold by retail on the prescription of Gynaecologists only. | 17.05.2019 |

| 10 | Azilsartan medoxomil tablet 20/40/80mg (Additional lower Strength) | Each uncoated tablet contains: Azilsartan Kamedoxomil equivalent to Azilsartan medoxomil 20/40/80mg | Indicated for the treatment of hypertension in adult patients either alone or in combination with other antihypertensive agents. With the condition: To be sold by retail on the prescription of a Cardiologist only. | 27.05.2019 |
|----|---|---|--|------------|
| 11 | Everolimus Tablet 0.25mg/0.5mg/0.75mg/1.0mg (Additional indication) | Each film coated tablet contains: Everolimus 0.25mg/0.5mg/0. 75mg/1.0mg | Prophylaxis of organ rejection in adult patients at low to moderate immunological risk receiving an allogeneic renal or cardiac transplant. In kidney and heart transplantation, Certican should be used in combination with ciclosporin for microemulsion and Corticosteroids. | 27.05.2019 |
| | | | With the condition: To be sold by retail on the prescription of Cardiologist only. | |
| 12 | Diclofenac sodium 2% Topical Solution (New route of administration) | Each 1 gram of solution contains: Diclofenac sodium 20mg | For the treatment of pain of osteoarthritis of the knees. | 03.06.2019 |
| 13 | Artesunate for injection 120mg with phosphate buffer 0.3M (Additional strength) | Each combi pack contains: a) Artesunate for injection 120mg Each vial contains: Artesunate IP 120mg b) Phosphate Buffer 0.3M (12ml) Each ampoule contains: Anhydrous Disodium Hydrogen Phosphate BP 4.0% w/v Potassium dihydrogen phosphate | | 26.06.2019 |
| | | BP 0.2% w/v | | |
| 14 | Tranexamic acid injection IP 100mg/ml (Additional Indication) | Each ml contains: Tranexamic acid IP 100 mg/ml | Indicated for short term use in prophylaxis and treatment in patients at high risk of per-and post-operative hemorrhage following: | 26.07.2019 |
| | | | a) Prostatectomy.b) Conisation of the cervix.c) Surgical procedures and extractions in hemophiliacs | |
| 15 | Rivaroxaban tablets 2.5mg (Add. Indication) | Rivaroxaban tablets 2.5mg | Rivaroxaban 2.5mg tablet, co-administered with acetylsalicylic acid (ASA), is indicated for the prevention of atherothrombotic events in adult patients with coronary artery disease (CAD) or symptomatic peripheral artery disease (PAD) at high risk of ischaemic events. With the condition: To be sold by retail on the prescription of Cardiologists only. | 30.07.2019 |
| 16 | Olaparib 100 mg and 150 mg tablet (Additional Indication) | Olaparib 100 mg and 150 mg tablet | for the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy With the condition: To be sold by retail on the prescription of Oncologist only. | 30.07.2019 |
| 17 | | Tapentadol Hydrochloride 25.000 milligram (mg) Capsules | | 30.07.2019 |

| 10 | T | The state of the s | The state of the s | 07.00.2010 |
|----|--|--|--|------------|
| 18 | Injection Powder for prolonged-release suspension for injection (Depot injection for I.M. use only) | acetate 22.5mg | Treatment of advanced Prostatic cancer. With the condition: To be sold by retail on the prescription of Oncologist only. | 07.08.2019 |
| | (Add. strength) | | | |
| 19 | Clobazam Tablets IP 2.5mg & 7.5mg (Add. lower strength). | Each tablet contains: Clobazam 2.5mg & 7.5mg | As an adjunctive therapy in patients with refractory epilepsy. With the condition: To be sold by retail on the prescription of Neurologist/Psychiatrist only. | 07.08.2019 |
| 20 | Sildenafil for oral suspension 10mg/ml (Add. Indication) | Each ml of reconstituted suspension contains: Sildenafil citrate IP equivalent to Sildenafil 10mg | For the treatment of pulmonary arterial hypertension (WHO Group I) in adults to improve exercise ability and delayed clinical worsening. With the condition: To be sold by retail on the prescription of Cardiologist only. | 09.08.2019 |
| 21 | Pantoprazole dual-release gastro- resistant tablet 80mg (New Dosage form) | Each dual-release gastro-resistant tablet contains: Pantoprazole sodium equivalent to Pantoprazole 80mg | For the treatment of moderate to severe refractory Gastro- oesophageal reflux disease (GERD) for 4 weeks only. | 02.09.2019 |
| 22 | Pirfenidone tablet 400 mg (Add. Strength) | Each film coated tablet contains : Pirfenidone 400mg | For the treatment of idiopathic pulmonary fibrosis. With the condition: To be sold by retail on the prescription of Pulmonologist only. | 02.09.2019 |
| 23 | Olopatadine HCI ophthalmic solution 0.7% w/v (Add. strength) | Each ml contains: Olopatadine HCI 0.7% w/v Benzalkonium chloride 0.015% w/v | | 06.09.2019 |
| 24 | Ibrutinib hard gelatin capsule 140mg (Add. indication) | Each capsule contains: Ibrutinib 140mg | For the treatment of patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL). With the condition: To be sold by retail on the prescription of Oncologist only. | 11.09.2019 |
| 25 | Ibrutinib hard gelatin capsule 140mg (Add. indication) | Each capsule contains: Ibrutinib 140mg | For the treatment of patients with Waldenstrom's macroglobulinemia | 11.09.2019 |
| 26 | Desonide Ointment 0.05% (Add. Dosage form | Desonide topical ointment 0.05% | for sub-acute to chronic steroid responsive dermatoses. | 26.09.2019 |
| 27 | Atorvastatin calcium tablet 30/60 mg (Add. Strength) | Each film coated tablet contains: Atorvastatin calcium equivalent to Atorvastatin 30/60 mg | Primary Hypercholesterolemia and Mixed Dyslipidemia (Type IIa and IIIb). | 27.09.2019 |
| | | | In adult patients with type II diabetes and without clinically evident coronary heart disease to reduce the risk of myocardial infarction and stroke. To reduce the risk of stroke in adult patients without clinically evident coronary heart disease but multiple risk factors for coronary heart disease | |
| | | | With the condition: To be sold by retail on the prescription of Cardiologist only. | |

| 28 | Desloratadine tablets 10mg (Additional strength) | Each film coated tablet contains: Desloratadine 10mg | For the treatment of Chronic Idiopathic Urticaria in patients who have not responded to the standard 5mg dose of Desloratadine. | 27.09.2019 |
|----|--|--|--|------------|
| 29 | Diclofenac sodium Rectal Solution 25 % w/v with applicator (Add. dosage form) | | Mild to moderate pain in adults. | 27.09.2019 |
| 30 | Perampanel tablets 2mg /4mg/6mg/8mg/1 Omg/12mg (Additional indication) | Each film coated tablet contains: Perampanel 2mg /4mg/6mg/8mg/1 0mg/12mg | For the adjunctive treatment of primary generalized tonic- clonic seizures in patients with epilepsy aged 12 years and older. With the condition: To be sold by retail on the prescription of Neurologist only. | 27.09.2019 |
| 31 | Ribociclib Film Coated Tablet 200mg (Additional Indication) | Ribociclib Succinate 254.40 mg equivalent to Ribociclib 200 mg | Indicated for the treatment of women with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER-2)-negative locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulestrant as initial endocrine based therapy or in women who have received prior endocrine therapy. Warning: To be sold by retail on the prescription of oncologist only. | 04-10-2019 |
| 32 | Cinidipine Tablet 20 mg (Additional Strength) | Each Film Coated Tablets contain Clinidipine 20mg | For the treatment of mild to moderate hypertension | 11-10-2019 |
| 33 | | | For the long term maintenance treatment of airflow obstruction in patients with chronic obstructive pulmonary disease (COPD) including chronic bronchitis and/or emphysema | 14-11-2019 |
| 34 | - | Each film coated sustained release Tablet contains: Cefpodoxime Proxetil, Equivalent to Cefpodoxime 200mg & 400mg | Acute bronchitis, exacerbations of chronic bronchitis, pneumonia, sinusitis, recurrent chronic tonsillitis, pharyngitis and acute otitis. | 03.12.2019 |
| 35 | Ivermectin Topical lotion 0.5% w/w | Each ml contains: Ivermectib 0.5% w/w | For the topical treatment of head lice infestation in patients 6months of age and older. | 03.12.2019 |
| 36 | Lenvatinib Capsules 4mg and 10mg (Additional Indication) | mesylate 4.90 mg equivalent to | In combination with everolimus, for the treatment of patients with advanced renal cell carcinoma (RCC) following one prior anti-angiogenic therapy. Warning: To be sold by prescription of a "Oncologist" only. | 04.12.2019 |
| 37 | Bilastine Oral Solution 2.500 mg/ml (Add. Dosage form) | Each ml contains: Bilastine 2.5000 mg | Symptomatic treatment of allergic rhino- conjunctivitis (seasonal and perennial) and urticaria. Bilastine is indicated in children aged 6 to 11 years with a body weight of at least 20 kg. | 10.12.2019 |
| 38 | Pregabalin Extended Release Tablets 82.50 mg and 165 mg | Each film coated Extended Release Tablet contains: Pregabalin 82.50 mg/165 mg | For treatment of peripheral neuropathic pain in adults. | 12.12.2019 |
| 39 | Clobetasol Propionate Topical Foam 0.5% w/w (Add.dosage form) | contains: Clobetasol Propionate 0.05% w/w Phenoxyethanol 1.0% w/w | Treatment of Inflammatory and pruritic manifestation of corticosteroid responsive dermatoses in patients 12 years of age or older. | 16.12.2019 |

| 40 | Clobetasol Propionate Cream, | Each gram of cream contains: | Indicated for the treatment of moderate to severe plaque | 16.12.2019 |
|----|------------------------------|------------------------------|--|------------|
| | 0.025% w/w | Clobetasol Propionate 0.25mg | psoriasis in patients 18 years of age and older. | |
| | (Add.Strength) | , | | |

List of drugs approved from SND Division from 01.01.2020 to 31.12.2020

| Sl. No. | Drug Name | Composition | Indication | Date of Approval |
|---------|--|---|---|------------------|
| 1 | Betahistine Dihydrochloride Orally Disintegrating Strips 16mg/24mg (New Dosage form) | | Indicated in the treatment of Menier's syndrome, Characterized by unilateral or bilateral tinnitus, vertigo, sensorineural hearing loss | 02-01-2020 |
| 2 | Posaconazole Gatro-Resistant Tablet 100 mg (Modified Release dosage form) | Each-Gastro-Resistant Tablet contains: Posaconazole 100 mg | For Prophylaxis of invasive Aspregilus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia form chemotherapy. Posaconazole Gastro Resistant tablets are indicated in patients 13 years of age and older Warning: To be sold by retail on the prescription of | 03-01-2020 |
| | | | Specialist only. | |
| 3 | Pirfenidone tablet 600 mg (Additional strength) | Each Film coated Tablet contains: Pirfenidone600mg | For the Treatment of idiopathic pulmonary fibrosis Warning: To be sold by retail on the prescription of Pulmonologist only. | 08-01-2020 |
| 4 | Saroglitazar Tablet 2mg/4 mg (Additional indication) | Each uncoated tablet contains: Saroglitazar2/4 mg | Type 2 Diabetes mellitus-as an add-on therapy to metformin | 13-01-2020 |
| 5 | Enzalutamide Soft Capsule 40 mg (Additional indication) | Each soft capsules contains Enzalutamide40 mg | In the treatment of adult men with metastic castration- resistant prostate cancer who are asymptomatic or mildly symptomatic after failure of androgen deprivation therapy in whom chemotherapy is not yet clinically indicated | 23-01-2020 |
| | | | Warning: To be sold by retail on the prescription of Oncologist only. | |
| 6 | Canagliflozin 100 mg and 300 mg Tablet (Additional Indication) | Each film coated tablet contains: Canagliflozin 100mg/300mg | To reduce the risk of major adverse cardiovascular events (cardiovascular death, nonfatal myocardial infarction and nonfatal stroke) in adults with type 2 diabetes mellitus and established cardiovascular disease. | 17-02-2020 |
| 7 | Norethisterone Acetate Controlled Release Tablet 10 mg (Additional Dosage form & Indication) | Norethisterone Acetate 10mg | To arrest bleeding in women with dysfunctional uterine bleeding due to hormonal imbalance. | 24-02-2020 |
| 8 | Lidocaine Hydrochloride 2% Solution (Additional Dosage Form) | Each ml contains: Lidocaine Hydrochloride 20mg | Indicated in adults: • For the production of topical anesthesia of irritated or inflamed mucous membranes of the mouth and pharynx. • For reducing gagging during the taking of X-ray pictures and dental impressions. | 25-02-2020 |

| 9 | Saroglitazar Tablets 4mg | Each uncoated tablet contains: | Noncirrhoic Non-Alcohol Steatohepatitis | 02-03-2020 |
|----|---|--|--|------------|
| | (Additional Indication) | Saroglitazar 4 mg | | |
| 10 | Itraconazole Capsule 200 mg (Additonal Strength) | Each capsule contains: Itraconazole 200 mg | For the treatment of onychymycosis of the toenail due to Trichophyton rubrum or T. mentagrophytes in non-immunocompromised patients | 03-03-2020 |
| | | | Warning: To be sold by retail on the prescription of a Specialist only. | |
| 11 | Nepafenac Ophthalmic suspension 0.3% w/v | Each ml contains: Nepafenac 3mg | Reeducation in the risk of postoperative macular oedema associated with cataract surgery in diabetic patients | 03-03-2020 |
| | (Additional Indication) | | Warning: To be sold by retail on the prescription of a Opthamologists only. | |
| 12 | Bisoprolol Fumarate tablet 1.25 mg (Additional strength) | Each film coated tablet contains: Biosoprolol Fumarate 1.25 mg | For the treatment of congestive heart failure (CHF) | 04-03-2020 |
| 13 | Pirfenidone Tablets 267 & 801 mg (Additional Strength) | Each film coated tablet contains: Pirfenidone 267mg/801 mg | For the treatment of idiopathic pulmonary fibrosis Warning: To be sold by retail on the prescription of a Pulmonologist only. | 04-03-2020 |
| 14 | Pegaspargase Injection 3750 IU/5ml (Additional Indication) | Each 5ml vial contains: Pegaspargase Injection 3750 IU | Injection as first-Line chemotherapeutic agent for treatment of patients with acute Lymphoblastic Leukemia (ALL) Warning: To be sold by retail on the prescription of a | 04-03-2020 |
| 15 | Controlled Release Tablet 15 mg | | Oncologist only. To arrest bleeding in women with Heavy Menstrual Bleeding (HMB)/abnormal Uterine Bleeding (AUB) due to hormonal imbalance | 09-03-2020 |
| | (Additional Strength) | | Warning: To be sold by retail on the prescription of a Gynaecologist/Obstertrician only. | |
| 16 | Estriol Tablets 1mg/2mg (Additional strength) | Each uncoated Tablet contains: Estriol 1mg/2mg | Estrogen deficiency symptoms due to menopause Warning: To be sold by retail on the prescription of a Gynaecologist only. | 09-03-2020 |
| 17 | Busulfan Injection 60 mg/10ml (Additional Strength) | Each 10 ml vial contains: Buslfan 60mg | In combination with cyclophosphamide as a conditioning regimen prior to allogenic hematopoietic progenitor cell transplantation for chronic myelogenous leukemia | 09-03-2020 |
| | | | Warning: To be sold by retail on the prescription of a Oncologist only. | |
| 18 | Ulipristal Acetate Tablet 30 mg | Each film coated tablet contains Ulipristal acetate 30 mg | Emergency contraception within 120 hrs (5 days) of unprotected sexual intercourse or contraceptive failure | 11-03-2020 |
| | (Additional strength& indication) | | Warning: To be sold by retail on the prescription of a Gynaecologist only. | |
| 19 | Canagliflozin Tablet 100/300 mg (Additional Indication) | Each film coated tablet contains: Canagliflozin 100/300 mg | To reduce the risk of end-stage kidney disease (ESKD), doubling of serum creathinine, cardiovascular (CV) death, and hospitalization for heart failure in adults with type 2 diabetes mellitus and diabetic nephropathy with albuminuria 300mg/day | 11-03-2020 |
| 20 | Nintedanib Soft Gelatin Capsules 100/150 mg (Additional Indication) | _ | For the treatment of systemic sclerosis associated interstitial Lung Disease (SSc-ILD) Warning: To be sold by retail on the prescription of a Pulmonologist only. | 20-03-2020 |
| | l | mg | 1 samonorogios om j. | |

| 21 | Oxaliplatin Injection 50 mg/vial & 100 mg/val | Each Vial contains: Sterile Solution of Oxaliplatin | For the treatment of advanced colorectal cancer. Warning: To be sold by retail on the prescription of an | 29-04-2020 |
|----|---|--|--|------------|
| | (Additional Dosage Form) | 50mg and 100mg | oncologist only. | |
| 22 | Prolonged Release Tablets | _ | For the treatment of Polycystic Ovary Syndrome (PCOS) with established insulin resistance | 05-05-2020 |
| | 500mg (Additional Indication) | Metformin Hydrochloride 500mg | WARNING: To be sold by retail on the prescription of a RMP only | |
| | | | In the package insert - | |
| | | | Special warnings and precautions: Pregnancy: | |
| | | | "There are no adequate and well-controlled | |
| | | | studies in pregnant women with Metformin and Metformin should not be used during pregnancy unless clearly needed" | |
| 23 | Olaparib 100 mg and 150 mg Tablets (Additional Indication) | Each Film coated tablet contains: Oloaparib100mg/150 mg. | Olaparib is indicated as monotherapy for the Maintenance treatment of adult patients with germline BRCA-mutated metastatic adenocarcinoma of the pancreas whose disease has not progressed on first-line platinum-based chemotherapy "WARNING: To be sold by retail on the prescription of Oncologist only" | 06-05-2020 |
| 24 | | Each Vial contains: | For the treatment of patients with multiple myeloma | 12-05-2020 |
| | 2.5mg/ml (Additional Pack Size) | Bortezomib 2.5 mg | • For the treatment of patients with mantle cell lymphoma who have received at least one prior therapy. | |
| | | | WARNING: To be sold by retail on the prescription of Oncologist only" which shall be in red box; | |
| 25 | (50mg/ml) (Dental | Each ml of Dental Suspension contains: | • For the prevention of caries in children and adults as part of a comprehensive control programme, for the: | 05-06-2020 |
| | Suspension) | Sodium Fluoride50 mg | - prevention of recurring (or marginal) caries | |
| | (New Dosage Form) | | - prevention of progression of caries | |
| | | | - prevention of decalcification around orthodontic appliances | |
| | | | - prevention of pit and fissure (occlusal) caries. | |
| | | | • For the desensitisation of hypersensitive teeth as part of a treatment regimen which includes the daily use of a suitable toothpaste. | |
| | | | Warning: "To be sold by retail on the prescription of a Dentist only" | |
| 25 | Paclitaxel Lipid Suspension for injection 100 mg/vial (Additional Strength) | Each vial contains: Paclitaxel100 mg | 1. For the treatment of breast cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated. | 16-06-2020 |
| | | | 2. For the adjuvant treatment of node-positive breast cancer administered sequentially to standard doxorubicin-containing combination chemotherapy. | |
| | | | 3. As first-line and subsequent therapy for the treatment of advanced carcinoma of the ovary. As first-line therapy, it is indicated in combination with cisplatin. | |
| | | | 4. In combination with cisplatin, is indicated for the first-line treatment of non-small cell lung cancer in patients who are | |

| | | | not candidates for potentially curative surgery and/or radiation therapy. | |
|----|---|---|---|------------|
| | | | 5. For the second-line treatment of AIDS- related Kaposi's sarcoma | |
| | | | Warning: "To be sold by retail on the prescription of a Oncologist only" | |
| 26 | Itraconazole Oral Solution 10 mg/ml | Each ml contains: | For the Treatment of Orophyaryngeal and Esophageal candidiasis | 25-06-2020 |
| | (New Dosage Form) | Itraconazole10 mg | Warning: "To be sold by retail on the prescription of a Specialist only" | |
| 27 | Bilastine Orodispersible Tablet 10 mg (New Dosage Form) | Each uncoated orodispersible tablet contains: Bilastine 10 mg | For symptomatic treatment of allergic rhino-conjunctivitis (seasonal and perennial) and urticarial. Bilastine is indicated in children aged 6 to 11 years with a body weight of at least 20 kg | 25-06-2020 |
| 28 | Itraconazole 50 mg Capsule (Modified Dosage Form) | Each capsule contains: Itraconazole50 mg | for the treatment of Systematic aspergillosis, candidiasis, cryptococcosis, sportrichosis, paracoccidioidomycossis, blastomycosis and other rarely occuring systemic or tropical mycosis "WARNING: To be sold by retail on the prescription of a | 30-06-2020 |
| | | | Specialist only". Caution: "This Itraconazole Capsules 50mg has equivalent bioavailability with that of itraconazole capsules 100mg. One capsule of this Itraconazole Capsule 50mg is therapeutically equivalent to one 100mg capsule of conventional Itraconazole capsules100mg. The recommended dose of this Itraconazole Capsules 50mg is therefore half the recommended dose for conventional itraconazole capsules 100 mg. This Itraconazole Capsules 50mg and conventional Itraconazole Capsules 100mg are not interchangeable". | |
| 29 | Meropenem dry powder for injection 2000 mg | Meropenem Trihydrate | Urinary Tract Infections like Complicated Urinary Tract Infections | 02-07-2020 |
| 30 | Dapagliflozin 10 mg | Each Film coated tablet contains: Dapagliflozin Propanediol Monohydrate equivalent to Dapagloflozin10 mg | 6. Dapagliflozin is indicated in adults for the treatment of heart failure with reduced ejection fraction. | 03-07-2020 |
| 31 | Ceftaroline fosamil powder for concentrate for solutin for infusion | Each Vial contains: Ceftaroline Fosamil600mg L-Arginine395mg | For the tratment of the adults with complicated skin and soft tissue infections (Cssti) | 06-07-2020 |

| 32 | Nintedanib 100mg/150mg | Each soft Gelatin capsule contains: Nintedanib Ethanesulfonate equivalent to Nintedanib100mg/150mg | For the treatment of other chronic fibrosing interstitial Lung Diseases (ILDs) with a Progressive phenotype | 09-07-2020 |
|----|---|--|--|------------|
| 33 | Ivabradine PR Tablets 10/15mg | Each film coated prolonged release tablet contains: Ivabradine hydrochloride equivalent to Ivabradine | For the treatment of chronic stable angina pectoris in patients with coronary artery disease and treatment of chronic heart failure. | 14-07-2020 |
| 34 | Paracetamol Orally Disintegrating 60/120 mg | Each Orally Disintegrating Strip contains: Paracetamol60mg/120 mg | For use in pediatric age group, for symptomatic treatment of pain and fever | 14-07-2020 |
| 35 | Sodium Citrate Granules 4.0 gm | Each sachet (5.6g) contains: Sodium Citrate 4.0 g | As an alkalising agent in cystitis in women | 16-07-2020 |
| 36 | Bedaquiline Tablet 100 mg | Each uncoated tablet contains: Bedaquiline fumarate equivalent to | In adult (>18 years) and pediatric patients (12 years to less than 18 years of age and weighing at least 30 kg) as part of combination therapy of pulmonary tuberculosis (TB) due to multi-drug resistant Mycobacterium tuberculosis. | 16-07-2020 |
| 37 | Trimetazidine Hydrochloride Extended Release Capsules 80mg | | For the treatment of Ischaemic heart disease (angina pectoris, sequelae of infarction). | 21-07-2020 |
| 38 | Regorafenib Tablet 40 mg (Additional Indication) | Each film coated tablet contains: Regorafenib40 mg | For the treatment of patients with hepatocellular carcinoma (HCC) who have been previously treated with Sorafenib Warning: "To be sold by retail on the Prescription of an Oncologist only | 03-08-2020 |
| 39 | Baclofen Extended Release tablet 30mg (Additional dosage & Strength) | Each uncoated extended release tablet contains:Baclofen30 mg | For the symptomatic treatment of neuronal spasticity due to multiple sclerosis, spinal chord, pathology & injury | 04-08-2020 |
| 40 | Tofacitinib Tablet 5 mg (Additional Indication) | Each film coated tablet contains: Tofacitinib citrate equivalent to tofacitinib5 mg | For the treatment of adult patients with active psoriatic arthritis (PsA) who have had an inadequate response or intolerance to methotrexate or other non-biologic disease modifying antirheumatic drugs (DMARDs) Warning: "To be the prescription of a Rheumatologist or | 04-08-2020 |
| 41 | Enzalutamide 80 mg capsules (Additional strength) | Each hard gelatin capsule contains: Enzalutamide80mg | Specialist in Internal Medicine only" For the treatment adults with metastatic castration-resistant prostate cancer (mCRPC) whose disease has progressed on or after Docetaxel therapy. Warning: "To be sold by retail on the Prescription of an Oncologist only" | 06-08-2020 |
| 42 | Oxaceprol Sustained Release Tablet 600 mg (Additional strength) | Each film coated SR tablet contains: Oxaceprol600 mg | Treatment of pain associated with osteoarthritis of knee in adult patients | 07-08-2020 |

| 43 | Olaparib 100/150 mg | Each film-coated tablets contains: | Olaparib is indicated as monotherapy for the treatment of | 17-08-2020 |
|----|---|---|--|------------|
| | (Additional Indication) | Olaparib100mg/15 0 mg | adult patients with metastatic castration-resistant prostate cancer and homologous recombination repair gene mutations (germline and/or somatic) who have progressed following a prior new hormonal agent. | |
| | | | Warning: "To be sold by retail on the Prescription of an Oncologist only" | |
| 44 | Levofloxacin Dispersible Tablets 100 mg (New Dosage Foam) | Each dispersible tablet contains: Levofloxacin hemihydrate Equivalent to Levofloxacin | Indicated in combination with other anti-tuberculosis agents, for the treatment of tuberculosis caused by Mycobacterium tuberculosis in children. | 18-08-2020 |
| | | 100 mg | Warning: "To be sold by the prescription of a Medical Specialist only" | |
| 45 | Trihexyphenidyl Hydrochloride 1 mg (Additional strength) | Each uncoated tablet contains: Trihexyphenidyl hydrochloride1 mg | It is indicated in all forms of Parkinsonism (postencephalitic, arteriosclerotic and idiopathic). It is often useful as adjuvant therapy when treating these forms of Parkinsonism with levodopa. Trihexyphenidyl is also indicated to control extrapyramidal disorders due to central nervous system drugs such as reserpine and the phenothiazines. Warning: "To be sold by the prescription of Neurologist | 18-08-2020 |
| | | | only" | |
| 46 | Olaparib 100/150 mg (Additional Indication) | Each film-coated tablets contains: Olaparib100mg/1 50 mg | Olaparib in combination with bevacizumab is indicated for the Maintenance treatment of adult patients with advanced high-grade epithelial ovarian, fallopian tube or primary peritoneal cancer who are in response (Complete response or partial response) to first-line platinum-based chemotherapy with bevacizumab. | 21-08-2020 |
| | | | Warning: "To be sold by retail on the Prescription of an Oncologist only" | |
| 47 | Ferric Carboxymaltose Solution for Injection/Infusion 20mg/vial (Additional Pack Size) | Each ml contains: Ferric Carboxymaltose equivalent to Elemental Iron50 mg. | For the treatment of iron deficiency in adults when oral iron preparations are ineffective or cannot be used. | 27-08-2020 |
| 48 | Lipid Based Tacrolimus Ointment 0.1% w/w (Additional indication) | Each gram contains: Tacrolimus Monohydrate Equivalent to Tacrolimus0.1% w/w | For the short-term and non-continuous chronic treatment of symptomatic lichen planus (oral and mucosal forms) in non-immunocompromised patients Warning: "To be the prescription of a Dermatologists only" | 28-08-2020 |
| 49 | Moxifloxacin Dispersible Tablets 100 mg (New Formdosage) | Each dispersible tablet contains: Moxifloxacin Hydrochloride Equivalent to Moxifloxacin 100 mg | Indicated in combination with other antituberculosis agents for the treatment of tuberculosis caused by Mycobacterium tuberculosis. as a second-line anti-mycobacterial drug when use of first line drugs is not appropriate due to resistance or intolerance | 02-09-2020 |
| 50 | Tapentadol Hydrochloride Nasal Spray 225 mg/ml (New Form dosage) | Each Spray (0.1 ml) contains: Tapentadol Hydrochloride22.5 mg Benzalkonium chloride (50%) 0.02%w/v (As Preservative) | For treatment of moderated to severe post-operative pain in hospital admitted patients | 02-09-2020 |
| 51 | Favipiravir 800mg Tablet (Additional strength) | Each film coated tablet contains: Favipiravir 800 mg | For the treatment of patients with mild to moderate Covid-19 disease, in light of Covid 19 outbreak, for restricted emergency use in the country. "WARNING: To be sold on the prescription of medical specialist" | 24-09-2020 |
| | | | SP | |

| 52 | Isotretinoin Capsules 8/16/24 mg (Additional Strength & Indication) | Each soft gelatin capsule contains: Isotretinoin8mg/16mg/24mg | For the treatment of severe recalcitrant nodular acne in non-pregnant patients 12 of age and older with multiple inflammatory nodules with a diameter of 5 mm or greater. Because of significant adverse reactions associated with its use, it is reserved for patients with severe nodular acne who are unresponsive to conventional therapy, including systemic antibiotics. | 29-09-2020 |
|----|--|--|--|------------|
| | | | "Warning: To be sold by retail on the prescription of Dermatologist only" | |
| 53 | Brivaracetam Tablets (Additional Strength) | Each film coated tablet contains: Brivaracetam 25mg | As adjunctive therapy in the treatment of partial-onset seizures in patients 16 years of age and older with epilepsy | 01-10-2020 |
| 54 | Perampanel Tablets 2 mg/4mg/6 mg/8 mg/12 mg (Additional Indication) | Each film coated tablet contains: Perampanel 2mg/4mg/6mg/8mg/10 mg/12mg | As an adjunctive treatment of partial-onset seizures with or without secondarily generalized seizures in patients with epilepsy 4 years of age and older "Warning: To be sold by retail on the prescription of Neurologist only" | 01-10-2020 |
| 55 | Acotiamide 300 mg Tablet (Additional dosage foam) | Each film coated Extended release tablet contains: Acotiamide Hydrochloride Hydrate 300mg | For the treatment of bloating after meals, epigastric bloating and early satiety in functional dyspepsia | 05-10-2020 |
| 56 | Azathioprine 75 mg Tablet (Additional Strength) | Each film coated tablet contains: Azathioprine 75 mg | Indicated for facilitating the survival and function of organ transplants "Warning: To be sold by retail on the prescription of Medical Specialist only" | 14-10-2020 |
| 57 | Travoprost Eye Drops 0.004% w/v (Preserved with an ionic buffer system) | Each ml contains Travoprost .0.004% w/v | For the reduction of elevated intraocular pressure in patients with open angle glaucoma or ocular hypertension "Warning: To be sold by retail on the prescription of Ophthalmologist only" | 16-10-2020 |
| 58 | Remdesivir for Injection 100 mg/vial (lyophilized) (Expansion of Indication) | Each vial contains: Remdesivir 100 mg (lyophilized) | For treatment of suspected or laboratory confirmed corona virus disease 2019 (COVID-19) in adults and children hospitalised with moderate to severe disease. In light of Covid 19 outbreak, for restricted emergency use in the country. "Warning: To be sold by retail on the prescription of specialist for use in Hospital/Institutional Setup only" | 19-10-2020 |
| 59 | Lidocaine Spray 10% w/v (Additional Strength) | Each ml contains: Lidocaine | Lidocaine Spray 10% w/v is indicated for the prevention of pain associated with the following procedures: Otorhinolaryngology: Anesthesia prior to minor non-invasive procedures in the nasal cavity, pharynx and epipharynx including rhinos copy and laryngoscopy. Obstetrics: During the final stages of delivery and before episiotomy and perineal suturing as supplementary pain control. Insertion of instruments and catheters into the respiratory and digestive tract: Provides surface anesthesia for the oropharyngeal and tracheal areas to reduce reflex activity, attenuate hemodynamic response and to facilitate insertion of the catheter or the passage of instruments during endotracheal intubation, laryngoscopy, bronchoscopy, esophagostomy and gastroscopy. Dental practice: Before minor dental procedures where local anesthesia is desired. | 02-11-2020 |

| 60 | Combipack OF Favipiravir Tablets (Combipack) | Each Combipack contains: Favipiravir Tablets 200 mg (2 tablets) Each film coated tablets contains: Favipiravir | For restricted emergency use for the treatment of patients with mild to moderate Covid-19 disease "WARNING: To be sold on the prescription of medical specialist" which shall be in box with red background. | 27-11-2020 |
|----|--|--|--|------------|
| 61 | Vildagliptin Sustained Release Tablets 100 mg (OROS) (Additional Strength) | Each film coated sustained release bi-layered tablet contains: Vildagliptin100mg | As an adjunct to diet and exercise to improve glycemic control in patients with type-II diabetes mellitus (T2DM) | 15-12-2020 |
| 62 | Thyroxine Sodium Tablets 112 mcg (Additional Strength) | Each uncoated tablet contains: Thyroxine Sodium | For the treatment of hypothyroidism | 17-12-2020 |
| 63 | Triamcinolone Acetonide Cream 0.1% w/w (New Dosage Foam) | Each gm contains: Triamcinolone Acetonide1mg Benzyl alcohol (as preservative)2% w/w In a creambaseq.s. | Inflammatory and pruritic manifestations of corticosteroid responsive dermatoses "WARNING: To be sold by retail on the prescription of Dermatologist only" | 22-12-2020 |
| 64 | Palbociclib Tablets 75/100/125 mg | Each film coated tablet contains: Palbociclib | 1. Palbociclib is a kinase inhibitor indicated for the treatment of hormone receptor (HR) positive, human epidermal growth factor receptor 2 (HER2) - negative advanced or metastatic breast cancer in combination with fulvestrant in women with disease progression following endocrine therapy. 2. Palbociclib is a kinase inhibitor indicated in combination with Letrozole for the treatment of postmenopausal women with estrogen receptor (ER) positive, human epidermal growth factor receptor 2 (HER2)-negative advanced breast cancer as initial endocrine based therapy for their metastatic disease. "WARNING: To be sold by retail on the prescription of an Oncologist only" | 22-12-2020 |
| 65 | Lurasidone Hydrochloride Tablets 20mg (Additional Strength) | Each film coated tablets contains: Lurasidone Hydrochloride | For treatment of patients with Schizophrenia "WARNING: To be sold by retail on the prescription Neurologist only" | 24-12-2020 |
| 66 | Hydrocortisone Sodium Succinate Injection (Additional Strength) | Each vial contains: Hydrocortisone Sodium Succinate (Buffered) Eq. to Hydrocortisone | Allergic states: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, perennial or seasonal allergic rhinitis, serum sickness, transfusion reactions. Dermatologic diseases: Bullous dermatitis herpetiformis, exfoliative erythroderma, mycosis fungoides, pemphigus, severe erythema multiforme (Stevens-Johnson syndrome). Endocrine disorders: Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; synthetic analogs may be used in conjunction with mineralocorticoids where applicable; in infancy, | 28-12-2020 |

mineralocorticoid supplementation is of particular importance), congenital adrenal hyperplasia, hyper calcemia associated with cancer, nonsuppurative thyroiditis. Gastrointestinal diseases: To tide the patient over a critical period of the disease in regional enteritis (systemic therapy) and ulcerative colitis. Hematologic disorders: Acquired (autoimmune) hemolytic anemia, congenital (erythroid) hypoplastic anemia (Diamond-Blackfan anemia), idiopathic thrombocytopenic purpura in adults (intravenous administration only; intramuscular administration is contraindicated), pure red cell aplasia, selected cases of secondary thrombocytopenia. Miscellaneous: Trichinosis with neurologic or myocardial involvement, tuberculous meningitis with subarachnoid block or impending block when used concurrently with appropriate antituberculous chemotherapy. Neoplastic diseases: For the palliative management of leukemias and lymphomas. Nervous System: Acute exacerbations of multiple sclerosis; cerebral edema associated with primary or metastatic brain tumor, or craniotomy. Ophthalmic diseases: Sympathetic ophthalmia, uveitis and ocular inflammatory conditions unresponsive to topical corticosteroids. Renal diseases: To induce diuresis or remission of proteinuria in idiopathic nephritic syndrome or that due to lupus erythematosus. Respiratory diseases: Berylliosis, fulminating or disseminated pulmonary tuberculosis when used concurrently with appropriate antituberculous chemotherapy, idiopathic eosinophilic pneumonias, symptomatic sarcoidosis. Rheumatic disorders: As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in acute gouty arthritis; acute rheumatic carditis; ankylosing spondylitis; psoriatic arthritis; rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy). For the treatment of dermatomyositis, temporal arteritis, polymyositis, and systemic lupus erythematosus Saroglitazar Tablets 4mg Each uncoated tablet contains: Non-Alcoholic Fatty Liver Disease (NAFLD) with 28-12-2020 67 comorbidities (Either Obesity, Type 2 Diabetes Mellitus, (Additional Indication) Saroglitazar 4 mg. Dyslipidemia Or Metabolic Syndrome)

Website Update 01-01-2021 to 31-07-2021

| Sl. No. | Drug Name | Composition | Indication | Date of Approval |
|---------|--|--|--|------------------|
| 1 | Atropine Sulfate Ophthalmic Solution USP 0.01% w/v (Additional Strength) | Each ml contains: Atropine Sulfate | It is indicated to control the progression of myopia in children of 5 years and above "WARNING: To be sold by retail on the prescription of Ophthalmologist only" With the Condition to conduct phase IV Clinical trial to assess long term safety of the product. | 12-01-2021 |
| 2 | Colistimethate Sodium For Injection IP 3 MIU/ 4.5 MIU | (i) Colistimethate Sodium for Injection IP 3 MIU | For the treatment of some serious infections caused by Gram-negative bacteria, including those of the lower respiratory tract and urinary tract, when more commonly | 15-01-2021 |

| | (Additional Strength) | Each Vial contains Colistimethate Sodium3 000000 IU | used systemic antibacterial agents may be contraindicated or may be ineffective because of bacterial resistance. | |
|----|--|---|---|------------|
| | | (ii) Colistimethate sodium for Injection IP 4.5 MIU | "WARNING: To be sold by retail on the prescription of a Medical Specialist only" | |
| | | Each Vial contains: | | |
| | | Colistimethate Sodium45 00000 IU | | |
| 3 | Dapagliflozin Tablets 10 mg Additional Indication | Each film-coated tablets contains: Dapagliflozin propanediol monohydrate equivalent to Dapagliflozin 10 mg | For the treatment of patients of Chronic Kidney Disease (CKD) up to Stage III (eGFR of greater than or equal to 30ml/min/1.73m2) "WARNING: To be sold by retail on the prescription of Nephrologist only" | 03-02-2021 |
| 4 | Brivaracetam Oral Solution Solution 10mg/ml Additional Dosage Foam | Each ml contains: Brivaracetam10mg | As adjunctive therapy in the treatment of partial onset seizures in patients 16 years of age and older with epilepsy | 03-02-2021 |
| 5 | Drospirenone Tablets 4 mg Additional Strength | Each for uncoated Tablets Drospirenone4 mg | Drospirenone is a Progestin intended for its use by females of reproductive potentials to prevent pregnancy | 15-02-2021 |
| 6 | in 20ml vial Apomorphine Hydrochloride 10 mg/ml solution for injection in 3ml cartridge | solution for infusion in 20ml vial Each ml contains: Apomorphine Hydrochloride | For the treatment of disabling motor fluctuations ("on-off" phenomena) in patients with Parkinson's disease which persist individually titrated treatment with Levodopa (with a peripheral decarboxylase inhibitor) and/or other dopamine agonists. "WARNING: To be sold by retail on the prescription of Neurologists only" | 16-02-2021 |
| 7 | Thyroxine Sodium Tablets 88 mcg/200mcg Additional Strength | Each uncoated tablet contains: Thyroxine Sodium | For the treatment of hypothyroidism | 22-02-2021 |
| 8 | Brivaracetam injection 50mg/5ml (10mg/ml) Additional dosage form & New route of administration | Each ml contains: Brivaracetam10mg | As adjunctive therapy in the treatment of partial-onset seizures in patients 16 years of age and older with epilepsy | 24-02-2021 |
| 9 | Aripiprazole powder and solvent for prolong-release suspension for injection 300mg/vial and 400 mg/vial (Modified Dosage Form) | Aripiprazole 300mg / | Maintenance treatment of schizophrenia in adult patients stabilized with oral aripiprazole "WARNING: To be sold by retail on the prescription of an Psychiatrist only" | 01-03-2021 |
| 10 | Rebamipide Ophthalmic Suspension 2% w/v with Stabilized Oxychloro complex (as preservative) | Each ml contains: Rebamipide%w/v Stabilized Oxychloro Complex0.01% w/v (As Preservative) | Indicated for the treatment of dry eye signs and symptoms | 01-03-2021 |
| 11 | Pancreatin Tablets USP 200 mg (Additional Strength/New dosage form/New Indication) | Each uncoated tablet contains: Pancreatin200mg Equivalent to Lipase 10,440 USP units Amylase 39,150 USP units | Indicated in combination with a proton pump inhibitor in adults for the treatment of exocrine pancreatic insufficiency due to chronic pancreatitis or pancreatectomy | 01-03-2021 |

| | | Protease39150 USP units | | |
|----|--|--|--|------------|
| 12 | Pemetrexed for injection 100mg/500mg (Additional Indication) | Each vial contains: Pemetrexed Disodium equivalent to Pemetrexed100mg/500mg | In combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous NSCLC, with no EGFR or ALK genomic tumor aberrations "WARNING: To be sold by retail on the prescription of oncologist only" | 08-03-2021 |
| 13 | Ibuprofen solution for infusion 400mg/100ml (New Dosage Form) | Each ml contains: Ibuprofen4mg | Indicated in adults for the short-term symptomatic treatment of acute moderate pain, and for the short-term symptomatic treatment of fever, when administration by intravenous route is clinically justified when other routes of administration are not possible | 08-03-2021 |
| 14 | Cholecalciferol oral drops 800IU/ml (Additional Strength/New dosage Form) | Each ml contains: Cholecalciferol 800IU | For the treatment of vitamin D3 deficiency | 08-03-2021 |
| 15 | Osimertinib Tablets 40mg and 80mg (Additional Indication) | Each film-coated tablets contains: Osimertinib (as mesylate) 40 mg/80mg | Osimertinib as monotherapy is indicated for: The adjuvant treatment after complete tumour resection in patients with nonsmall cell lung cancer (NSCLC) whose tumours have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations "WARNING: To be sold by retail on the prescription of oncologist only" | 09-03-2021 |
| 16 | Dolutegravir Dispersible Tablets 10mg (Additional Dosage Form) | Each Tablet contains: Dolutegravir sodium equivalent to Dolutegravir 10mg | In combination with other antiretroviral agents for the treatment of Human Immunodeficiency Virus type1 (HIV-1) infection in adults (treatment-naive or -experienced) and in pediatric patients (treatment-naive or -experienced but integrase strand transfer inhibitor [INSTI]-naive) aged at least 4 weeks and weighing at least 3 kg "WARNING: To be sold by retail on the prescription of a | 23-03-2021 |
| 17 | 12.5% w/v | Each Spray (0.2ml) delivers Sildenafil citrate equivalent to Sildenafil25 mg | Specialist only" For the treatment of Erectile Dysfunction in men "WARNING: To be sold by retail on the prescription of an | 26-03-2021 |
| 18 | (Additional Dosage Form) Dexamethasone Tablets 6 mg (Additional Strength) | Each film coated tablet contains: Dexamethasone6 mg | Urologist, Psychiatrist and Endocrinologist only" Dexamethasone is indicated for treatment of variety of disorders in which systemic glucocorticoids are indicated including: • Neurology: Cerebral oedema (only with symptoms of intracranial pressure evidenced by computerised tomography) caused by a primary or metastatic brain tumour, craniotomy or head injury. • Dermatology: Initial treatment of extensive, severe, acute, skin diseases responding to glucocorticoids, e.g. erythroderma, pemphigus vulgaris • Autoimmune disorders/rheumatology: Initial treatment of autoimmune disorders like systemic lupus erythematodes, in acute gouty arthritis ,acute arthritis carditis, ankylosing spondylitis ,psoriatic arthritis , Severe progressive course of active rheumatoid arthritis Severe systemic course of | 06-04-2021 |
| | | | juvenile idiopathic arthritis (Still's disease). • Haematological disorder: Idiopathic thrombocytopenic | |

| | | | purpura in adults. | |
|----|--|---|--|------------|
| | | | Oncologists only- for steroid responsive neoplastic disease like leukemias and lymphomas in adults only. | |
| 19 | Betahistine Hydrochloride Orally Disintegrating tablet 8 mg/16mg/24 mg (New Dosage Form) | Each Uncoated Orally Disintegrating Tablet contains: Betahistine Hydrochloride mg/16mg/24mg Excipients q.s. | Indicated for the symptomatic treatment of vertigo caused due to conditions including Menier's syndrome. | 06-04-2021 |
| 20 | Pegaspargase powder for solution for injection/ infusion (New Dosage Form & Indication) | | Indicated as a component of anti-neoplastic combination therapy in acute lymphoblastic leukaemia (ALL) in paediatric patients from birth to 18 years, and adult patients | 09-04-2201 |
| 21 | Drospirenone Tablets 4mg (New Dosage Form) | Each Film Coated Tablet contains: Drospirenone4mg | Drospirenone is a Progestin intended for its use by females of reproductive potentials to prevent pregnancy | 22-04-2021 |
| 22 | Baricitinib Tablets 1mg, 2mg, & 4 mg (Additional Strength & Additional Indication) | Each Film Coated Tablet contains: Baricitinib1mg/2mg/4 mg | Baricitinib, in combination with Remdesivir, for treatment of suspected or laboratory confirmed coronavirus disease 2019 (COVID-19) in hospitalized adults requiring supplemental oxygen, invasive mechanical ventilation, or extracorporeal membrane oxygenation (ECMO) "WARNING: To be sold by retail on the prescription of | 01-05-2021 |
| 23 | Remdesivir Injection 100mg/20 ml (without Cyclodextrin) | Each ml contains: Remdesivir5mg | specialist for use in Hospital/ Institutional set up only" For treatment of suspected or laboratory confirmed corona virus disease 2019 (COVID-19) in adults and children hospitalized with moderate to severe disease "WARNING: To be sold by retail on the prescription of specialist for use in Hospital/ Institutional set up only" | 06-05-2201 |
| 24 | Favipiravir Oral suspension 100mg/ml (New Dosage Form) | Each ml contains: Favipiravir 100mg | For the treatment of patients with mild to moderate Covid- 19 disease "Warning: To be sold on prescription of medical specialist" | 19-05-2021 |
| 25 | Posaconazole Delayed- Release Tablets (Additional Indication) | Each Delayed-Release Tablets contains: Posaconazole100mg | For the treatment of oropharyngeal candidiasis, including oropharyngeal candidiasis refractory to Itraconazole and/or fluconazole for prophylaxis of invasive Aspergillus and Candida infections in patients, 13 years of age and older, who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy. Zygomycosis (Mucormycosis) in patients with diseases refractory to other therapy, or or patients who are intolerant | 28-05-2021 |
| | | | to other therapy "Warning: To be sold on prescription of medical specialist" | |
| 26 | Posaconazole Gastro-Resistant Tablets 300 mg (Additional Strength) | Each Gastro-Resistant Tablet contains: Posaconazole3 00 mg | For prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with | 07-06-2021 |

| | T | <u> </u> | | |
|----|---|---|--|------------|
| | | | prolonged neutropenia from chemotherapy. | |
| | | | Posaconazole Gastro Resistant Tablets are indicated in patients 13 years of age and older. | |
| | | | "WARNING: To be sold by retail on the prescription of a Medical Specialist only" | |
| 27 | | Each vial contains: | • For the treatment of severe sepsis. | 24-06-2021 |
| | 200000 IU (Liquid) (Additional Strength) | Ulinastatin200000 IU | For the treatment of mild and severe acute pancreatitis. | |
| 28 | Imatinib Oral Solution 80 mg/ml (New Dosage Foam) | Each ml contains: Imatinib (As mesilate) 80mg | 1. Treatment of adult and pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) integrated with chemotherapy. | 24-06-2021 |
| | | | 2. Treatment of adult patients with relapsed or refractory Ph+ ALL as monotherapy. | |
| | | | 3. Adjuvant treatment of adult patients following resection of gastro-intestinal stromal tumors (GIST), and 4. Treatment of CML in blast crisis, accelerated phase or in | |
| | | | chronic phase after failure of interferon alpha therapy. "WARNING: To be sold by retail on the prescription of oncologist only" | |
| 29 | Clozapine Tablet 12.5 mg | Each uncoated tablet contains: | Indicated in the management of schizophrenic patients | 29-06-2021 |
| | (Additional Strength) | Clozapine 12.5 mg | "WARNING: To be sold by retail on the prescription of Neurologist only" | |
| 30 | Sorafenib Tablets 400mg (Additional Strength) | Each film coated tablets contains: Sorafenib Tosylate eq. to Sorafenib400 mg | For the treatment of hepatocellular carcinoma, advanced renal cell carcinoma and locally advance or metastatic differentiated thyroid carcinoma refractory to radioactive iodine | 30-06-2021 |
| | | | "WARNING: To be sold by retail on the prescription of Oncologist only" | |
| 31 | Favipiravir Powder for Oral suspension 800mg | Each sachet contains: | For the treatment of patients with mild to moderate Covid- 19 disease | 06-07-2021 |
| | (Additional Dosage Form) | 1.5g of Favipiravir powder contains: | "WARNING: To be sold on the prescription of medical specialist" | |
| | | Favipiravir 8 00mg Excipients q.s | | |
| 32 | | Each ml contains: | For Prevention of infusion associated phlebitis | 09-07-2021 |
| | Solution 1000 IU/ml (Additional Indication) | Heparin Sodium1000 IU/ml AbsoluteAlcohol10.00%v/v Topical Solution baseq.s. | "WARNING: To be sold by retail on the prescription of Specialists only" | |
| 33 | Pemetrexed for injection 1000mg (Lyophilized) (Additional Strength) | Each vial contains: Pemetrexed Disodium heptahydrate IP equivalent to Pemetrexed1000mg | 1. Indicated as a monotherapy for the maintenance treatment of locally advanced or metastatic Non-Small Cell Lung Cancer (NSCLC) other than predominantly squamous cell histology in patients whose disease has not progressed immediately following platinum-based chemotherapy. First line treatment Should be a platinum doublet with gemcitabine, paclitaxel or docetaxel. | 16-07-2021 |
| | | | 2. In combination with cisplatin is indicated for the first line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly | |

| | | T | | 1 |
|-----|--|---|---|------------|
| | | | squamous cell histology. 3. As monotherapy for the second-line treatment of patients | |
| | | | with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology. | |
| | | | 4. As monotherapy for the maintenance treatment of locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology in patients whose disease has to progressed immediately following platinum based chemotherapy. "WARNING: To be sold by retail on the prescription of | |
| 2.4 | A . 10 1' 1' A '1 | E 1E66 | oncologist only' | 16.07.2021 |
| 34 | Acetyl Salicylic Acid Effervescent Tablets 500 mg | Each Effervescent tablet contains: Acetyl Salicylic Acid500 mg | Analgesic & Anti pyretic | 16-07-2021 |
| | (Additional Dosage Foam) | Acetyl Salicylic Acid 500 mg | | |
| 35 | Pentosan polysulfate sodium sterile solution for intravesical instillation 10 mg/ml (30ml) (Additional Dosage form & Route of administration) | | For intravesical treatment of the bladder pain syndrome/interstitial cystitis "WARNING: To be sold by retail on the prescription of a Specialist only" | 16-07-2021 |
| 36 | Amikacin Injection 750mg/3ml & 1000 mg/4ml | Amikacin Injection IP 750 mg Each 3ml contains: | Amikacin Injection is indicated in the treatment of serious infections due to Amikacin sensitive organisms | 23-07-2021 |
| | (Additional pack size /Additional Strength) | Amikacin Sulphate IP Equivalent to Amikacin | | |
| | | Amikacin Sulphate IP Equivalent to Amikacin1000 mg | | |
| | | Water for Injections IP q.s Methylparaben IP0.04w/v Propylparaben IP0.01w/v (As preservatives) (Suitable buffering agent and antioxidant added) | | |
| 37 | Riboflavin Sodium Phosphate IP eq. to Riboflavin 0.1 % w/v with Hypromellose Ophthalmic Solution in PFS (Pack Size is 3.0ml) (Additional Pack Size) | Each ml contains: Riboflavin Sodium Phosphate IP Eq. to Riboflavin0.1% w/v Hypromellose I.P%w/v | To treat keratoconus and corneal ectasia following refractive surgery | 27-07-2021 |
| 38 | Amorolfine Lotion 0.25% w/v (Additional Dosage Form) | Each ml contains: Amorolfine Hydrochloride Equivalent to Amorolfine0.2 5% w/v | For the treatment of dermatomycosis caused by dermatophytes, cutaneous candidiasis and pityriasis versicolor "WARNING: To be sold by retail on the prescription of an Dermatologist only" | 29-07-2021 |

| 39 | Silodosin Tablet 8 mg | Each film coated tablets contains: | For the treatment of signs and symptoms of benign prostatic | 30-07-2021 |
|----|--------------------------|------------------------------------|---|------------|
| | (Additional Dosage Form) | Silodosin8 mg | hyperplasia (BPH) in adults only | |
| | - | | "WARNING: To be sold by retail on the prescription of a | |
| | | | Urologist only" | |